

Perception of arable crop farmers on usage of organic fertilizer in maize production in Ido Local Government Area of Oyo State, Nigeria

Abstract

One major problem facing maize production can be attributed to the fertility status of most Nigerian soils which is considerably low. The study was conducted to examine arable crop farmers' perception of the use of organic fertilizer in maize production in Ido Local Government Area of Oyo State. A total of one hundred and twenty respondents were randomly selected for the study from 10 villages. Primary data were collected using a well-structured questionnaire and interview guide. Data were analyzed with descriptive statistics such as frequency, percentages and mean. Chi-square was used to test the hypothesis of the study. Poultry manure (90.0%) was the major source of organic fertilizer and constitutes the most frequently used (70.8%). It was found that bulkiness of organic fertilizer was the most severe constraint that affect its usage ($\bar{x}=2.43$). However, there was an unfavourable perception on use of organic fertilizer in maize production. Test of hypothesis revealed that age ($\chi^2= 4.46$, $df= 1$) and educational level ($\chi^2= 8.54$, $df = 3$) were found significantly related with the perception of respondents on usage of organic fertilizer ($p = 0.05$). Advocacy should be intensified on usage of organic fertilizer in crop production.

Keywords: Perception; arable crop; organic fertilizer; maize production

Introduction

Maize is one of the most important food crops in Africa and the developing world in general (Forum for Agricultural Research in Africa (FARA), 2009). Together with rice and wheat, maize provides at least 30% of the food calories to more than 4.5 billion people in 94 developing countries (Von Braun, Byerlee, Chartres, Lumpkin, Olembo and Waage, 2010). About 67% of the total maize production in the developing world comes from low and lower middle income countries; hence, maize plays an important role in the livelihoods of millions of poor farmers (FAOSTAT, 2010).

According to Food and Agricultural Organization (FAO) (2000), maize is a staple food that can remedy food insecurity in the tropics. In spite of the economic importance of maize to the teeming populace in Nigeria, its production has not over time met food and industrial needs of the country. The demand for maize as reported by Adebawale (2004) exceeds its supply as a result of its additional uses as livestock feeds, baking and brewing industries. There is need for intensification of the production. However, the production is facing the major challenge that is attributed to the fertility status of most Nigerian soils which is considerably low. There has been much concern on how to improve the fertility and productivity of the soil for increased agricultural production (Azagaku and Anzaku 2002).

The use of inputs as soil amendments in the form of mineral fertilizer or organic manure have been found useful in improving the productivity of the soil. Recognizing the myriad of deficiencies inherent in the use of inorganic fertilizer in crop production, agricultural based agencies have advocated the use of organic manures as alternatives to inorganic fertilizer (Adeoye, 1985). Usage and benefits derived from organic fertilizer have been discussed by many authors (Adediran *et al.*, 2003; Adetunji, 2010). Unlike inorganic fertilizer, nutrients contained are released more slowly and are stored for a longer time in the soil, ensuring longer residual effects, improved root development and higher crop yields (Abou El Magd *et*

al., 2006). Organic fertilizer also helps to improve soil by lowering bulk density, reduce soil erosion and improve soil fertility (Wheeler, 2008). Issues of underground water pollution by some of the inorganic fertilizers and the need to have a less polluted environment also justify the use of organic manure.

However, the use of organic fertilizer nowadays is being advocated in the production of organic food products which plays a great role in the health of man. The study therefore examined the perception of arable crop farmers' on use of organic manure in maize production.

Objectives of the study

The general objective of the study was to examine the perception of arable crop farmers on use of organic fertilizer in maize production in Ido Local Government Area of Oyo State, Nigeria. Specifically, the study was meant to:

- i. describe **personal** characteristics of maize farmers in the study area;
- ii. **identify** sources of organic fertilizer for maize production;
- iii. ascertain level of usage of organic fertilizer on maize production;
- iv. examine the challenges associated with **usage** of organic fertilizers in maize production;
- v. examine famers' perception of use of organic fertilizer for maize production.

Methodology

The study was carried out in Ido Local Government Area, Oyo State, with its headquarters at Ido. The study area has a land mass covering an area of **986 km²** and a total population of 103,261 (NPC, 2006). Ido Local Government Area covers the area spanning Apata, Ijokodo, Omi-Adio, Akufo and Apete. The climatic condition of Ido is equatorial, notably with dry and wet seasons with relatively high humidity.

The population of this study was the arable crop farmers in Ido Local Government Area of Oyo State. **Ten (10) villages were purposively selected for the study because of their dominant involvement in maize production.** The villages include Sinkooke, Sinkoisale, Odetola, Onigbagbo, Alakaso I, Alakaso II, Adeseku, Tade, Oderemi and Alagba. Twelve arable crop farmers were randomly selected each from the ten villages making a total of one hundred and twenty respondents. Data were collected with the use of well-structured questionnaire and interview guide. **The Statistical Package for Social Science (SPSS) version 16 software was used to analyze the data.**

Measurements of variables

The dependent variable of the study was the perception of arable crop farmers on the use of organic fertilizers. A list of perception statements was presented to respondents. Responses were recorded on a four-point likert- type scale of strongly agree (SA), agree (A), disagree (D) and strongly disagree (SD) and scored as 3, 2, 1 and 0 respectively for positive statements and 0, 1, 2 and 3 respectively for negative statements. The average score is 1.5, values greater than or equal 1.5 was regarded as favourable perception while scores less than 1.5 was regarded as unfavourable perception.

Frequency of usage of organic fertilizer for arable crop production was measured on a three point scale and scored as very often = 3, often = 2 and rarely = 1 with an average score of 2. Perceived constraints associated with organic fertilizer were measured using three point likert type scale and scored as very severe (VS) = 2, Severe (S) = 1 and not severe (NS) = 0. Meanwhile, perceived benefit associated with the use of organic fertilizer were recorded on a two-way response format of yes = 1, and no = 0.

Data Analysis

The data collected were analyzed using descriptive statistics such as frequency count, percentages and mean. An inferential statistics such as Chi-square was used to test the study hypothesis.

Results and discussion

Personal characteristics of respondents

Results on age distribution of respondents as found in Table 1 indicate that less than half of the respondents (41.7%) were between the age range of 26 - 35years. The mean age was found to be 41years which implies that the respondents were still in their active age. This indicates a good supply of agile workforce in maize production which tends to have positive effect on individual's efficiency and seriousness. Findings further revealed that most of the respondents (77.5%) were male. This suggests that maize production in the study area was male dominated. This may be attributed to the accessibility and control of resources as well as land tenure system which tend to favour men. It was also revealed that 72.5% of the respondents were married which implies that respondents were responsible for the provision of means of survival for their family. Spouses and children may constitute source of labour for farming operations. Findings on religious status show that a little below average (46.7%) were Christian. This implies that the respondents were predominantly Christians. About half (51.7%) had secondary school education, 37.5% had primary school education, 6.7% possessed higher education while 4.2% of the respondents had non-formal education. The findings indicate that the respondents had the basic education which is needed for interpretation of agro-information. Similarly, farmers with basic knowledge on how to read and write can appreciate and use most of the crop production technologies available. Findings further revealed that the respondents had an average household size of 5 which means that respondents had considerable smaller number of household. The distribution based on farming experience among the respondents indicates that 59.2% had between 1 to 10years of farming experience while very few (33.3%) had between 11 to 20years experience. Years of experience plays a prominent role in farming activities because it determines the quantum of knowledge gathered over time for better production.

Table 1: Personal characteristics of respondents (n = 120)

Variables	Frequency	Percentage (%)	Mean
Age (Years)			
16 – 25	23	19.2	41years
26 – 35	50	41.7	
36 – 45	34	28.3	
46 and above	13	10.8	
Sex			
Male	93	77.5	
Female	27	22.5	
Marital status			
Single	17	14.2	
Married	87	72.5	
Divorced	5	4.2	

139	Widowed	11	9.2	
140	Religion			
141	Christianity	56	46.7	
142	Islam	50	41.7	
143	Traditional	14	11.7	
144	Educational background			
145	No formal education	5	4.2	
146	Primary education	45	37.5	
147	Secondary education	62	51.7	
148	Tertiary education	8	6.7	
149	Household size			
150	1-5	63	54.2	5
151	6-10	42	34.1	
152	10 and above	15	2.5	
153	Years of farming experience			
154	1 to 10	71	59.2	
155	11 to 20	40	33.3	17 years
156	21 to 30	6	5.0	
157	31 and above	3	2.5	
158	Source: Field survey, 2015			

159

160 **Sources of organic fertilizer for arable crop production**

161 Findings in Figure 1 show that poultry manure (90.0%) constitute the major source of organic
 162 fertilizer used in arable crop production. This may be attributed to its availability and
 163 reported nutrient contents which were found to be more compare to other sources of organic
 164 fertilizers. This is supported by Nweke, Ijearu, Igili, (2013) findings that poultry manure was
 165 the richest manure among goat and pig manure applied at the same rate. This was also
 166 corroborated by Weevil (2002) that application of poultry manure to the soil enhances
 167 satisfactory growth in crops such as okro. Aliyu (2000) and Dauda *et al.* (2008) made similar
 168 observation in their separate studies with poultry manure in the growth and yield of crops.
 169 However, other sources of organic fertilizer used by the respondents include cow dung
 170 (71.7%), compost (48.3%), household waste (42.5%), crop residue (42.5%) while equal
 171 percentage of the respondents (36.7%) indicated green manure and pig manure as other
 172 sources organic fertilizer.




Fig 1: Sources of organic fertilizer for arable crop production (n= 120)

Frequency of usage of organic fertilizer in maize production

Frequency of usage of organic fertilizers in maize production as found in Table 2 revealed that poultry manure ($\bar{x}$ = 2.62) and cow dung ($\bar{x}$ = 2.47) were the most frequently used source of organic fertilizers. The findings are expected since poultry and cow dung constitute the most available source of organic fertilizer as earlier reported. However, urban waste ($\bar{x}$ =1.59) was found to be the least frequently used source of organic fertilizer. This may be attributed to its unavailability within the farm communities.

The overall findings on frequency of usage of organic fertilizer revealed that organic fertilizer was rarely used in maize production as indicated by 42.5% of the respondents despite the benefits. However, usage of organic fertilizer is gradually gaining recognition in Nigerian farming system because of the advocacy for production of natural/ chemical free food.

Table 2: Frequency of usage of organic fertilizer by the respondents (n=120)

Category	Always Freq.	Occasionally Freq.	Not at all Freq.	Mean	Standard deviation
Poultry manure	85(70.8)	24(20.0)	11(9.2)	2.62*	0.65
Cow dung	68(56.7)	40(33.3)	10(10.0)	2.47*	0.67
Household waste	21(17.5)	51(42.5)	48(40.0)	1.78	0.73
Compost	31(25.8)	41(34.2)	48(40.0)	1.86	0.80
Green manure	25(26.8)	40(33.3)	55(45.8)	1.75	0.78
Urban waste	16(13.3)	39(32.5)	65(54.2)	1.59	0.72
Crop residue	21(17.5)	48(40.0)	51(42.5)	1.75	0.74
Pig manure	27(22.5)	34(28.3)	59(49.2)	1.73	0.81

199 Grand Mean 15.55
 200 Mean level 1.94
 201 Source, Field survey, 2015
 202 Figures in parentheses are percentages
 203 Overall frequency of usage: Mostly used = 35.0%, rarely used = 42.5%, Never used 22.5%

204
 205
 206
 207

208 **Perceived constraints associated with the use of organic fertilizers in maize production**
 209 Results in Table 3 reveal the constraints facing the usage of organic fertilizer in maize
 210 production. It was found that organic fertilizer is usually bulky ($\bar{x}$ = 2.34). This tends to
 211 create pressure on storage and transportation and consequently add to the cost of production
 212 in maize. It was also observed that large quantity of organic fertilizer will be needed to meet
 213 up with crop requirement ($\bar{x}$ = 2.35) unlike inorganic fertilizer. Organic fertilizer was also
 214 found to emits offensive odour ($\bar{x}$ = 2.33) which may predisposes farmers to health hazard.

215

216 **Table 3: Constraints facing the use of organic fertilizer (n=120)**

217 Constraints	218 Very severe Freq	Severe Freq	Not severe Freq	Mean	SD
219 Bulkiness	58(48.3)	55(45.8)	58(48.3)	2.43*	0.60
220					
221 Need for large quantity to					
222 meet crop requirement	57(47.5)	48(40.0)	15(12.5)	2.35*	0.69
223 Nutrients are					
224 released slowly	51(42.5)	43(35.8)	26(21.7)	2.21	0.78
225					
226 Emission of offensive					
227 odour	51(42.5)	58(48.3)	11(9.2)	2.33*	0.64
228					
229 Usage can lead to					
230 health hazard	49(40.8)	56(46.7)	15(12.5)	2.28	0.68
231 Presence of					
232 impurities	45(37.5)	59(49.2)	16(13.3)	2.24	0.67
233 Limited information					
234 on quantity to be used	55(45.8)	41(34.2)	24(20.0)	2.26	0.77
235 Grand mean				16.1	
236 Mean level				2.30	

237 Source, Field survey, 2015
 238 Figures in parentheses are percentages
 239 SD= Standard deviation
 240
 241

Perception on usage of organic fertilizer

Findings on perception of usage of organic fertilizer as found in table 4 show that more than half of the respondents (51.7%) strongly agreed that large quantity is needed to meet crop requirement. Fifty percent (50.0%) of the respondents strongly agreed that organic manure is easy to produce while less than half (44.2%) strongly agreed that it requires special skills for production and application. However, it was found that little above half of the respondents (52.5%) had unfavourable perception on use of organic fertilizers in maize production. This implies that maize farmers were more disposed to conventional farming which involves the use of inorganic fertilizer.

Table 4: Perception arable crop farmers on the use of organic fertilizers for maize production (n=120)

Category	SA	A	D	SD	Mean	Sd
Organic fertilizer is easy to produce	60(50.0)	33(27.5)	15(12.5)	12(10.0)	3.18	1.00
Large quantity is needed to meet crop requirement	62(51.7)	29(24.2)	24(20.0)	5(4.2)	3.23	0.91
Preparation is labour intensive	47(39.2)	48(40.0)	17(14.2)	8(6.7)	3.11	0.89
Usage increases soil micro organisms	45(27.5)	68(56.7)	4(3.3)	3(2.5)	3.29	0.65
Special skill is needed for production and application	53(44.2)	43(35.8)	15(12.5)	9(7.5)	3.17	0.92
Organic manure does no harm to soil microbes	42(35.0)	58(48.3)	16(13.3)	4(3.3)	3.15	0.77
Transportation is expensive	45(37.8)	48(40.0)	20(16.7)	7(5.8)	3.09	0.88
Usage is dangerous on human health	36(30.0)	44(36.7)	27(22.5)	13(10.8)	2.86	0.97
It is environmentally friendly	40(33.3)	57(47.5)	16(13.3)	7(5.8)	3.08	0.84
Application is very easy	43(35.8)	60(50.0)	10(8.3)	7(5.8)	3.16	0.81
It reduces incidence of pest and diseases	35(29.2)	54(45.0)	22(18.3)	9(7.5)	2.96	0.88
Not effective for early maturing crops	31(25.8)	40(33.3)	38(31.7)	11(9.2)	2.76	0.94
It is not economical	33(27.5)	35(29.2)	27(22.5)	25(20.8)	2.63	1.10

Source: Field survey, 2015

Figures in parentheses are percentages

SA = strongly agree, A= Agree, Disagree, SD = strongly disagree, Sd = Standard deviation

Overall perception: Favourable 47.5%, unfavourable 52.5%

Test of relationship between personal characteristics of respondents and perception of organic fertilizer usage

The result of chi-square analysis in Table 5 shows that age ($\chi^2 = 4.46$, $df = 1$) and educational level ($\chi^2 = 8.541$, $df = 3$) were found to be significantly related with the perception of respondents on use of organic fertilizer in maize production at 5% level of significance. This implies that perception of farmers on usage of organic fertilizer is developed overtime and may be attributed to the experience gained as they advance in age and education.

Table 5: Test of relationship between personal characteristics of respondents and perception of organic fertilizer usage

Socio-economic characteristics	χ^2	df	p-value	Decision
Sex	4.461	1	.07	not significant
Age	13.470	3	.00	significant
Education level	8.541	3	.04	significant
Marital status	6.061	3	.11	not significant
Religion	6.150	2	.07	not significant
Year of farming experience	3.736	3	.29	not significant

Significant at 5% probability level

Conclusion

Despite the relative advantages associated with the usage of organic fertilizer, there is a reservation towards it as most of the farmers had unfavourable attitude towards its usage. This is similar to the findings of Iqbal, Sattar, Islam, Islam and Mollah (2004). This may be associated with the identified challenges which include bulkiness, need for large quantity to meet crop requirement and emission of offensive odour which may lead to health challenge. It is suggested that age and educational level can positively change the attitude of the farmers towards usage of organic fertilizer in maize production. It is therefore recommended that farmers should be adequately informed on usage and benefits of organic fertilizer for maize production

REFERENCES

- Abou El-Magd, M. A. El-Bassiony, M. and Fawzy, Z. F. (2006). Effect of organic manure with or without chemical fertilizer on growth, yield and quality of some varieties of Broccoli plants. *Journal of Applied Science Research* 2(10):791-798. <http://www.insipub.com/jasr/2006/791-798.pdf>.
- Adebawale, E.A., (2004). Tropical uses of maize. Institute of African research and training Ibadan.
- Adediran, J. A., Taiwo, L. B. and Sobulo, R. A. (2003). Effects of organic waste and method of composting on compost maturity, nutrient composition and yield of two vegetable crops. *Journal of Sustainable Agriculture* 22 (4), 95-109.
- Adeoye, K. B. (1985). Effect of some management practices on soil crusting in the Savannah Soils. *Proceedings of International Conference on soil fertilizer, Soil tilth and Post-clearing of land in the humid tropics*. Soil Science Society of Nigeria. pp. 302-309.
- Adetunji, M.O. (2010). Factors affecting the use of organic fertilisers among small scale

330 farmers in Ogbomoso agricultural zone, Oyo State, Nigeria. pp. 66-71. In: Olabiyi,
331 T.I. et al. (eds.). Proceedings of the first technical workshop on Organic Agriculture.
332 Nigerian Organic Agriculture Network.

333 Aliyu L. (2002). Analysis of the chemical composition of some organic manure and their
334 effect on the yield and composition of pepper. *Crop Res.* 23: 362-368.

335 Azagaku, E. D. and Anzaku, H.. (2002). Effect of NPK Fertilizer Levels on the Growth and
336 Development of early Maturing Maize Cultivar (TZESR –W) . In M. Iloeje, G. Osuji,
337 Udoh H. and G Asumugha (eds). *Agriculture: A Basis for Poverty Eradication and*
338 *Conflict Resolution. Proceedings of the 36th Annual Conference of the Agricultural*
339 *Society of Nigeria Held at the Federal University of Technology Owerri, Nigeria, 20-*
340 *24 October 2002, PP.106-111.*

341 Dauda S. N., Ajayi, F. A., Ndor, E. (2008). Growth and Yield of Watermelon (*Citrullus*
342 *lanatus*) as Affected by Poultry Manure Application. *J. Agric. Social Sci.* 121–124.
343 <http://www.fspublishers.org> (accessed 2009 November 10).

344 Forum for Agricultural Research in Africa (FARA) (2009). Patterns of change in maize
345 production in Africa: implications for maize policy development. Networking Support
346 function 3. Regional Policies and Markets.

347 Food and Agriculture Organization of the United Nations (FAO). (2000). Tropical maize.
348 Improvement and production. Food and Agriculture Organization of the United
349 Nations (FAO) Plant Production and Protection Series, Vol. 28. R.L. Paliwal, Ed.: 1–
350 363, Rome: Food and Agriculture Organization of the United Nations.

351 FAOStat. (2010). [faostat.fao.org]

352 Forum for Agricultural Research in Africa (FARA), (2009). Networking Support Function
353 3: Regional Policies & Markets. Patterns of Change in Maize Production in Africa:
354 Implications for Maize Policy Development. Ministerial Policy Brief Series (Number
355 3). 8pages.

Iqbal, S. M. A., Sattar, M. A., Islam, N., Islam G. M. M., and Mollah M. R. A. (2014).
Assessment of Farmers Perception on the Application of Chemical Fertilizers and
Organic Manures in Chuadanga Region. *Journal of Environmental Science &*
Natural Resources, 7(2): 69-72,

356 National Population Commission (2006). Official Website of National Population
357 Commission (NPC). Abuja.

358 Nweke, I. A., Ijearu, S. I. and Igili, D. N. (2013). Effect of different sources of animal
359 manure on the growth and yield of okra (*Abelmoschus esculentus* L.Moench) In
360 Ustoxic Dystropept at Enugu South Eastern, Nigeria. *International Journal of Science*
361 *and Technology Research* 2 (3):135-137.

362 Tian G, Kang B. T., Brussaard T. (1992). Biological effects of plant residues with contrasting
363 chemical compositions under humid tropics incomplete

364 Von Braun, J., Byerlee, D., Chartres, C., Lumpkin, T., Olembo, N. and Waage, J. (2010)
365 Draft Strategy and Results Framework for the CGIAR. CGIAR, The World Bank,
366 Washington D.C

367 Wheeler, S. A. (2008). What influences agricultural professionals' views towards organic
368 agriculture? *Ecological Economics* 65, 145-154.

369