

An improvement of robust estimator using known values of Probability weighted moment for finite population variance

*M A Bhat, S Maqbool, S A Mir, Immad.A.Shah

Division of Agricultural Statistics SKUAST-Kashmir (190025)

Abstract: In this study new improved robust estimator has been proposed for precise estimation of finite population variance in simple random sampling by incorporating as auxiliary information of probability weighted moment. Properties associated with proposed estimators are assessed by mean square error and bias through numerical demonstration. We have also provided theoretical efficiency comparison of the study.

Keywords: Ratio estimator, Probability Weighted moment; SRSWOR, MSE, Bias and Efficiency.

1. Introduction

Here we consider a finite population $U = \{U_1, U_2, \dots, U_N\}$ of N distinct and identifiable units.

Let Y be a real variable with value Y_i measured on $U_i, i=1,2,3,\dots,N$ given a vector $[Y_1, Y_2, Y_3, \dots, Y_N]$. Sometimes in sample surveys information on auxiliary variable X correlated with study variable Y , is available can be utilized to obtain the efficient estimator for the estimation of Population variance. With this this modification our aim is to estimate

the population variance $S_y^2 = \frac{1}{N-1} \sum_{i=1}^N (y_i - \bar{y})^2$ on the bases of random sample selected from

population U to obtain the improved and precise estimates. Various authors have proposed different ratio estimators for finite population variance by utilizing the ancillary information of different parameters of population such as Isaki [2], Kadilar and Cingi [3], Subramani and Kumarapandian [5] and Bhat *et al.* [1].

2. Material and Methods:

2.1 Notations: N = Population size. n = sample size. $\gamma = \frac{1}{n}$, Y = study variable. X =

Auxiliary variable. $\bar{X}, \bar{Y}$ = Population means. $\bar{x}, \bar{y}$ = Sample means. S_y^2, S_x^2 = Population variances. s_y^2, s_x^2 = sample variances. C_x, C_y = Coefficient of variation. ρ = Correlation coefficient. $\beta_{1(x)}$ = Skewness of the auxiliary variable. $\beta_{2(x)}$ = Kurtosis of the auxiliary variable. $\beta_{2(y)}$ = Kurtosis of the study variable. $B(.)$ = Bias of the estimator.

$MSE(.)$ = Mean square error. $\hat{S}_R^2$ = Ratio type variance estimator. S_{pw} = probability Weighted Moments

3. Suggested Estimator:

$$\hat{S}_{MS1}^2 = s_y^2 \left[\frac{S_x^2 + S_{pw}}{s_x^2 + S_{pw}} \right]$$

We have derived here the bias and mean square error of the proposed estimator to first order of approximation as given below:

Let $e_0 = \frac{s_y^2 - S_y^2}{S_y^2}$ and $e_1 = \frac{s_x^2 - S_x^2}{S_x^2}$. Further we can write $s_y^2 = S_y^2(1 + e_0)$ and $s_x^2 = S_x^2(1 + e_1)$

and from the definition of e_0 and e_1 we obtain:

$$E[e_0] = E[e_1] = 0. \quad E[e_0^2] = \frac{1-f}{n} (\beta_{2(y)} - 1). \quad E[e_1^2] = \frac{1-f}{n} (\beta_{2(x)} - 1). \quad E[e_0 e_1] = \frac{1-f}{n} (\lambda_{22} - 1)$$

The proposed estimator is given below:

$$\begin{aligned}\hat{S}_p^2 &= s_y^2 \left[\frac{S_x^2 + \alpha S_{pw}}{s_x^2 + \alpha S_{pw}} \right] \\ \Rightarrow \hat{S}_p^2 &= s_y^2 (1 + e_0) \left[\frac{S_x^2 + \alpha S_{pw}}{s_x^2 + e_1 S_x^2 + \alpha S_{pw}} \right] \\ \Rightarrow \hat{S}_p^2 &= \frac{S_y^2 (1 + e_0)}{(1 + A_p e_1)} \text{ where } A_p = \frac{S_x^2}{(S_x^2 + \alpha S_{pw})}\end{aligned}$$

$$\Rightarrow \hat{S}_p^2 = S_y^2 (1 + e_0) (1 + A_p e_1)^{-1}$$

$$\Rightarrow \hat{S}_p^2 = S_y^2 (1 + e_0) (1 - A_p e_1 + A_p^2 e_1^2 - A_p^3 e_1^3 + \dots)$$

Expanding and neglecting the terms more than 3rd order, we get

$$\hat{S}_p^2 = S_y^2 + S_y^2 e_0 - S_y^2 A_p e_1 - S_y^2 A_p e_0 e_1 + S_y^2 A_p^2 e_1^2 \quad (1)$$

$$\Rightarrow \hat{S}_p^2 - S_y^2 = S_y^2 e_0 - S_y^2 A_p e_1 - S_y^2 A_p e_0 e_1 + S_y^2 A_p^2 e_1^2 \quad (2)$$

By taking expectation on both sides of (1), we get

$$E(\hat{S}_p^2 - S_y^2) = S_y^2 E(e_0) - S_y^2 A_p E(e_1) - S_y^2 A_p E(e_0 e_1) + S_y^2 A_p^2 E(e_1^2)$$

$$BiasE(\hat{S}_p^2) = S_y^2 A_p^2 E(e_1^2) - S_y^2 A_p E(e_0 e_1) \quad (3)$$

$$BiasE(\hat{S}_p^2) = \gamma S_y^2 A_p [A_p (\beta_{2(x)} - 1) - (\lambda_{22} - 1)]$$

Squaring both sides of (2), neglecting the terms more than 2nd order and taking expectation, we get

$$E(\hat{S}_p^2 - S_y^2)^2 = S_y^4 E(e_0^2) - S_y^4 A_p^2 E(e_1^2) - 2S_y^4 A_p E(e_0 e_1)$$

$$MSE(\hat{S}_p^2) = \gamma S_y^4 [(\beta_{2(y)} - 1) + A_p^2 (\beta_{2(x)} - 1) - 2A_p (\lambda_{22} - 1)]$$

4. Efficiency conditions In this section, we have derived the efficiency conditions of proposed estimators with other existing estimators under which proposed estimator are performing better than the existing estimators

The bias and Mean square error of existing ratio type estimators up to the first order of approximation is given by

$$BiasE(\hat{S}_K^2) = \gamma S_y^2 R_K [R_K (\beta_{2(x)} - 1) - (\lambda_{22} - 1)] \quad (1)$$

$$MSE(\hat{S}_K^2) = \gamma S_y^4 [(\beta_{2(y)} - 1) + R_K^2 (\beta_{2(x)} - 1) - 2R_K (\lambda_{22} - 1)] \quad (2)$$

Where $R_K = \text{Constant}$, $K = 1, 2, 3$

Bias, MSE and constant of proposed estimators is given by

$$Bias(\hat{S}_p^2) = \gamma S_y^2 R_p [R_p (\beta_{2x} - 1) - (\lambda_{22} - 1)] \quad (3)$$

$$MSE(\hat{S}_p^2) = \gamma S_y^4 [(\beta_{2y} - 1) + R_p^2 (\beta_{2x} - 1) - 2R_p (\lambda_{22} - 1)] \quad (4)$$

$R_p = \text{Constant}$, $p = 1$

From Equation (4) and (6), we have

$$\begin{aligned}MSE(\hat{S}_p^2) &\leq MSE(\hat{S}_K^2) \text{ if } \lambda_{22} \geq 1 + \frac{(R_p + R_K)(\beta_{2x} - 1)}{2} \\ MSE(\hat{S}_p^2) &\leq MSE(\hat{S}_K^2) \\ \gamma S_y^4 [(\beta_{2y} - 1) + R_p^2 (\beta_{2x} - 1) - 2R_p (\lambda_{22} - 1)] &\leq \gamma S_y^4 [(\beta_{2y} - 1) + R_K^2 (\beta_{2x} - 1) - 2R_K (\lambda_{22} - 1)]\end{aligned} \quad (5)$$

$$\Rightarrow [(\beta_{2y} - 1) + R_p^2(\beta_{2x} - 1) - 2R_p(\lambda_{22} - 1)] \leq [(\beta_{2y} - 1) + R_K^2(\beta_{2x} - 1) - 2R_K(\lambda_{22} - 1)] \quad (6)$$

$$\Rightarrow [1 + R_p^2(\beta_{2x} - 1) - 2R_p(\lambda_{22} - 1)] \leq [1 + R_K^2(\beta_{2x} - 1) - 2R_K(\lambda_{22} - 1)] \quad (7)$$

$$\Rightarrow (\beta_{2x} - 1)(R_p^2 - R_K^2) [-2R_p(\lambda_{22} - 1)] \leq [-2R_K(\lambda_{22} - 1)] \quad (8)$$

$$\Rightarrow (\beta_{2x} - 1)(R_p^2 - R_K^2) [-2(\lambda_{22} - 1)(R_p - R_K)] \leq 0 \quad (9)$$

$$\Rightarrow (\beta_{2x} - 1)(R_p^2 - R_K^2) \leq [2(\lambda_{22} - 1)(R_p - R_K)] \quad (10)$$

$$\Rightarrow (\beta_{2x} - 1) \leq \frac{2(\lambda_{22} - 1)(R_p - R_K)}{(R_p^2 - R_K^2)} \quad (11)$$

$$\Rightarrow (\beta_{2x} - 1) \leq \frac{2(\lambda_{22} - 1)(R_p - R_K)}{(R_p - R_K)(R_p + R_K)} \quad (12)$$

$$\Rightarrow (\beta_{2x} - 1)(R_p + R_K) \leq 2(\lambda_{22} - 1) \quad (13)$$

By solving equation (13), we get

$$MSE(\hat{S}_p^2) \leq MSE(\hat{S}_K^2) \text{ if } \lambda_{22} \geq 1 + \frac{(R_p + R_K)(\beta_{2x} - 1)}{2}$$

5. Numerical Illustration:

We use the data of Murthy [4] page 228 in which fixed capital is denoted by X (auxiliary variable) and output of 80 factories are denoted by Y (study variable). we apply the proposed and existing estimators to this data set and the data statistics is given below:

N=80, $S_x=8.4542$, $n=20$, $C_x=0.7507$, $\bar{X} = 11.2624$, $\beta_{2(x)} = 2.8664$, $\bar{Y} = 51.8264$, $\beta_{2(y)} = 2.8664$, $\rho = 0.9413$, $\beta_{1(x)} = 1.05$, $\lambda_{22} = 2.2209$, $S_y=18.3569$, $Md=7.5750$, $Q1=9.318$, $C_y=0.3542$, $G = 9.0408$, $D = 8.0138$, $S_{pw} = 7.9136$

Table 1: Bias and Mean Square Error of the existing and the proposed estimators

Estimators	Bias	Mean Square Error
Isaki	10.8762	3925.1622
Kadilar&Cingi	10.4399	3850.1552
Subramani&Kumarapandiyam	6.1235	3180.7740
Proposed	5.2264	2474.4535

Table 2: Percent relative efficiency of proposed estimators with existing estimators

Estimators	Isaki	Kadilar and Cingi	Subramani and Kumarapandiyam
Proposed	158.6274	155.5961	128.5445

Conclusion:

From the above study we revealed that our suggested estimator perform better than the existing estimators as the mean square error as well as bias s too much lower than existing estimators.

Literature cited:

- [1]. Bhat M. A., Maqbool, S., Saraf, S. A., Rouf A. and Malik, S. H. Variance estimation using linear combination of skewness and quartiles. *Journal of Advances in Research* 13(2): 1-6, 2018 Article no.37321. ISSN: 2348-0394, NLMID: 101666096
- [2]. Isaki, C.T. (1983). Variance estimation using auxiliary information. *Journal of the American Statistical Association*, 78, 117-123.
- [3]. Kadilar, C. & Cingi, H. (2006a). Improvement in Variance estimation using auxiliary information *Hacettepe Journal of mathematics and Statistics*, 35(1). 117-115.
- [4]. M. N Murthy, Sampling theory, *Theory and Methods*, Statistical publishing Society, Calcutta, 1967.

[5]. Subramani, J. and Kumarapandiyan, G. (2015). Generalized modified ratio type estimator for estimation of population variance. *Sri-Lankan journal of applied Statistics*, vol 16-1, 69-90.