

Bacterial Diseases of Plants in Nepal: A Review

ABSTRACT

Bacterial diseases are the important disease next to the fungal diseases in Nepal. In Nepal, major bacterial diseases are Bacterial leaf blight of rice (*Xanthomonas oryzae* pv. *oryzae*), Bacterial wilt of potato and tomato (*Ralstonia solanacearum*), Citrus greening (*Candidatus liberibacter*), Citrus canker (*Xanthomonas campestris* pv. *citri*), Soft rot of potato (*Erwinia carotovora* pv. *atroseptica*) and Black rot of crucifers (*Xanthomonas campestris* pv. *campestris*) as they are prevailed in most of areas and cause the devastating losses. Other are the minor with less economic importance like Bacterial stalk of maize (*Erwinia chrysanthemi* pv. *zeae*), bacterial postulates of soybean (*Xanthomonas campestris* pv. *phaseoli*), Potato scab (*Streptomyces scabies*). Some of the are minor diseases like Stewart's wilt of corn (*Pantoea stewartii*), Bacterial speck of tomato (*Pseudomonas syringae* pv. *tomato*), Bacterial spot of tomato (*Xanthomonas campestris* pv. *vesicatoria*), Bacterial sheath rot of wheat (*Pseudomonas fuscovaginitiae*), Crown gall (*Agrobacterium tumefaciens*), Bacterial leaf strip (*Xanthomonas rubrilineans*) and Bacterial spots of pumpkin (*Xanthomonas cucurbitae*) some are only recorded in Nepal.

Keywords: Bacterial; Diseases; Economic importance; major

INTRODUCTION:

Diseases of the crop plants are the major constraints in production of agricultural crops. In Nepal, bacterial diseases are most prevalent in the crops next to the fungal diseases. Most destructive bacterial diseases are Bacterial leaf blight of rice, bacterial leaf streak of rice, wilts of solanaceous crops, citrus greening and canker and black rot of crucifers. And other are considered as a minor diseases. The research works on bacterial disease was initiated in Nepal after the establishment of Plant Pathology Section, established in 1954. Now, Nepal Agriculture Research Council, Institute of Agriculture and Animal Sciences and Agriculture and Forestry University, NAST, Prubanchal University are active in research of phytopathological diseases in Nepal. Phytopathological research are mainly focused in the fungal disease and less research are conducted in bacterial, viral and nematodes.

1. BACTERIAL DISEASES OF RICE

1.1. Bacterial blight

BB [bacterial blight] caused by *Xanthomonas oryzae* pv. *oryzae* was first observed in the terai region with the introduction of high-yielding rice varieties. In Nepal, BLB was reported as early as 1968 in Kathmandu valley [1]. However, it spread rapidly in subsequent years and is now prevalent throughout the terai and hills.

1.1.1. Occurrence and Economic importance

The disease was reported to occurring widely in Terai and inner Terai region of Nepal causing yield loss [2] and incidence was higher in eastern Terai than in the western part [3]. The combination of rainy weather, strong winds, and moderate temperatures in Kathmandu favoured rapid disease spread [4]. The disease causes severe damage in rice production [5] and can reduce grain yields up to 26% especially during prolonged periods of rainy weather from August

to October. Similarly, yield losses of 13-32% were reported on susceptible cultivar Taichung Native-1 [6]. It reduced the yield from 5-60 % in Terai and mid—hills during hot and humid periods [7].

1.1.2.Pathogenic variability:

The pathogenic races varied from location to location [8]. There are two major groups of *X. oryzae* pv. *oryzae* in Nepal. One group consists of strains with high molecular polymorphism and many pathotypes that are either virulent to the 11 major resistance genes or virulent only to Xa21. Strains in the second group have low molecular polymorphism and are avirulent to Xa4, xa5, Xa7, and Xa21 [9]. This disease has been confined to hilly regions and at altitude of about 1,238 m of Katmandu valley [4]. Nine pathogenic races were identified based on differential interactions on eight rice cultivars [10].

1.1.3.Virulence

The ability of an isolate to cause lesions with different lengths across the lines was interpreted as virulence. The races were virulent to many important resistance genes and the genetic diversity was high (0.98) in the collection from Nepal [11]. Adhikari et al. found that a line with a four-gene combination, NaH56, showed higher level of resistance to *Xanthomonas oryzae* pv. *oryzae* than the other lines[12]. Most strains from South Asia (Nepal and India) were virulent to cultivars containing the bacterial blight resistance gene xa-5 [11].

1.1.4. Host resistance:

Continues research on host resistance of bacterial blight has been conducted and found different resistant and susceptible varieties in different time and locations. IR36, IR54, IR60, and IR64 showed high levels of resistance to *Xanthomonas oryzae* pv. *oryzae* [5]. The research was conducted on resistance of different varieties to the different strains of *Xanthomonas* and found IR-BB8 exhibited a high level of resistance to nearly two-thirds of the strains of *Xanthomonas oryzae* pv. *oryzae* that were evaluated [13].

Rice cultivars BR-34-13, PAU-50-B-25, Lawmi, Sabitri, BW293-21, IR7167-33, Rodina, are resistance [14]. Burlakoti et.al. reported Chandina, Bindeshwori and Chaite-2 cultivars as a moderately resistant [7]. Also, Research carried out under Bhairahawa conditions showed BR 4962-12-4, Barkhe 2001, BRR1 Dhan-26, NR 1988-2-3-5, JR 7699-44-3-3-1 and Barkhe 2045 were the best genotypes having stable and higher level of resistance against bacterial blight[15] . Bhatta et.al. reported BLB resistance genes Xa-10, Xa-13, Xa-7, Xa-3, Xa-4, Xa-5, Xa-8 in Nepal. The Xa-3, Xa-4, xa-5, Xa-7, xa-8, Xa-10 and xa-13 was found on several rice accessions, but not Xa-21[16]. Nepalese rice accessions lack the Xa-21 genes which is more important for controlling BLB epidemics throughout the world [17].

Hardinath-1, Tarahara-1, Sukkhadhan-1, Sukkhadhan-2, Sukkhadhan3, Hardinath-2, Swarnas ub1, Samba mahasuri sub1 and Mithila were resistant to blast and BLB [18]. Dangal, et.al, carried screening of 134 genotypes out of which 34 genotypes were moderately resistant to BLB [19]. The survey was conducted in three districts of mid-eastern regions of Nepal. Low disease incidence were observed in Makawanpur, Radha 4 and Sabitri varieties of rice at Baridya and Dang, varieties SukhaDhan 1 and SukhaDhan 2 are resistant at Banke[20].

1.1.5.Chemical control:

Blitox or Validamycine @ 2-3 gm/litre of water was effective in controlling BLB [18]. Foliar spray of 100 ppm streptocycline has been recommended against the disease bacterial leaf blight of rice [7].

1.1.6.Cultural control

The application of 2-3 handfuls of cow dung in 10 litre of water each day for 14 helps to reduce the bacterial leaf blight of rice. Also application of aqueous extract of agave is effective for control of BLB of rice.

1.2. Bacterial sheath brown rot

Bacterial sheath brown rot is an important disease of rice in high altitude and high rainfall areas of Nepal. Sheath brown rot caused by *Pseudomonas fuscovaginae* was first reported from Hokkido, Japan and has been found to be induced by chilling temperatures. The disease is seed borne and from seeds in early 1990s [21]. The greenish, light brown highly fluorescent colonies are isolated from the sample above 1400m while some suspected colonies were observed in lower altitude like ~~1010m~~ Siukhet [22].

2. BACTERIAL DISEASES OF MAIZE

Maize is a most important cereal crop in Nepal and also the principal staple cereal diet of most of the Nepalese people who mainly lives in hilly region. Maize plants are affected by wide range of pathogens with fungal and bacterial diseases being the most important in Nepal. Maize is susceptible to a lesser no of pathogenic bacteria. Three bacterial species have been recorded pathogenic to maize crop in Nepal [23].

2.1. Bacterial stalk rot of maize

Bacterial stalk rot [*Erwinia chrysanthemi* pv. *zeae* (Sabet) Victoria, Arboleda & Munoz] was first reported in 1968 at Rampur [24]. Stalk rot is distributed throughout the country, but it is most prevalent in the hot and humid areas [25]. It is mostly confined to terai and inner terai as a major disease also it is extended to some pockets of the mountains and Kathmandu valley but as minor disease [24]. Bacterial stalk rot caused up to 80% yield loss along with fungal diseases in maize in the plains. Sarlahi Seto and E2 Illinois were moderately resistant [7].

2.2. Bacterial leaf strip and Stewart Wilt of Maize

Xanthomonas rubrilineans is the causal organism of minor disease bacterial leaf strip of maize which was first reported by Manandhar in 1976 at Rampur. The Stewart wilt of maize caused by *Erwinia stewartii* (E.F. Smith Dye) Dowson also reported from Rampur for the first time in 1978 [24].

3. BACTERIAL DISEASE OF WHEAT

3.1. Bacterial sheath rot

The bacterial sheath rot of wheat is caused by bacterium *Pseudomonas fuscovagittcae*. The bacterium was noted from Nigale farm in 1994 and Lumle Agriculture Center in 1995. The bacteria was identified by cultural characteristics biochemical and pathogenicity test. This was the first report of the disease. Annapurna-1, Annapurna-2, Annapurna-3 and WK.685 noted to be highly infected [22].

4. BACTERIAL DISEASES OF CITRUS

4.1. Citrus greening

Citrus greening also known as Huanglongbing (HLB) disease is one of the major disease of citrus caused by a vectored pathogen. The causative agent is *Candidatus Liberibacter*, fastidious phloem-restricted, gram-negative bacteria. Transmission is by the Asian citrus psyllid (*Diaphorina citri*) and also by graft transmission [26]. Citrus decline was first reported in Pokhara valley by Thrower in 1968 [27]. Based on visual observation, Knorr et al suspected that the citrus decline was caused by greening disease entered with the planting materials introduced to Horticulture Research Station, Pokhara from Saharanpur, India [28]. HLB has been considered the number one among many biotic factors contributing to low productivity and threat to citrus industry in Nepal [29].

4.1.1. Occurrence and economic importance:

Up to 39-55 % mandarin trees were found infected with greening disease in Pokhara valley [28]. The symptoms of Citrus Greening was noted on about 55% of citrus trees in Pokhara valley and 100% in Horticulture Research Station in 1980s [30]. More recent HLB is widespread in many citrus pockets of Kaski, Syanja, Tanahu, Lamjung and Dhading districts [29, 31]. This disease had resulted to massive

decline of citrus orchards especially located below 1000 m altitude where population of vector – psyllid is abundant [32].

4.1.2.Host resistance:

Most of the commercial citrus species of Nepal like mandarin and sweet orange are very susceptible to the disease while acid lime is slightly tolerant, but it carry HLB bacterium which source of inoculums of disease [32].

4.1.3.Chemical Control:

In citrus, greening was epidemic throughout the country. Proper management of orchard followed by copper fungicide spray reduced the infection [7].

4.2. Citrus Canker:

Citrus canker is one of the problematic diseases of citrus which affect almost all types of important citrus crops. The disease causes extensive damage to citrus which probably originated in Southeast Asia or India, and now occurs in more than 30 countries [33]. The disease is caused by *Xanthomonas campestris* (=axonopodis)pv. citri(Xcc) [34].

4.2.1.Management:

The local varieties/cultivars of mandarin and sweet oranges were found resistant to citrus canker. The lime was very susceptible and special care should be taken to control it with copper based chemical sprays [35]. Also reported as Copper hydroxide, basic copper chloride, copper oxychloride, and tribasic copper sulfate are the most effective bacterial sprays for protecting leaves and fruit from attack of *X. citri*. In citrus, canker can be managed by proper management of orchard followed by copper fungicide spray reduced the infection [7].

5. BACTERIAL DISEASES OF TOMATO

5.1. Bacterial wilt of tomato

Bacterial wilt caused by *Ralstonia solanacearum* (Formerly *Pseudomonas solanacearum* E.F. Smith) is one of the most important and wide spread bacterial diseases of crop plants in Nepal. The disease was first reported in early 1960s in the Kathmandu valley 1300 m above sea level [36].

5.1.1.Occurrence and Economic Importance

Bacterial wilt (*Ralstonia solanacearum*) incidence was recorded up to 100%in tomato, in terai and mid-hills under hot and humid conditions [7].

5.1.2.Pathogenic variability:

Adhikari et.al. isolated the 25 strains of *Pseudomonas solanacearum* and they placed 8 isolates from high hills in biovar 2 and the rest from lowland subtropics in biovar 3. On the basis of pathogenicity tests, 8 strains were of race 3 and 17 strains were of race 1. Race 1 and biovar 3 are first reports from Nepal [37].

5.1.3.Host resistance:

The partially resistant tomato lines are CL1131-0-0-43-4 (CL1131) and Pusa Ruby a susceptible tomato cultivar [38]. Hybrids such as Srijana, CLN 2026 C and CLN 2026 D are the resistant varieties for the bacterial wilt in Nepal [39].

5.1.4.Cultural control:

Crop rotation to corn, lady's finger, or cowpea, and cultivar resistance appeared to be useful management strategies to control bacterial wilt of tomato in Nepal [38].Grafting technology using resistant root stocks has been successful for managing bacterial wilt in solanaceous vegetables.

5.2. Bacterial spot of tomato:

The first incidence of the disease was observed in cultivar Pusa Ruby and in three different commercial farms at Banke district, Mid-western region with incidence ranging 50-60% in Nepal in 2009 [40]. The dark, circular and water-soaked lesions of about 3 mm in diameter were observed on leaves of tomato plants (*Solanum lycopersicum*).

5.3. Bacterial speck of tomato

Bacterial speck of tomato is caused by *Pseudomonas syringae* pv. *tomato*. In local cultivar Baglung Local (BL) small necrotic flecks surrounded by chlorotic haloes 1.5–3.0 mm in diameter were observed on leaves of tomato plants during the spring of 2007 in an experimental farm in Kirtipur Nepal [41]. This was the first report in Nepal. Lamichhane et.al. tested 10 cultivars of tomato for susceptibility of disease, cultivars Thims 16, C.L. and Spectra 737 were found least susceptible in the field, while in the tunnel all the local cultivars (C.L., Panjabi, B.L. and Lapsi Gede) were very low susceptible [42].

6. BACTERIAL DISEASES OF POTATO

6.1. Bacterial wilt

Ralstonia solanacearum [43], caused wilt of potato and solanaceous crops including other host plants is formerly known as *Pseudomonas solanacearum* EF, is the second most limiting phyto pathological factor to potato production in Nepal. In Nepal, the disease is considered as the most important one that causes a considerable yield loss every year [44].

6.1.1. Occurrence and economic importance:

The disease is one of the destructive diseases of potato in hills and high hills. The bacterial wilt of potato was severe in the subtropical to cool highlands. Reports indicate that crop infection increased from 5-40% in 1977 to 25 80% in 1980 and tuber rotting in stores ranged from 10 to 20% in Nuwakot, Rasuwa, Sindhupalchok, Dolakha, Kaski, Makwanpur, and Palpa districts [45]. Bacterial wilt was also observed on some government farms and stations at Kakani (1500 m), Kitripur (1350 m), Daman (2500 m), and Nigale (2500 m) of the central hills, and Pokhara (900 m) of the western hills. Burlakoti et.al. recorded up to 100%, 90% and 32.5% incidence of wilt in tomato, brinjal and potato, respectively, in terai and mid-hills under hot and humid conditions [7]. Up to 100% rotting of potatoes in the store has since been recorded in severe cases [44].

6.1.2. Pathogen Variability

Isolates belonging to biovar 2 were all from potato plants isolated from the high hills [45]. Adhikari reported the race 3 and the biovar II in the potato from mid to high hill region by hypersensitivity, biochemical, cultural and serological test [46]. Race 3 and biovar II of the pathogen was widely spread over potato growing areas of mid and high hills of Nepal [47]. It was concluded that pathotypes and biotypes of bacterial wilt pathogens of potato were remained the same in Nepal from the last two decades.

6.1.3. Host Resistance:

According to Dhital, Potato genotypes BR-63.65, BS-63.5, BS-70.23, BS-35.25, BS-35.24, BS-35.22 and BS-36.3 were found free from the disease in two or more tested [48].

6.1.4. Chemical control

Research on chemical control of bacterial wilt of potato shows that use of full rate of Stable Bleaching Powder at 25 kg/ha in infested soil is highly effective for the control of diseases in terms of reduction of disease incidence and pathogen population [49].

6.1.5. Cultural management

Three year crop rotation with no-host crops, eradication of volunteer potato plants and farmers education on crop hygiene and disease management are the way of management of bacterial wilt of potato [48]. Hogger and Shrestha reported flooding paddy fields prior to potato crop reduces bacterial wilt so, farmers in irrigated areas of the mid hills and plains are advised to use a crop rotation with paddy for at least two seasons and use clean seed [50].

6.2. Potato Soft Rot

Potato soft (*Erwinia carotovora* pv *atroseptica*) was reported in the Kathmandu District, of central region of Nepal. The disease was suspected in local cultivars Rato Alu and SetoAlu.. The pathogen may have been introduced to this region of Nepal via seed potato tubers from other countries [51].

6.3. Potato scab

Potato scab is caused by ~~a bacterium-like organism~~, *Streptomyces scabies*. It is one of the feared diseases of potato in Nepal. The management of the potato scab can be done by dipping tubers in solution of acetic acid+zinc sulfate then in the solution of Blitox-50 and Emisan-6 [52]

7. BACTERIAL DISEASE OF CRUCIFERS

7.1. Black rot of crucifers

Black rot caused by *Xanthomonas campestris* pv. *campestris* (Pammel) Dowson (Xcc) is one of the economically important diseases of brassicaceous vegetable crops of Nepal. The disease is found in terai and inner terai regions where cauliflower and other cruciferous vegetables are cultivated. The black rot of crucifers in Nepal has been reported as early as 1977 .

7.1.1. Occurrence and economic importance

Black rot is a globally important disease of crucifers. The disease can cause significant losses, particularly in warm and humid environments. Burlakoti reported that black rot caused up to 24.5% infection in local cauliflower and up to 39% in mustard seeds [7]. In Nepal more than 60 % yield reduction in cauliflower has been reported [53].

7.1.2. Pathogens Variability

Cabbage strains belonged to five races (races 1, 4, 5, 6, and 7), with races 4, 1, and 6 the most common. All cauliflower strains were race 4 and race 6 was the strain of leaf mustard. Nepalese *Xanthomonas campestris* pv. *campestris* strains clustered separately from other *Xanthomonas* spp. and pathovars [54].

7.1.3. Management

~~Cabbage variety Copenhagen market 1 was found to be highly infected and least infection was found in Drum head [53].~~ As black rot of the crucifers is seed borne disease, seed soaking with 0.2 % of Copper sulphate and Zinc sulphate found effective.

8. OTHER BACTERIAL DISEASES IN NEPAL

8.1. Bacterial Postulates of soybean

Bacterial postulates of soybean, caused by bacterium *Xanthomonas campestris* pv. *phaseoli*. Neupane and co-workers have reported the bacterial pustule as a major disease of soybean in the mid hills [55]. Since exact losses due to soybean diseases have not been assessed under Rampur condition estimated yield losses up to 25% have been observed due to bacterial pustule [56]. The occurrence of the bacterial postulates (*Xanthomonas campestris* pv. *phaseoli*) of soybean disease varied with elevation [57]. It was observed Genotype SB 00122 to be highly resistant while PK 327, Pooja, G-8522, F 778817, JARS 87-I, TGX-31 1-230, Cobb, Kavre, G-1946 and TGX-15 19-ID were resistant to the bacterial postulates of soybean [56].

8.2. Crown Gall of Apple

Crown gall of apple, caused by *Agrobacterium tumefaciens* was major problem in pome fruits in Nepal which is the quarantine disease of Nepal. The control of crown gall of apple can be done by treating the root dipping in 750 ppm of agrimycin before transplanting of seedling [7].

8.3. Bacterial leaf spot of Pumpkin

Xanthomonas cucurbitae cause the disease bacterial leaf spot of pumpkin. J.R. Lamichhane et al. observed the small necrotic spots with chlorotic haloes on the leaf surface and also scab-like lesions on fruits of pumpkin on variety Arka Chandan in Bhaktapur and Kathmandu districts. This was the first report of the disease on pumpkin in Nepal [58].

CONCLUSION

This paper analyzed major works in the bacterial diseases of crops in Nepal. The researches are being conducted in major diseases like Bacterial leaf blight of rice Bacterial wilt of tomato and potato under heading pathogens types, economic importance, occurrence and disease management. Bacterial diseases of plants are usually very difficult to manage. Frequently, combinations of control measures are required to manage the diseases. So, there are some works done in the control measures of diseases in Nepal. Most of the works are on host resistance and few on chemical and cultural management of diseases.

References

1. Khadka BB, Shah SM, Lawat K. Plant disease in Nepal-A supplementary list. *FAO Plant protection Bulletin, Technical document*. 1968;66.
2. Shah SM .Important disease problems of some major crops in rainfed agriculture in Nepal.In:*Reviews of Pest, Disease and Weed problems in rainfed crops in Asia and the Far East*. FAO No. RAPE 23.1975;174-185
3. Upadhyay BP. Rice Disease Survey Report1977.In: *The proceeding of the fifth rice improvement workshop*, NRIP, Nepal. 1978; 65.
4. Manandhar HK, Thapa BJ, Amatya, P. Bacterial blight (BB) in hilly regions of Nepal. *International Rice Research Newsletter (Philippines)*. 1987.
5. Adhikari TB, Mew, TW .Bacterial blight (BB) resistance in some Nepal rice cultivars. *International Rice Research Newsletter (Philippines)*. 1989.
6. Thapa BJ. Report on rice pathology research work at Khumaltar. In: *The proceeding of the Eighth Rice Improvement Workshop*, NRIP, Parwanipur. 1981.
7. Burlakoti RR, Khatri-Chhetri GB. Bacterial diseases of crop plants in Nepal: a review. *Journal of the Institute of Agriculture and Animal Science*. 2004; 25: 1-10.
8. Chaudhary B, Yadav M, Akthar T, Yadav RB, Gharti DB, Bhandari D. Evaluation of Rice genotypes for resistance to Bacterial Blight. *Rice Research in Nepal Proceeding of 24th summer crop workshop*. Hardianth,Nepal: NARC,National Rice Research Programme.2004; 344-349.
9. Adhikari TB, Mew TW, Leach JE.Genotypic and pathotypic diversity in *Xanthomonasoryzaepv.oryzae* in Nepal. *Phytopathology*. 1999; 89(8):687-694.
10. Adhikari TB, Mew TW, Teng PS. Phenotypic diversity of *Xanthomonas oryzae p. oryzae* in Nepal. *Plant Disease*.1994;78,68-72.
11. Adhikari TB, Cruz C., Zhang Q, Nelson RJ, Skinner DZ, Mew TW, Leach JE. Genetic Diversity of *Xanthomonas oryzae pv.oryzae* in Asia. *Applied and Environmental Microbiology*. 1995; 61(3), 966-971.
12. Adhikari TB, Basnyat RC, Mew TW. Virulence of *Xanthomonas oryzae pv.oryzae* on rice lines containing single resistance genes and gene combinations. *Plant disease*. 1999; 83(1), 46-50.
13. Adhikari, T. B. Assessment of near-isogenic rice lines for resistance to *Xanthomonas oryzae pv. oryzae* in Nepal. *International journal of pest management*. 1993; 39(3), 293-296.

14. Adhikari TB, Mew TW. Resistance of rice to *Xanthomonas oryzae* pv. *oryzae* in Nepal. *Plant disease*. 1994; 78(1), 64-67.
15. Bhandari D. Evaluation of level of Resistance in Paddy genotypes against Bacterial Leaf Blight under Bhairahawa conditions. *Summer Crops Research in Nepal Proceedins of The 25th Summer Crops Workshop* Hardinath, Nepal: National Rice Research Programme. 2007: 455-459.
16. Bhatta MR, Sah BP, Khatiwada SP, Amgai R .Molecular screening on rice and barley: a gateway to marker assisted selection. In Marker Assisted Screening on Cereal in Nepal. *Proceedings of the Research Papers Presented at Special Session-Marker Assisted Breeding during Working Group Meeting on Plant Breeding and Genetics from 24 to 25 April 2012 at Khumaltar, Lalitpur, Nepal*. 2012; 45-45.
17. Amgai RB, Niroula RK, Pantha S, Hamal SS, Tamang BG, Sah BP, Bhatta MR. Marker-assisted screening of Nepalese rice for bacterial leaf blight (BLB) resistance. *Nepal Journal of Biotechnology* . 2015 ;3(1):35-39.
18. Sah SN, Yadaw RB . Status of Rice Research and Development in Nepal. *Proceedings of the 27th National Summer Crops Workshop* Kathmandu, Nepal: Nepal Agricultural Research Council. 2013; 2: 433-442.
19. Dangal NK, Chaudhary B, Joshi P, Sherpa AT. Evaluation Of Rice Genotypes Against Bacterial Leaf Blight Under Field Conditions. *Proceedings of the 27th National Summer Crops Workshop* Kathmandu, Nepal: Nepal Agricultural Research Council. 2013; 1: 313-318.
20. Khadka RB, Adhikari S, Shrestha G. Bacterial Leaf Blight of Rice in Mid and Farwestern Nepal: Preliminary Study of Disease. *Proceedings of the 27th National Summer Crops Workshop* Kathmandu, Nepal: Nepal Agricultural Research Council. 2013;1: 298-303.
21. Shakya DD, Manandhar S. Bacterial sheath brown rot of rice caused by *Pseudomonas fuscovaginae* in Nepal. In: *Plant Pathogenic Bacteria* (M Lemattre, S Freigoun, K Rudolph and JG Swings, eds), (Eds). INRA, Paris (Les Colloques, no. 66). 1992: 73-78.
22. Sharma S, Sthapit BR, Pradhanang PM and Joshi KD. Bacterial sheath brown rot of rice caused by *Pseudomonas fuscovaginae* in Nepal. *Lumle Agriculture Research Center* 98/5. 1995;2-10.
23. Manandhar G. *25 years of maize research in Nepal (1972- 1997)*. Nepal Agricultural Research Council, National Maize Research Program, Rampur, Chitwan, Nepal; 1997.
24. Subedi S. A review on important maize diseases and their management in Nepal. *Journal of Maize Research and Development*. 2015;1(1), 28-52. DOI: 10.5281/zenodo.34292
25. Shah SM. Diseases of maize in Nepal. In: Fifth Inter-Asian corn improvement workshop, Department of agriculture and Kesetsart University, Bangkok, Thailand. 1968.
26. Adhikari D, Baidya S, Koirala DK. Citrus Greening Test on Sweet Orange (Junar) by Scratch Method at Sindhuli District. *Journal of the Plant Protection Society*. Plant Protection Society, Nepal. Vol.4. 2012; 263-268
27. Thrower LB. Report on visit to Nepal. *FAO Report* PL, T51, 1968.
28. Knorr LC, Shah SM, Gupta OP. Greening disease of citrus in Nepal. *Plant Report*. 1970;54(12):1092-1095.
29. Regmi C, Yadav BP. Present status of Huanglongbing in western districts of Nepal. *Proc. 4th Hort. Seminar, Jan 18-19, 2007, Kirtipur, Kathmandu, Nepal, Nepal Horticulture Society*. 2007:40-43.
30. Regmi C. Mycoplasma-like disease of citrus in Nepal and USSR. Ph. D. Dissertation. Moscow Agriculture Academy, USSR. 1982.
31. Regmi C, Devekota RP, Paudyal KP, Shrestha S, Ayers AJ, Murcia N, Bove JM, Duran VN. Shifting from seedling mandarin trees to grafted trees and controlling huanglongbing and viroids: a biotechnological revolution in Nepal. In: *Insect-Transmitted Procaryote. Proceedings of the 17th Conference of IOCV, Adana, Turkey (ME Hilf, LW TFimmer, RG Milne and JV da Graçaeds)*. 2010.

32. Paudyal KP. Technological Advances in Huanglongbing (HLB) or Citrus Greening Disease Management. *Journal of Nepal Agricultural Research Council*. 2016; 1:41-50.
33. Jetter KM, Civerolo EL, Sumner DA. Ex ante economics of exotic disease policy: citrus canker in California. *Exotic Pests and Diseases: Biology, Economics and Public Policy for Biosecurity*. 2000:121-49.
34. Goto M. "Citrus Canker." In J. Kumar, H.S. Chaube. U.S. Singh, and A. N. Mukhopadhyay (eds.), *Plant diseases of International Importance*. Englewood Cliffs: Prentice Hall. 1992;3: 170-208
35. Dhakal D, Regmi C, Basnyat SR. Etiology and control of citrus canker disease in Kavre. *Nepal Journal of Science and Technology*. 2009;10: 57-61.
36. Shrestha SK. Preliminary study on brown rot of potatoes in Nepal. *Nepalese Journal of Agriculture*. 1977;12:11-21
37. Adhikari TB, Manandhar JB, Hartman GL. Characterisation of *Pseudomonas solanacearum* and evaluation of tomatoes in Nepal. In *ACIAR PROCEEDINGS AUSTRALIAN CENTRE FOR INTERNATIONAL AGRICULTURAL RESEARCH*. 1993: 132-132
38. Adhikari TB, Basnyat RC. Effect of crop rotation and cultivar resistance on bacterial wilt of tomato in Nepal. *Canadian journal of plant pathology*. 1998;20(3), 283-287.
39. Timila RD, Joshi, S. Participatory Evaluation of Some Tomato Genotypes for Resistance to Bacterial Wilt. *Nepal Agriculture Research Journal*. 2014;8: 50-55.
40. Lamichhane JR, Balestra GM, Varvaro L. First report of bacterial spot caused by *Xanthomonas campestris* pv. *vesicatoria* race 2 on tomato in Nepal. *New Disease Reports*, 2010;22:25. [[doi:10.5197/j.2044-0588.2010.022.025](https://doi.org/10.5197/j.2044-0588.2010.022.025)]
41. Lamichhane JR, Kshetri MB, Mazzaglia A, Varvaro L, Balestra GM. Bacterial speck caused by *Pseudomonas syringae* pv. *tomato* race 0: first report in Nepal. *Plant Pathology*. 2010; 59(2): 401-401.
42. Lamichhane JR, Balestra GM, Varvaro L. Responses of commonly cultivated tomato cultivars in Nepal to bacterial speck. *Phytopathologia Mediterranea*. 2011;49(3): 406-413.
43. Yabuuchi E, Kosako Y, Yano I, Hotta H, Nishiuchi Y. Transfer of two *Burkholderia* and an *Alcaligenes* species to *Ralstonia* gen. nov., proposal of *Ralstonia pickettii* (Ralstonia. Palleroni and Doudoraff 1973) comb. nov., *Ralstonia solanacearum* (Smith 1896) comb. nov. and *Ralstonia entomophila* (Davis 1969) comb. nov. *Microbial Immunol*. 1995;39: 897-904.
44. Pradhanang PM, Pandey RR, Ghimire SR, Dhital BK, Subedi A. An approach to management of bacterial wilt of potato through crop rotation and farmers participation. In: *Bacterial wilt* (GL Hartman and AC Hayward, eds.). *Proceedings of an International Conference held at Kaohsiung, Taiwan*. 1993:362- 370.
45. Shrestha SK. Occurrence and spread of bacterial wilt disease of potatoes with its effect on seed potato production in Nepal. In: *Proceedings of National Conference on Science and Technology*. 1988;213-219.
46. Adhikari TB. Identification of biovars and races of *Pseudomonas solanacearum* and sources of resistance in tomato in Nepal. *Plant Disease*. 1993; 77:905-907.
47. Dhital SP, Thaveechai N, Shrestha SK. Characteristics of *Ralstonia solanacearum* strains of potato wilt disease from Nepal and Thailand. *Nepal Agriculture Research Journal*. 2001; 4: 42-47.
48. Dhital SP. Bacterial wilt of potato in Nepal. In *Integrated Management of Bacterial Wilt. Proceedings of an International Workshop held in New Delhi, India*. 1995 :41-50.
49. Dhital SP, Thaveechai N, Kosiratana W, Piluek K, Shrestha SK. Effect of chemical and soil amendment for the control of bacterial wilt of potato in Nepal caused by *Ralstonia solanacearum*. *Kasetsart Journal, Natural Sciences*. 1997;31(4): 497-509.
50. Hogers CH, Shrestha SK. Control of brown rot of potato with crop rotation in Kathmandu valley in Nepal [infection diseases of potato, upland valley]. *Journal of the Institute of Agriculture and Animal Science (Nepal)* 1982..

- 426 51. Lamichhane JR, Balestra GM, Varvaro L. Occurrence of potato Soft Rot caused by *Erwinia*
427 *carotovora* (synonym *Pectobacterium carotovorum*) in Nepal: A First Report. *Plant Disease*.
428 2010;94(3): 382-382. <https://doi.org/10.1094/PDIS-94-3-0382C>
- 429 52. Chaurasia PCP, Chaudhari DK, Bhattarai DP. Management of common scab (*Stereptomyces*
430 *scabies*) of potato in 1998/99. 1999.
- 431 53. Timila RD. Seed-borne Infection of *Xanthomonas campestris* pv. *Campestris* in Cabbage and
432 its Control through Seed Treatment. *Nepal Journal of Science and Technology*. 2015; 3(1).
- 433 54. Jensen BD, Vicente JG, Manandhar HK, Roberts SJ. Occurrence and diversity of
434 *Xanthomonas campestris* pv. *campestris* in vegetable Brassica fields in Nepal. *Plant Disease*.
435 2010;94(3):298-305.
- 436 55. Neupane RK, Sah RP, Darai R. Varietal investigation on soybean in terai region of Nepal. In 23.
437 *National Summer Crops Research Workshop (Grain Legumes), Khumaltar, Lalitpur (Nepal), 2-*
438 *3 Jun 2002*. National Grain Legumes Research Program. 2003.
- 439 56. Gharti DB, Darai R, Banstola BS. Effect of Level of Field resistance To Major Diseases In
440 Available Genotypes Of Soybean At Nglrp, Rampur During 2005-2007. *Summer Crops*
441 *Research in Nepal Proceedins of The 25th Summer Crops Workshop* Hardinath, Nepal:
442 National Rice Research Proramme. 2007: 402-409.
- 443 57. Manandhar JB, Sinclair JB. Occurrence of soybean diseases and their importance in
444 Nepal. *FAO plant protection Bulletin*. 1982; 30(1):13-16.
- 445 58. Lamichhane JR, Balestra GM, Varvaro L. Bacterial leaf spot caused by *Xanthomonas*
446 *cucurbitae* reported on pumpkin in Nepal. *New Disease Reports*. 2010;22:20.