

1 **Biological Invasion of Tomato Leaf Miner, *Tuta absoluta* (Meyrick) in Nigeria:**
2 **Problems and Management Strategies Optimization: A review**

3
4
5
6
7
8

ABSTRACT

Nigeria is the largest producer of tomato in Africa while the Northern regions are the main producers of tomatoes in the country. Tomato (*Lycopersicon esculentum* Mill.) is widely cultivated and consumed in every part of Nigeria. It is an important commercial crop and good source of vitamins, minerals, essential amino acids, sugar and dietary fibers. The study aimed to report the outbreak of Tomato leafminer (*Tuta absoluta*) in tomato growing regions of Nigeria with a view to bring out its management strategies in Nigeria. The study was conducted in the major tomato producing regions which lies between latitudes 7.5o 11' and 13.0 ° N and a temperature range of 22 - 30°C. *Tuta absoluta* has been attributed to causing tomato shortage in Nigeria as a result of its devastating attack. The outbreak caused critical reduction in fruit yield which resulted in high cost of tomato. The damage is done by the larvae which tunnel into the stems, apical buds as well as the green and ripe fruits reducing their quality and subsequently the yields. Nonetheless, about 80 -100% yield loss had been reported. *Tuta absoluta* was reported to have developed resistance to a number of pesticides used in the country. Meanwhile, the Government had either less concern or neglect on the pest invasion or areas of researches especially on invasive species, thus poor funding which over stresses the existing challenges in controlling the pest. This paper intends to review some important highlights of Integrated Pest management (IPM) approaches for effective and efficient management strategies of the pest in Nigeria and beyond.

9
10
11
12
13

Keywords: [Tomato in Nigeria; *Tuta absoluta*; Tomato leaf miner; Tray Trap Technique; NIHORT-Lyptol and NIHORT-Raktin.]

14 **1. INTRODUCTION**

15
16
17
18
19
20
21
22
23
24

Nigeria is the largest producer of tomato in Africa and significantly contributes to the production of other major agricultural commodities [1]. Agro-ecological regions within the country are diverse and different regions possess environmental advantages in terms of optimum climatic and soil adaptability for the production of specific crops. The Northern regions are the main producers of tomatoes [2, 3]. It is an important component of daily diet, consumed fresh and in paste form. In Nigeria, tomatoes grow well in northern states of the country including Kano, Katsina, Jigawa, Zamfara, Bauchi, Sokoto and Taraba states. However, Kano state has the most suitable comparative advantages for tomato production in commercial scale [1]. The production is beset with many biotic and abiotic stresses such as

* E-mail address: nbszango@yahoo.com.

diseases, nematode and insect pests, thus resulting in reduced yield. Smallholder farmers cultivating 0.5 to 4 hectares of land account for about 90% of tomato production in the country, with the remaining (10%) being supplied by large-scale producers [1]. Nigeria has the largest area of land harvested for fresh tomato in Africa with 541, 800ha followed by Egypt with 214, 016ha [2], and the country remains the largest producer of tomato in Africa [3]. Unfortunately, huge amount of the crop is lost annually due to post-harvest losses, poor handling and lack of modern storage and processing facilities [4]. In 2014 for instance, Nigerian farmers averagely generated a significantly lowest yield of 4.0m tons/ha of tomato as compared to the quantity produced by Egypt (38.7m tons/ha) and South Africa (78.7m tons/ha) [2].

35

1.1 Invasive/introduced species

The invasive species are non-native that significantly modify or disrupt new ecosystems colonized. They may arrive in new areas through natural migration but are often introduced by human activities and/or other species [3]. In 2016, a sporadic attack by an invasive tomato leaf miner, *T. absoluta* was reported to cause more than 80% yield loss in Nigeria [3]. This had caused price hikes of up to 400% in three months as *T. absoluta* destroyed the annual harvest affecting entire tomato farms across northern states in the country (Fig. 1). The incident resulted in tomato shortage to the then newly commissioned Africa largest tomato processing company (Dangote Farms Tomato Processing Factory in Kano, Nigeria). Nonetheless, *Tuta absoluta* was resurfaced in 2017 and spreads to the north Eastern states especially Gombe state [5].

47

According to the Federal Ministry of Agriculture and Rural Development (FMARD) Nigeria currently produces 1.8 million tons annually, signifying that the output lost from the pest invasion is equivalent to 720, 000 metric tons (MT). However, the root cause of *T. absoluta* introduction to Nigeria is unknown and had become the most devastating pest with severe destructions in tomato producing areas [6]. Huge economic losses and rapid spread of the

* E-mail address: nbszango@yahoo.com.

devastating pest had been recorded in recent years. Complete eradication of the pest might be possible by the adoption of Integrated Pest Management (IPM) strategies. These involve the use of modern methods such as Tray Trap Technology (TTT) and biological control methods such as use of parasitoids and predators as well as minimized synthetic chemicals usage. Recently, National Horticultural Research Institute, Nigeria (NIHORT) developed and tested the efficacy of three bio-pesticides techniques: NIHORT-Raktin, NIHORT-Lyptol, and O-Ruptur. Similarly, gene silencing methods using RNA interference (RNAi) technology had also demonstrated potentials in controlling insect pests [7]. The present study had reviewed the ecology; biology and potential management strategies of *T. absoluta* for the implementation of efficient, effective and sustainable IPM strategies in Nigeria and elsewhere.




Fig. 1. Tomato Prices during 2015 Outbreak (Naira per Basket)

Source: [9] Sanda et al, 2017

3. RESULTS AND DISCUSSION

3.1 GEOGRAPHICAL DISTRIBUTIONS

The Tomato leafminer, *T. absoluta* (Meyrick) (Lepidoptera: Gelechiidae) is native to South America (Peru) and was declared a major pest of tomato in 1964 in Argentina [8, 10]. It was first noticed in Spain in 2006; since then, *T. absoluta* spread across Europe, Middle East and continues to spread rapidly across African countries and is found everywhere across the world including Nigeria. In Nigeria, it is commonly called “tomato Ebola or Sharon”. (Table 1) [20].

Table 1: The world's geographical distribution of *Tuta absoluta*

S/N0	Region	Country	Year Introduced	Reference
1	South America	Brazil (Peru)	Native	[8, 10]
2	South America	Argentina	Declared pest 1964	[8, 10]
3	Europe	Spain	2006	[10]
4	North Africa	Algeria	2008	[11, 12]
5	North Africa	Morocco	2009	[11, 12]
6	North Africa	Egypt	2010	[13]
7	West Africa	Sudan/Southern Sudan	2011	[14, 15]
8	East Africa	Ethiopia	2012	[16]
9	East Africa	Kenya	2013	[17]
10	East Africa	Tanzania	2014	[18]
11	West Africa	Senegal	2014	[19]
12	West Africa	Nigeria	2015	[20]
13	West Africa	Niger Republic	2016	[20]

83

84 In Nigeria, the major tomato producing areas lie between latitudes 7.5° 11' and 13.0 ° N,
85 within a temperature range of 22-30°C (Fig. 3). Tomato leafminer had become widespread

* E-mail address: nbszango@yahoo.com.

86 and an important pest throughout the regions [21]. According to the Ministry of Agriculture
87 and Rural Development of the Federal republic of Nigeria six tomato producing states
88 inclusive of Kano, Katsina, Jigawa, Gombe, Plateau and Abuja had been adversely affected
89 by the pest in 2017 [24]. [Channels TV news station on 24 May 2016].

90

91 **3. BIOLOGY AND LIFE CYCLE**

92 *Tuta absoluta* is an important invasive pest of tomato and a noctuid moth that is almost
93 always active at night. It mates and lays eggs during the night period. Adults (Figure 2a) are
94 spackle brown or silver-colored with black spotted wings. The adults are often 5-7 mm long,
95 with wing length of 8-10 mm. *T. absoluta* is a holometabolous insect with high reproduction
96 rate. Females can lay up to 260 - 300 eggs leading to 12 generations per year depending on
97 environmental conditions [12, 4].

98



99

100 **Fig. 2.** Adult (A) and larvae (B) of tomato leafminer

101 **Source:** [25, 26]

102



Fig. 3. Distributions of *Tuta absoluta* in Tomato Growing Areas in Nigeria

Source: [3, 27, 28]

They undergo complete metamorphosis involving four developmental stages which include egg, larva, pupa and adult stages that are completed within 24-30 days at favorable environmental conditions [23, 24]. The newly emerged female adults release sex pheromone that attracts the male during mating [25, 27]. Females usually deposit their eggs (which are elliptical in shape, measuring up to about 1mm) on tomato leaves within 7 days of mating. The egg color varies from creamy white to bright yellow. The emerging larvae burrow into the leaves where they feed on the mesophyll tissues and develop within 7-11 days [28]. The larvae (caterpillars) (Fig. 2b) appear creamy in the first stage before turning greenish and pinkish in the final stage. They measure from 0.6mm to 0.8mm in length in their first larval stage and 7.3mm to 8mm in the fourth stage [25]. The caterpillars possess two narrow black bands on the head, one lateral and the other ventral. The pupa (Chrysalis) measures 4mm

119 to 5mm in size and is initially green in color before turning brown. Pupation lasts for 10-11
120 days in soil to get rid of chemicals and strong heat. However, *T. absoluta* was reported to
121 exhibit different pupation behaviors depending on the environment and can survive the
122 winter as egg, larva or adult depending on the environmental conditions [25].

123

124 **4. IMPACTS OF *Tuta absoluta* ON TOMATO PRODUCTION IN NIGERIA**

125 In Nigeria, *T. absoluta* was first reported in 2015 in the Arumeru district and later discovered
126 in Kadawa irrigation valley of Kano state, the major tomato producing area in the country and
127 where the famous Dangote's Dansa Tomato Company is sited. The pest inflict its damage at
128 the larval instar stage and it entirely attacks the tomato plant thus damaging all of the
129 productive plant parts [29] (Fig. 4). Specifically, the larvae penetrate the apical buds,
130 flowers, fruits, leaves and/or stems immediately after hatching. Leaf damages are often due
131 to the feeding habit on the mesophyll tissue which forms irregular mines on leaf surfaces and
132 later become necrotic. Moreover, sometimes the larvae form large galleries on the stem and
133 fruits which alter plants' general development and it also develop open wounds that serve as
134 entry points of pathogens leading secondary infections [30]. *T. absoluta* feeds on both green
135 and mature tomato fruits and can destroy the entire tomato vegetation within two days (Fig.
136 3) and the damage is always severe especially in young plants. Tomato is considered the
137 primary host for *T. absoluta*; however, it also attacks eggplants, peppers, potatoes, beans
138 and tobacco, hence the pest remains persistent in both cultivated and uncultivated areas for
139 long time.

140



141

Fig. 4. Symptoms of *Tuta absoluta* Damage on Leaf, Fruit and Stem of tomato

Source: [9] Sanda et al 2017

5. Management strategies of *Tuta absoluta*

Considering the vicious nature and crop destruction potential of *T. absoluta* in the invaded regions, there exist no feasible, effective, efficient and sustainable control method(s) that can effectively be used by resource poor farmers to manage pest in Nigeria. However, there various control strategies which if properly and effectively implemented can help manage *T. absoluta* in the country and beyond [6]. According to the National Horticultural Research Institute, Nigeria (NIHORT), the integrated pest management of *T. absoluta* must focused on realizing the following national goals: 1) to contain the economic damage of leafminer, 2) to prevent the development of resistant strains of *T. absoluta* and 3) to prevent the accumulation of insecticide residues in the food chain. Therefore, these management strategies can be categorized as follows; i) Chemical control ii) The use of pheromone traps and mass trapping (Tuta trap tray) ii) Cultural control iv) Biological control v) Resistant varieties and vi) Bio-pesticides.

5.1 CHEMICAL CONTROL

Chemical control of *T. absoluta* often fails not only due to the resistance of the pest against many pesticides, but also because a big part of its development takes place inside the plant or the soil; out of reach of pesticides. A resistance management strategy for *T. absoluta* is critical in the search for a sustainable solution to chemical control of *T. absoluta* in the medium. The new active ingredient which is likely to be the mainstay of the IPM programme is Chlorantraniliprole (Trade name: Rynaxypyr) from Dupont. At present, there are no records of resistance by *T. absoluta* to it and is known to be compatible with other natural enemies. Although a number of pesticides had been registered for the management of the pest, they are of low to moderate effectiveness due to the cryptic nature of the larvae and the high biotic potential of the insect. In addition, several cases of insecticide resistance

* E-mail address: nbszango@yahoo.com.

171 have been reported including resistance to organophosphates, Ampligo, Pyrethroids,
172 Abamectin, Chlorantraniliprole, Flubendiamide, Permethrin and Spinosad [31, 32, 33].

173 **5.2 PHEROMONE**

174 Sex pheromones are powerful chemicals secreted by female insect to attract the male
175 counterparts for mating. They can either be natural or synthetic. These chemicals are
176 detected by the males, assisting them in locating the females for mating. Many pheromones
177 from different species have been identified and are synthetically produced for use in insect
178 pest management. The majority of female sex pheromones identified in Lepidoptera consist
179 of a mixture of two or more compounds which not only evoke long-range male attraction but
180 also elicit courtship behavior [32,]. The female sex pheromone of *T. absoluta* for instance,
181 consists of two components. The major component, which represents about 90% of the
182 volatile material found in the sex gland of calling females was (3E, 8Z, 11Z)-3,8,11-
183 tetradecatrien-1-yl acetate or TDTA [33]. Sex pheromones have been widely used to monitor,
184 forecast or control populations of moth pests [34]. The most widespread and successful
185 application of sex pheromones is that used in the detection and population monitoring [35].

186 The insect populations could be controlled or monitored by two main techniques; the mass
187 annihilation and mating disruption techniques [35]. Sex pheromones management
188 strategies are based on the sexual reproduction of the targeted pest. However, a study by
189 [36, 37, 38] recently confirmed that *T. absoluta* females were also able to reproduce without
190 mating (*i.e.* parthenogenically). Asexual reproduction as well as the polygenic nature of
191 *T. absoluta* males could have strong implications on the efficiency of sex pheromones
192 management strategies and must be considered imperative for further studies especially in
193 *T. absoluta* management [37, 38]. Although the female sex pheromone of *T. absoluta* was
194 identified in 1995, virgin females were already used to capture more than 100 males per trap
195 per day [39]. According to [40] virgin females could be used to compare different trap
196 designs, heights, and displacement in tomato fields for capturing *T. absoluta* males. A high
197 specificity and sensitivity of traps baited with virgin females was recorded and they seemed

198 to be more economical and convenient than light traps. Moreover, wind tunnel experiments
199 under laboratory conditions indicated that the mixture of the two pheromone components
200 was more effective in evoking long-range female location and short-range courtship behavior
201 in males than TDTA alone [37]. However, it has been observed in laboratory experiments
202 that *T. absoluta* males were far less sensitive to the absence of the minor component than
203 most other Lepidopterans [40]. To monitor *T. absoluta*, pheromone lures are principally
204 coupled with Delta traps [41, 42]. Various companies such as ISCA Technologies (United
205 States), Russell IPM Ltd. (United Kingdom), Koppert Biological Systems (The Netherlands),
206 and PRI Pherobank (The Netherlands) had manufactured different kind of these traps for *T.*
207 *absoluta* management [29].

208

209 5.3 TUTA TRAP TRAY AND TRAY TRAP TECHNOLOGY (TTT)

210 Syngenta Bioline conducted trials in 2013 to test Black Sticky Traps as a new tool for
211 capturing *T. absoluta* in tomato crops and high capture rates were achieved [43], however,
212 correct placement of the traps is critical for successful captivity. Traps placed at 15-20cm
213 above the ground captured 5 times as many adult pests as ones placed at 1m above the
214 ground. Interestingly, the traps do not attract predators of the *T. absoluta* such as
215 *Nesidiocoris spp* [43]. Similarly, the National Horticultural Research Institute, Nigeria
216 (NIHORT) developed simple, effective, efficient and sustainable technology known as Tray
217 Trap Technology (TTT) that can attract, trap and kill more than five thousand adults per tray
218 [9]. It was the only control measure employed and recorded massive successes in curtailing
219 the menace of leafminer pest in Nigeria. The tool is sold to local farmers at affordable rates
220 within the outbreak areas. Moreover, the technology consists of a tray filled with water
221 containing small amount of detergent (without foam). A lamp is placed on a solid support at
222 the center but not completely dipped into the water (Fig. 5). The whole system is placed in
223 the farm but away from canopies and the lamp attracts adult *T. absoluta* pests during nights.
224 The adults flap around the light and suddenly fall into the solution and remain captive.

225

226 5.4 BIOLOGICAL CONTROL MEASURES

227 Predators of *T. absoluta* are present in Africa but may not be found in the field in sufficient
228 number at the time when the adult leafminer moths inflict damage on the crop and to begin
229 to lay eggs. For example, predatory mites, bugs as well as parasitic wasps usually migrate
230 into the crop in high number only when there is a lot of the specific prey present. This may
231 be too late to prevent damage from occurring.



232
233
234
235

Fig. 5. Tray Trap Technique (TTT) in Nigeria

Source: [22]National Horticultural Research Institute, Ibadan, Nigeria (NIHORT)

236 Nonetheless, to tackle this problem, predators could be reared in mass and released prior to
237 domination by the destructive pests. Several scientific reports [44, 45, 46] had established
238 the importance of the control agents in the management of the epidemics invasive species to
239 a significantly lower threshold levels. However these control agents might be severely
240 affected with improper planning especially when chemical control methods are the ultimate
241 options. Pyrethroids (e.g. Alphacypermethrin, deltamethrin, cypermethrin etc.) can kill up to
242 75% of the population of natural enemies and the harmful effect could last for up to 12
243 weeks after a single spray. Massive production of the bio-agents could be expensive;
244 therefore the grower must be aware of the importance of using compatible pesticides that
245 are feasible, cost effective and environmentally friendly. There are several bio-agents that
246 are currently used for the control of *T. absoluta* among which are: a) *Trichogramma*

247 *pretiosum* b) *Trichogramma achaeae* c) *Macrolophus pygmaeus* d) *Nesidiocoris tenuis* e)
248 *Nabis pseudoferus* f) *Amblyseius* spp.

249 **5.4.1 *Trichogramma achaeae***

250 In South American the egg parasitoid wasp, *Trichogramma achaeae* had been identified as
251 agent for biological control of the Tomato Pinworm, *T. absoluta*. In a greenhouse research,
252 higher efficacy of 91.74 % damage reduction was obtained when 30 adults were released
253 per plant (equivalent to 75 adults/ m²) in every 3 - 4 days in south-eastern Spain [45].
254 Moreover, biological control of *T. absoluta* using the damsel bug, *Nabis pseudoferus*, has
255 also been conducted in Spanish greenhouses [45]; while two semi field bioassays on tomato
256 plants under controlled conditions had shown significant reduction in the number of eggs of
257 *T. absoluta* between 92 and 96 %, when 8-12 first stage nymphs of *N. pseudoferus* were
258 released per plant [45]. *Trichogramma* is normally introduced onto crops as an irradiated
259 *Ephestia kuehniella* caterpillar egg parasitized by the *Trichogramma* wasp. These wasps are
260 extremely sensitive to broad-spectrum chemicals and can only be used in spray
261 programmes that are carefully designed to conserve *Trichogramma* parasitoids. Moreover,
262 *Trichogramma* parasitoids can be used in combination with *M. pygmaeus* for effective
263 biological control of *T. absoluta* on tomatoes.

264 **5.4.2 *Amblyseius* spp**

265 *Amblyseius swirskii* is a beneficial predatory mite endemic to the Eastern Mediterranean
266 regions. This species is considered generalized predator and readily predate small soft-
267 bodied arthropod pest species as well as pollen or plant exudates. *A. swirskii* has attracted
268 substantial interest as a biological control agent of mites, thrips and whiteflies in greenhouse
269 and nursery crops and is currently reared and sold commercially in Europe and North
270 America [46]. Both *Amblyseius swirskii* and *A. cucumeris* had been observed feeding on

271 very young *T. absoluta* caterpillars. In laboratory studies, these and other predatory mites
272 had been observed feeding on the eggs of *T. absoluta*, though they could not complete their
273 life cycle on leafminer eggs alone [48, 49]

274 **5.4.3 Nesidiocoris spp**

275 *Nesidiocoris tenuis* (Reuter) (Hemiptera: Miridae) is an effective predator of pests of tomato
276 crops and a promising bio-control agent of *T. absoluta* in the Mediterranean. *N. tenuis* has
277 catholic habits, feeding on almost every small arthropod dwelling on tomato plants [50]. This
278 bug is indigenous to Kenya and is commercially reared and has been widely used in Spain
279 against *T. absoluta* in tomato farms [50]. It has also been tested in south eastern Algeria.
280 However, one should be cautious when using them as bio-control agents; because they are
281 capable of causing serious economic damage to tomato plants if they remain in large
282 number after a successful targeted control. Natural migration of *N. tenuis* to tomato
283 vegetation is unlikely to be of any risk to the crops due to a much lower population of bugs
284 as compared to their artificially large inundatory release.

285

286 **5.4.4 Macrolophus spp**

287 *Macrolophus pygmaeus* and *M. caliginosus* had been reported feeding on young leafminer
288 caterpillars. *Macrolophus* can be a common insect in natural vegetation and has been
289 recorded as becoming established inside tomato greenhouses in Northern Europe [51].
290 Moreover, both *Nesidiocoris* and *Macrolophus* had been observed to prey on *T. absoluta*
291 immediately after its detection in the Spanish Mediterranean coastal region [52, 53]. *M.*
292 *pygmaeus* and *N. tenuis* have already been tested under laboratory conditions to assess
293 their suitability as *T. absoluta* predators [54]. Adults were able to prey on more than 100
294 eggs per individual per day with no significant differences between them. In contrast,
295 nymphs of *M. pygmaeus* consumed significantly fewer eggs than that of *N. tenuis* [53].
296 Furthermore, both species prey on the larvae at all larval stages with greater preference for

297 first- instar larvae [45, 55], although the number of larvae preyed upon was significantly
298 lower than that of eggs. Also [45] reported that when *M. pygmaeus* and *N. tenuis* were
299 established in the crop, they were able to reduce up to 75% and 97% of leaflet infestations
300 or 56% and 100% of fruit infestations respectively.

301 **5.4.5 *Bacillus thuringiensis***

302 *Bacillus thuringiensis* var. *kurstaki* has exhibited satisfactory efficacy against *T. absoluta*
303 larval infestations in Spanish outbreaks. The bacterium ingest a lot of leafminer larva to a
304 lethal dose while delayed application of *B. thuringiensis* may cause higher insect mortality if
305 the insects become more susceptible to the pathogen after a longer period of feeding on the
306 resistant crop [55]. [56] reported that in a combined application of mass release of *T.*
307 *pretiosum* Riley and *B. thuringiensis* resulted in fruit damage only by 2 % in South America.
308 Therefore the only life stage that is likely to be affected by *B. thuringiensis* is the first instar,
309 free -living stage. Due to many over-lapping generations of *T. absoluta* this will require a
310 very intensive spray programme to provide sufficient coverage of the vulnerable life stage.
311 This raises questions of resistance management and cost-effectiveness of the programme
312 [56]. *B. thuringiensis* is proved to be highly efficient in reducing the damage produced by the
313 first, second, and third *T. absoluta* larval instars with statistically significant differences
314 between controls and treated plants in all cases.

315 **5.4.6 ENTOMOPATHOGENIC NEMATODES (EPN) AND FUNGI**

316 Insect killing nematodes are massively and commercially produced and have been used in
317 high value protected crops against a range of pests. Entomopathogenic nematodes have
318 been tested for the management of *T. absoluta*. Both Laboratory and field experiments had
319 revealed high larval mortality (78.6-100%). However, low pupal mortality (10%) had been
320 recorded when *Steinernema feltiae* was evaluated against the pest [57]. The efficiency of
321 Entomopathogenic fungi on *T. absoluta* has not been widely investigated although, several

322 fungal species including *Metarhizium anisopliae* and *Beauveria bassiana* were reported to
323 attack the eggs, larvae and adults of the Leafminer. Studies revealed up to 54% mortality of
324 *T. absoluta* adults by *M. anisopliae* [58] while the Entomopathogenic fungus *Metarhizium*
325 *anisopliae* could result in female mortality of up to 37.14%. Laboratory studies indicated
326 *Beauveria bassiana* caused 68% larval mortality [59, 60]. It was reported that complete
327 control of *T. absoluta* pupae in laboratory bioassays with another commercial *Metarhizium*
328 have been achieved [59]. Moreover, in Algeria similar results were reported [60] that the
329 eggs and early instars of *T. absoluta* were also readily infected and destroyed by *M.*
330 *anisopliae*. According to reports from Turkey compared the efficacy of *B. bassiana* and *M.*
331 *anisopliae* on Tuta eggs and first instar larvae; these entomo-pathogens were 42 –67% and
332 92 to 100% efficient in controlling tuta larvae, respectively. However, only *Metarhizium*
333 seemed to be effective against the tuta egg stage, providing 92% effective control, whilst
334 *Beauveria* provided only 12.5% control of Tuta eggs. These results show the potential of *M.*
335 *anisopliae* to control pupae of the tomato leafminer in integrated pest management
336 programs.

337 **5.4.7 BIO - PESTICIDAL CONTROL**

338 Azadirachtin is a complex of Tetranortri-terpenoidlimonoid from Neem seeds and is a major
339 active component responsible for both anti-feeding and toxic effects in insects. Other
340 limonoid and sulphur-containing compounds with repellent, antiseptic, contraceptive,
341 antipyretic and anti-parasitic properties were also found in other parts of the tree (leaves,
342 flowers, bark and roots) [61]. The Neem seed extract, Azadirachtin acts as a contact and
343 systemic insecticide against *T. absoluta*. A soil application caused 48.9 -100 % larval
344 mortality [62]. Application of Neem oil on the adaxial surface of the foliage causes 57-100%
345 larval mortality. However, it was reported that direct application on larvae caused 52.4-95 %
346 mortality [63]. Azadirachtin has a knockdown action on *T. absoluta* larvae while in a
347 laboratory test; aqueous Neem seed extract induced high larval mortality by both systemic

348 and trans-laminar actions [64]. It prevents oviposition by inhibiting oogenesis and synthesis
349 of ovarian ecdysteroid. In males, Azadirachtin interrupts the meiotic process responsible for
350 sperm production [65]. Nonetheless, a laboratory study reported that isolates of *B. bassiana*
351 high mortality to tomato leafminer eggs and larvae [59, 60] where the eggs appeared more
352 susceptible than the first instars larvae [58]. Thus, the use of Azadirachtin significantly
353 reduces *T. absoluta* population on tomato plants in the field. It is much more effective on first
354 and second instar larvae compared to the third and fourth instars.

355 Generally the efficacy in Azadirachtin is low during the first few days of application and later
356 suddenly increases until the 14th day, during which it exerts its maximum efficacy [65].
357 Recently, the National Horticultural Research Institute, Nigeria (NIHORT) developed three
358 new bio-pesticides for the control of *T. absoluta* at different stages (Fig. 6). These are the
359 NIHORT-Raktin, NIHORT-Lyptol and O-Ruptur. The NIHORT-Raktin is an emulsifiable
360 suspension from Neem tree that contains Azadirachtin as the primary active ingredient.
361 NIHORT-Raktin is a natural extract that controls such insects as *T. absoluta*, army worm,
362 aphids, biting and sucking insects. The NIHORT-Raktin is sprayed on tomato leaves after a
363 dilution of 1 litre NIHORT-Raktin in 15 liters of water. NIHORT-Lyptol is another emulsifiable
364 suspension from Eucalyptus tree with Cineole as the major active ingredient. It has powerful
365 poisoning effect against *T. absoluta* larvae after direct contact. It is also sprayed on tomato
366 leaves after a dilution similar to NIHORT-Raktin. O-Ruptur is a wettable concentrate
367 containing the active ingredient Tropine. It has contact poison against tomato leafminer
368 pupae. O-Ruptur can be applied at the rate of 250 g in 7 liters of water to an area of 4 m².

369 **5.5 PHYTOSANITARY MEASURES**

370 Preventive approaches are best techniques to counter leafminer epidemic as the pest is very
371 difficult to control. Sound farm sanitation reduces the pest survival rate since most eggs and
372 larvae are being destroyed. The selective removal and destruction of infested plant parts
373 and the total destruction of plants after harvest [66] as well as the management of alternate

* E-mail address: nbszango@yahoo.com.

374 wild hosts on the other hand. Establishment and maintenance of habitats within and
375 surrounding open-field grown tomato crops that supply food and provide shelter for
376 oviposition and overwintering encourage natural antagonists of *T. absoluta* [54].



377
378
379 **Fig. 6. NIHORT-Raktin (A), NIHORT-Lyptol (B) and O-Ruptur (C)**

380
381 *Source: [22] National Horticultural Research Institute, Ibadan, Nigeria (NIHORT)*

383 **5.6 RESISTANT VARIETIES**

384 Breeding tomatoes for *T. absoluta* resistance had so far been unsuccessful and so there are
385 no commercial tomato hybrids available that have an acceptable degree of resistance to *T.*
386 *absoluta* [67]. This is thought to be due mainly to a lack of genetic variability for this trait
387 within the gene pool used for domestic tomato production, resulting from long periods of
388 breeding/selection for other traits. However, to-date there seems limited or no clear
389 information as whether there are successful tomato varieties which are resistant to *T.*
390 *absoluta*. Efforts to develop resistant varieties are going on in different parts of the world and
391 this may need more research efforts to identify the adaptation mechanisms and areas of
392 weakness for effective control [20].

394 **5.7 GENETIC CONTROL USING RNA INTERFERENCE (RNAi)**

395 RNA interference (RNAi) technology has demonstrated an effective potential to control
396 insect pests. However, the efficiency of RNAi can vary greatly between the different insect
397 orders. It is a new and promising technique because interfering with insect RNA transcripts

* E-mail address: nbszango@yahoo.com.

398 using RNAi may lead to total death of the pests [68]. [8] reported the possibility of using
399 RNAi as an alternative method for controlling the tomato leafminer. They selected two target
400 genes (coding for Vacuolar ATPase-A and Arginine kinase) based on the RNAi response
401 reported for these genes in other pest species. The first was based on the uptake of dsRNA
402 by leaflets and the second was based on “in planta-induced transient gene silencing
403 (PITGS) method. It has been demonstrated that the *T. absoluta* larvae that fed on the leaves
404 containing *dsRNA* of the target genes had about 60% reduction in the accumulation of the
405 target gene transcript and an increased larval mortality resulting in less leaf damage [69].
406 Furthermore, they generated transgenic ‘Micro-Tom’ tomato plants that expressed hairpin
407 sequences for both transcripts and observed a reduction in foliar damage by *T. absoluta* in
408 those plants [8].

409

410 **6.0 CONCLUSIONS**

411 Currently, *T. absoluta* is present everywhere in Africa and Nigeria where tomato is grown;
412 the pest is a potential impending threat to tomato production that spreads across Africa.
413 Moreover, with all its damaging impacts on tomato production, there exist no effective,
414 efficient and sustainable management options that will help eradicate the pest. Multilateral
415 measures are proposed in controlling tomato leafminer which consider combination of
416 biological control methods, massive trapping and chemical or microbiological applications.
417 Others include legislative and agricultural extension approaches to convey practical
418 agricultural solutions to farmers; policy makers to establish appropriate control policies; and
419 farmers to adopt new innovations towards sustainable tomato production in Africa. The use
420 of breeding and molecular approaches in order to bring forward resistant varieties could be
421 the most sustainable control strategy. Moreover, the bio-pesticides (NIHORT-Raktin,
422 NIHORT-Lyptol and O-Ruptur. NIHORT-Raktin) of the NIHORT should be given wider
423 spread and be cost effective. Therefore, there is need for more collaborative efforts of the

European and African research scientists and the management specialists as well as tomato growers to tackle *T. absoluta* menace in Nigeria and beyond.

426

427 **7.0 COMPETING INTERESTS**

428

429 All authors have declared that no competing interests exist.

430

431 **8.0 AUTHORS' CONTRIBUTIONS**

432

433 1. Corresponding Author Sanda N. B.: initiated and designed the study, performed the
434 basic review, wrote the protocol, and wrote the first draft of the manuscript.

435 2. 'Wudil B. S.', Abdullahi A.M. and 'Sunusi M.': Critical review of the manuscript.

436 3. 'Hamisu H. S. and Sule H.: managed the literature searches and provided the field
437 information.

438 4. All authors read and approved the final manuscript.

439

440

441 **REFERENCES**

442

443 1. FAO [Food and Agriculture Organization of the United Nations]. (2014). FAOSTAT.
444 Available: <http://faostat3.fao.org/faostatgateway/go/to/download/Q/QC/E>. Accessed
445 2014 Feb 5.

446 2. Etebu E, Enaregha E. Postharvest quality of commercial tomato (*Lycopersicon*
447 *esculentum* Mill.) fruits brought into Yenagoa Metropolis from Northern Nigeria. J.
448 Biol. Agric. Healthcare. 2013; 3(11):2224-3208.

449 3. Borisade, O. AL, Kolawole, AO, Adebo, GM and Uwaidem, YI. Journal of Agricultural
450 Extension and Rural Development. The tomato leafminer (*Tuta absoluta*)
451 (Lepidoptera:Gelechiidae) attack in Nigeria: Effect of climate change on over-sighted
452 pest or agro-bioterrorism? 2017; 9(8): 163 – 171.

453 4. Gebremariam G. *Tuta absoluta*: a global looming challenge in tomato production,
454 review paper. J Biol Agric Healthc. 2015; 5(14):57-62.

455 5. Bello BO, Ullah H, Olawuyi O, Adebisi O. Microorganisms causing post-harvest
456 tomato (*Solanum lycopersicum* L.) fruit decay in Nigeria. Scientia. 2016; 13(2):93-6.

- 457 6. Cook DC, Fraser RW, Paini DR, Warden AC, Lonsdale WM, De Barro PJ.
458 Biosecurity and yield improvement technologies are strategic complements in the
459 fight against food insecurity. PLoS One. 2011 Oct 12; 6(10):e26084.
- 460 7. Borisade OA, Kolawole AO, Adebo GM, Uwaidem YI. The tomato leafminer (*Tuta*
461 *absoluta*) (Lepidoptera: Gelechiidae) attack in Nigeria: effect of climate change on
462 over-sighted pest or agro-bioterrorism? Journal of Agricultural Extension and Rural
463 Development. 2017 Aug 31; 9(8):163-71.
- 464 8. Germain JF, Lacordaire AI, Cocquempot C, Ramel JM, Oudard E. A new tomato
465 pest in France: *Tuta absoluta*. PHM Revue Horticole. 2009(512):37-41.
- 466 9. Sanda N. B., Hamisu, H. S. and You-ming Hou. Outbreak of an Invasive pest
467 Tomato leaf miner (*Tuta absoluta*) on Tomato (*Lycopersicon lycopersicum*)
468 productions in Nigeria: Paper presented at the 3rd International Congress on
469 Biological Invasion, Hanzhou, China. (2017, November, 19-23) (pp. 184)
- 470 10. Camargo RA, Barbosa GO, Possignolo IP, Peres LE, Lam E, Lima JE, Figueira A,
471 Marques-Souza H. RNA interference as a gene silencing tool to control *Tuta*
472 *absoluta* in tomato (*Solanum lycopersicum*). PeerJ. 2016 Dec 15; 4:e2673.
- 473 11. Edrees AE. Effect of Neem Seeds Ethanolic Extracts on Mortality of Tomato leaf-
474 miner *Tuta absoluta* (Lepidoptera: Gelechiidae) (Doctoral dissertation, Sudan
475 University of Science and Technology).
- 476 12. Pastrana JA, Braun K. Los lepidópteros argentinos: sus plantas hospedadoras y
477 otros sustratos alimenticios. San Miguel de Tucumán: Sociedad Entomológica
478 Argentina; 2004.
- 479 13. EPPO First record of *Tuta absoluta* in Algeria. EPPO Reporting
480 Services, 2008b. 7 (135). Accessed January 15, 2010. Available:
481 http://www.eppo.org/publications/reporting/reporting_service.htm

- 482 14. EPPO First record of *Tuta absoluta* in Morocco (2008/174). EPPO Reporting
483 Services, 2008c, 9(174). Accessed January 15, 2010. Available:
484 http://www.eppo.org/publications/reporting/reporting_service.htm
- 485 15. Moussa S, Sharma A, Baiomy F, El-Adl FE. The Status of Tomato leafminer, *Tuta*
486 *absoluta* (Meyrick)(Lepidoptera: Gelechiidae) in Egypt and potential effective
487 pesticides. Academic Journal of Entomology. 2013;6(3):110-5.
- 488 16. Pfeiffer DG, Muniappan R, Sall D, Diatta P, Diongue A, Dieng EO. First record of
489 *Tuta absoluta* (Lepidoptera: Gelechiidae) in Senegal. Florida Entomologist. 2013
490 Jun; 96(2):661-2.
- 491 17. Brévault T, Sylla S, Diatte M, Bernadas G, Diarra K. *Tuta absoluta* Meyrick
492 (Lepidoptera: Gelechiidae): a new threat to tomato production in Sub-Saharan
493 Africa. African Entomology. 2014 Jul; 22(2):441-4.
- 494 18. Goftishu M, Seid A, Dechassa N. Occurrence and Population Dynamics of Tomato
495 Leaf Miner [*Tuta absoluta* (Meyrick), Lepidoptera: Gelechiidae] in Eastern Ethiopia.
496 East African Journal of Sciences. 2014;8(1):59-64.
- 497 19. Mohamed ES, Mahmoud ME, Elhaj MA, Mohamed SA, Ekesi S. Host plants record
498 for tomato leaf miner *Tuta absoluta* (Meyrick) in Sudan. EPPO Bulletin. 2015 Apr
499 1;45(1):108-11.
- 500 20. Biondi A, Zappalà L, Desneux N, Aparo A, Siscaro G, Rapisarda C, Martin T, Tropea
501 Garzia G. Potential toxicity of α -cypermethrin-treated nets on *Tuta absoluta*
502 (Lepidoptera: Gelechiidae). Journal of economic entomology. 2015 Mar
503 21;108(3):1191-7.
- 504 21. Tonnang HE, Mohamed SF, Khamis F, Ekesi S. Identification and risk assessment
505 for worldwide invasion and spread of *Tuta absoluta* with a focus on Sub-Saharan
506 Africa: implications for phytosanitary measures and management. PloS one. 2015
507 Aug 7; 10(8):e0135283.

- 508 22. Abiola Oke Scientists at the National Horticultural Research Institute (NIHORT),
509 Ibadan have invented a technology to reduce the impact of tuta absoluta, known as
510 tomato ebola, 2015. Accessed May 6th, 2018. Available:
511 [https://www.dailytrust.com.ng/news/agriculture/new-tray-and-](https://www.dailytrust.com.ng/news/agriculture/new-tray-and-lamp-tech-to-end-tomato-ebola/189880.html)
512 [lamp-tech-to-end-tomato-ebola/189880.html](https://www.dailytrust.com.ng/news/agriculture/new-tray-and-lamp-tech-to-end-tomato-ebola/189880.html)
- 513 23. Haougui Adamou, Basso Adamou, Madougou Garba, Salissou Oumarou, Bague
514 Gougari, Moumouni Abou, Aissa Kimba and Patrick Delmas. Confirmation of the
515 presence of *Tuta absoluta* (meyrick) (lepidoptera: gelechiidae) in niger (west africa).
516 In. J. of Sc, Env. 2016. Vol. 5, No 6, 2016, 4481 – 4486
- 517 24. NAPPO. Surveillance Protocol for Tomato leaf Miner, *Tuta absoluta*, for NAPPO
518 Members Countries (2012).
- 519 25. Nigerian government attempts to contain *Tuta absoluta* spread by channelstv.com.
520 Accessed 24 May 2016. [https://www.channelstv.com/.../nigeria-home-grown-](https://www.channelstv.com/.../nigeria-home-grown-solution-tomato-pest-tuta-absoluta)
521 [solution-tomato-pest-tuta-absoluta](https://www.channelstv.com/.../nigeria-home-grown-solution-tomato-pest-tuta-absoluta)
- 522 26. Vercher R, Llopis VN, Porcuna JL, Marí FG. La polilla del tomate," *Tuta absoluta*".
523 Phytoma Espana: La revista profesional de sanidad vegetal. 2007(194):16-23.
- 524 27. *Tuta absoluta* Accessed January 20,
525 2018 https://en.wikipedia.org/wiki/Tuta_absoluta
- 526 28. Never Zekeya, Patrick A. Ndakidemi, Musa Chacha and Ernest Mbega Tomato
527 Leafminer, *Tuta absoluta* (Meyrick 1917), an emerging agricultural pest in Sub-
528 Saharan Africa: Current and prospective management strategies, Afr. J. Agric. Res.,
529 2017. Vol. 12(6), pp. 389-396, 9 February, 2017
- 530 29. Aynalem B. Tomato leafminer [(*Tuta absoluta* Meyrick) (Lepidoptera: Gelechiidae)]
531 and its current ecofriendly management strategies: A review. Journal of Agricultural
532 Biotechnology and Sustainable Development. 2018 Feb 28; 10(2):11-24.

- 533 30. Batalla-Carrera L, Morton A, García-del-Pino F. Efficacy of entomopathogenic
534 nematodes against the tomato leafminer *Tuta absoluta* in laboratory and
535 greenhouse conditions. *BioControl*. 2010 Aug 1; 55(4):523-30.
- 536 31. Harizanova V, Stoeva A, Mohamedova M. Tomato leaf miner, *Tuta absoluta*
537 (Povolny) (Lepidoptera: Gelechiidae)—first record in Bulgaria. *Agricultural science*
538 *and technology*. 2009; 1(3):95-8.
- 539 32. APHIS U. New pest response guidelines: tomato leafminer (*Tuta absoluta*).
540 Washington, DC: United States Department of Agriculture. 2011.
- 541 33. Siqueira HÁ, Guedes RN, Picanço MC. Insecticide resistance in populations of *Tuta*
542 *absoluta* (Lepidoptera: Gelechiidae). *Agricultural and Forest Entomology*. 2000 May
543 1;2(2):147-53.
- 544 34. Haddi K, Berger M, Bielza P, Cifuentes D, Field LM, Gorman K, Rapisarda C,
545 Williamson MS, Bass C. Identification of mutations associated with pyrethroid
546 resistance in the voltage-gated sodium channel of the tomato leaf miner (*Tuta*
547 *absoluta*). *Insect biochemistry and molecular biology*. 2012 Jul 1; 42(7):506-13.
- 548 35. Roditakis E, Vasakis E, Grispou M, Stavrakaki M, Nauen R, Gravouil M, Bassi A.
549 First report of *Tuta absoluta* resistance to diamide insecticides. *Journal of Pest*
550 *Science*. 2015 Mar 1; 88(1):9-16.
- 551 36. Linn CE, Campbell MG, Roelofs WL. Pheromone components and active spaces:
552 what do moths smell and where do they smell it? *Science*. 1987 Aug 7;
553 237(4815):650-2.
- 554 37. Prasad Y, Prabhakar M. Pest monitoring and forecasting. Integrated pest
555 management: principles and practice. Oxfordshire, UK: Cabi. 2012:41-57.
- 556 38. Megido RC, Haubruge E, Verheggen FJ. First evidence of deuterotokous
557 parthenogenesis in the tomato leafminer, *Tuta absoluta* (Meyrick)(Lepidoptera:
558 Gelechiidae). *Journal of Pest Science*. 2012 Dec 1; 85(4):409-12.

- 559 39. Witzgall P, Kirsch P, Cork A. Sex pheromones and their impact on pest
560 management. Journal of chemical ecology. 2010 Jan 1; 36(1):80-100.
- 561 40. Silva, Shênia Santos. "Fatores da biologia reprodutiva que influenciam o manejo
562 comportamental de *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae)." (2008).
- 563 41. Megido RC, Haubruge E, Verheggen FJ. First evidence of deuterotokous
564 parthenogenesis in the tomato leafminer, *Tuta absoluta* (Meyrick) (Lepidoptera:
565 Gelechiidae). Journal of Pest Science. 2012 Dec 1; 85(4):409-12.
- 566 42. Caparros Megido R, Haubruge E, Verheggen F. Pheromone-based management
567 strategies to control the tomato leafminer, *Tuta absoluta* (Lepidoptera: Gelechiidae).
568 A review. Biotechnologie, Agronomie, Société et Environnement= Biotechnology,
569 Agronomy, Society and Environment [= BASE]. 2013; 17(3):475-82.
- 570 43. Uchoa-Fernandes MA, Della Lucia TM, Vilela EF. Mating, oviposition and pupation
571 of *Scrobipalpuloides absoluta* (Meyr.)(Lepidoptera: Gelechiidae). Anais da
572 Sociedade Entomologica do Brasil. 1995; 24(1):159-64.
- 573 44. HASSAN N, AL-ZAIDI S. *Tuta absoluta*—Pheromone mediated management
574 strategy. International pest control. 2010; 52(3):158-60.
- 575 45. Real insight on *Tuta absoluta*, Real PM, Kenya (Ltd). [http://realipm.com/wp-](http://realipm.com/wp-content/uploads/2015/10/Real-insights-Tuta-Absoluta.pdf)
576 [content/uploads/2015/10/Real-insights-Tuta-Absoluta.pdf](http://realipm.com/wp-content/uploads/2015/10/Real-insights-Tuta-Absoluta.pdf). Accessed March` 6th,
577 2018
- 578 46. Retta AN, Berhe DH. Tomato leaf miner—*Tuta absoluta* (Meyrick), a devastating pest
579 of tomatoes in the highlands of Northern Ethiopia: A call for attention and action.
580 Research Journal of Agriculture and Environmental Management. 2015 Jun;
581 4(6):264-9.
- 582 47. Castañé C, Arnó J, Gabarra R, Alomar O. Plant damage to vegetable crops by
583 *zoophytophagous mirid* predators. Biological control. 2011 Oct 1; 59(1):22-9.

- 584 48. González-Cabrera J, Mollá O, Montón H, Urbaneja A. Efficacy of *Bacillus*
585 *thuringiensis* (Berliner) in controlling the tomato borer, *Tuta absoluta*
586 (Meyrick)(Lepidoptera: Gelechiidae). BioControl. 2011 Feb 1; 56(1):71-80.
- 587 49. Cabello T, Gallego JR, Fernandez FJ, Gamez M, Vila E, Del Pino M, Hernandez–
588 Suarez E. Biological control strategies for the South American tomato moth
589 (Lepidoptera: Gelechiidae) in greenhouse tomatoes. Journal of economic
590 entomology. 2012 Dec 1;105(6):2085-96.
- 591 50. Buitenhuis R, Shipp L, Scott-Dupree C. Dispersal of *Amblyseius swirskii* Athias-
592 Henriot (Acari: Phytoseiidae) on potted greenhouse chrysanthemum. Biological
593 Control. 2010 Feb 1; 52(2):110-4.
- 594 51. Momen F, Metwally A, Nasr A, Ebadah I, Saleh K. First report on suitability of the
595 tomato borer *Tuta absoluta* eggs (Lepidoptera: Gelechiidae) for eight predatory
596 phytoseiid mites (Acari: Phytoseiidae) under laboratory conditions. Acta
597 Phytopathologica et Entomologica Hungarica. 2013 Dec 1; 48(2):321-31.
- 598 52. Park HH, Shipp L, Buitenhuis R. Predation, development, and oviposition by the
599 predatory mite *Amblyseius swirskii* (Acari: Phytoseiidae) on tomato russet mite (Acari:
600 Eriophyidae). Journal of Economic Entomology. 2010 Jun;103(3):563-9.
- 601 53. SanchezJ.a. & Iacasa. Impact of the zoophytophagous plant bug
602 *Nesidiocoris tenuis*(Heteroptera: Miridae) on tomato yield. 2008. J. Econ. Entomol.
603 101: 1864–1870
- 604 54. Goula M, Alomar O. Miridos (Heteroptera Miridae) de interés en el control integrado
605 de plagas en el tomate. Guía para su identificación. Bol. San. Veg. Plagas.
606 1994;20(1):131-43.
- 607 55. Pires LM, Marques EJ, de Oliveira JV, Alves SB. Selection of isolates of
608 entomopathogenic fungi for controlling *Tuta absoluta* (Meyrick)(Lepidoptera:
609 Gelechiidae) and their compatibility with insecticides used in tomato crop.
610 Neotropical entomology. 2010 Dec;39(6):977-84.

- 611 56. Arnó J, Sorribas R, Prat M, Matas M, Pozo C, Rodríguez D, Garreta A, Gómez A,
612 Gabarra R. *Tuta absoluta*, a new pest in IPM tomatoes in the northeast of Spain.
613 IOBC/wprs Bull. 2009; 49:203-8.
- 614 57. Desneux N, Wajnberg E, Wyckhuys KA, Burgio G, Arpaia S, Narváez-Vasquez CA,
615 González-Cabrera J, Ruescas DC, Tabone E, Frandon J, Pizzol J. Biological
616 invasion of European tomato crops by *Tuta absoluta*: ecology, geographic
617 expansion and prospects for biological control. Journal of pest science. 2010 Aug 1;
618 83(3):197-215.
- 619 58. Huang Z., C. Guan and X.Guan. Cloning, characterization and expression of a new
620 cry1Ab gene from *Bacillus thuringiensis* WB9. Biotechnol Lett. 2004, 26:1557-1561
- 621 59. Joel González-Cabrera, Judit Arnó and Rosa Gabarra Prospects for the biological
622 control of *Tuta absoluta* in tomatoes of the Mediterranean basin Alberto Urbaneja^{1,*},
623 Article first published online: 25 JUN 2012 Society of Chemical Industry
- 624 60. Garcia-del-Pino F, Alabern X, Morton A,. Efficacy of soil treatments of
625 entomopathogenic nematodes against the larvae, pupae and adults of *Tuta absoluta*
626 and their interaction with the insecticides used against this insect. BioControl, 2013,
627 58(6):723-731. <http://rd.springer.com/journal/10526>
- 628 61. Pires, L.M., Marques, E.J., de Oliveira, J.V., Alves, S.B. Selection of isolates of
629 entomopathogenic fungi for controlling *Tuta absoluta* (Meyrick) (Lepidoptera:
630 Gelechiidae) and their compatibility with insecticides used in tomato crop,
631 Neotropical Entomology, (2010): 39: 977-984.
- 632 62. Giustolin, T.A., Vendramim, J.D., Alves, S.B., Vieira, S.A. Associated effect between
633 tomato resistant genotype and *Bacillus thuringiensis* var. kurstak on the development
634 of *Tuta absoluta* Meyrick (Lep., Gelechiidae), Neotropical Entomology, (2001), 30:
635 461-465

* E-mail address: nbszango@yahoo.com.

- 636 63. Rodríguez, M.S., Gerding, M.P., France, A. Entomopathogenic fungi isolates
637 selection for egg control oftomato moth *Tuta absoluta* Meyrick (Lepidoptera:
638 Gelechiidae), AgriculturaTecnica, (2006). 66: 151-158
- 639 64. Flavia A.C., Silva D.A., Martinez S. Effect of Neem seed oil aqueous solutions on
640 survival and development of the predator *Cyclonedasan guinea* (L.) (Coleoptera:
641 Coccinellidae). Neotropical Entomology, (2004) 33:19.
- 642 65. Edrees A.E.E. Effect of Neem seeds ethanolic extracts on mortality of tomato leaf-
643 miner *Tuta absoluta* (Lepidoptera: Gelechiidae).Thesis of Master, Sudan University
644 of Science and Technology, (2014), 1-71.
- 645 66. Gonçalves-Gervásio, R.D.R., Vendramim, D.J. Bioactivity of aqueous Neem seeds
646 extract on the *Tuta absoluta*(Meyrick, 1917) (Lepidoptera: Gelechiidae) in three
647 ways of application, Ciência e Agrotecnologia, (2007):31 (1): 28-34
- 648 67. Martinez S., van Emden H.F. Growth disruption, abnormalities and mortality of
649 *Spodoptera littoralis* (Boisduval) (Lepidoptera: Noctuidae) caused by Azadirachtin.
650 Neo tropical Entomology, (2001) 30:113-125.
- 651 68. Ajten Berxolli and Shpend Shahini. Azadirachtin, a useful alternative for controlling
652 *Tuta absoluta* (myerick) European Journal of Physical and Agricultural Sciences
653 (2017) ,Vol. 5 No. 2, 2017 ISSN 2056-5879
- 654 69. Hill MP, Hoffmann AA, Macfadyen S, Umina PA, Elith J. Understanding niche shifts:
655 Using current and historical data to model the invasive red legged earth mite,
656 *Halotydeus destructor* Divers Distrib. (2012) 18(2):191–203

657

658 **FIGURE CAPTIONS**

- 659 1. Distributions of *T. absoluta* in Tomato Growing Areas in Nigeria
- 660 2. Adult (A) and larvae (B) of tomato leafminer
- 661 3. Symptoms of *T. absoluta* Damages on Leaf, Fruit and Stem

- 662 4. Tomato Prices during 2015 *T. absoluta* Outbreak (Naira per Basket)
- 663 5. Tray Trap Technique (TTT) in Nigeria
- 664 6. NIHORT-Raktin, NIHORT-Lyptol and O-Ruptur (National Horticultural Research
- 665 Institute, Ibadan, Nigeria).
- 666