

## Original Research Article

# Growth and Yield Benefit of Cabbage as Influenced by Nutrients and Leaf Plucking

### ABSTRACT

The experiment was carried out at the Horticulture Farm of Sher-e-Bangla Agricultural University, Dhaka during the period from September 2016 to December 2016 to find out the growth, yield and economic benefit of cabbage as influenced by nutrients and leaf plucking. The research comprises of two factors: Factor A: NPK nutrients (four levels) as-  $N_0$  = control,  $N_1 = N_{100} P_{30} K_{100}$  kg ha<sup>-1</sup>,  $N_2 = N_{120} P_{40} K_{110}$  kg ha<sup>-1</sup>,  $N_3 = N_{140} P_{50} K_{120}$  kg ha<sup>-1</sup> and Factor B: leaf plucking (three levels) as-  $L_0$  = No leaf plucked,  $L_1$  = 4-leaves plucked and  $L_2$  = 6-leaves plucked. The research was set up in Randomized Complete Block Design with three replications. In case of nutrients, the highest gross yield (90.53 t ha<sup>-1</sup>) and marketable yield (68.95 t ha<sup>-1</sup>) were obtained from  $N_3$ , while the lowest gross yield (60.26 t ha<sup>-1</sup>) and marketable yield (44.24 t ha<sup>-1</sup>) from  $N_0$ . For dissimilar levels of leaf plucking, the highest gross yield (80.64 t ha<sup>-1</sup>) and marketable yield (62.08 t ha<sup>-1</sup>) were recorded from  $L_1$ , whereas the lowest gross yield (74.13 t ha<sup>-1</sup>) and marketable yield (56.96 t ha<sup>-1</sup>) from  $L_0$ . Due to combined effect, the highest gross yield (94.38 t ha<sup>-1</sup>) and marketable yield (71.91 t ha<sup>-1</sup>) were recorded from  $N_3L_1$ , whereas the lowest gross yield (58.75 t ha<sup>-1</sup>) and marketable yield (41.15 t ha<sup>-1</sup>) from  $N_0L_0$ . From the economic point of view, the highest benefit cost ratio (BCR) was 2.35 noted from  $N_3L_1$  and the lowest (1.63) from  $N_0L_0$ . It is evident that the  $N_3L_1$  gave the best performance for the growth, yield and economic benefit of cabbage. So,  $N_{140} P_{50} K_{120}$  kg ha<sup>-1</sup> with 4-leaves plucked may be considered as an optimum dose for cabbage production.

**Keywords:** Cabbage, NPK nutrients, leaf plucking, yield, Benefit Cost Ratio (BCR)

### 1. INTRODUCTION

Cabbage (*Brassica oleracea* var. *capitata* L.) is one of the most significant and nutritious winter leafy vegetables which belongs to the family Cruciferae. It is a biennial crop that is grown as an annual, unless it is grown for seed production [1] [2]. It can grow simply under wide range of environmental condition in both temperate and tropical, but cool moist climate is most suitable [3]. The origin of cabbage is the Western Europe and north shores of the Mediterranean Sea [4]. The edible portion of cabbage plant is head which is formed by the fleshy leaves overlapping one another. It has been reported that 100 g of green edible portion of cabbage contains 92% water, 24 kilocalories of food energy, 1.5 g of protein, 4.8 g of carbohydrate, 40 mg of calcium, 0.6 mg of iron, 600 IU of carotene, 0.05 mg of riboflavin, 0.3 mg of niacin and 60 mg of vitamin C [5]. It has been documented as a very necessary

vegetable to the farmers in providing income and nutrition worldwide [6]. Cultivation of cabbage is mainly done in winter season in Bangladesh. It is cultivated in 16.6 thousand hectares with a production of 220 thousand metric tons and the average yield is about 9 t ha<sup>-1</sup> [7]. Now, India is the second largest manufacturer of cabbage in the world, next to China, accounting for 16.55 per cent of the world area and 12.79 per cent of the world production [8]. Cabbage can play a critical role in elevating the nutritional status of Bangladesh, as it is rich in vitamins and minerals such as ascorbic acid, contains appreciable quantities of thiamin, riboflavin, calcium and iron [9]. Among the vegetables, it covers about 5% production under vegetable crops in Bangladesh [10]. However, low yield in Bangladesh may be attributed to a number of reasons viz. lack of quality seeds, nutrients unavailability, disease and insect infestation, improper or limited irrigation facilities etc.

The demand of cabbage as leafy vegetable is plentiful but the productivity of cabbage per unit area is quite low in Bangladesh due to excessive use of nutrients. Use of imbalanced nutrients in the soils may be harmful and causing our agricultural soil degraded and unproductive [11]. Nutrients may be applied through two sources viz., organic and inorganic sources. It enhances plant growth by providing amendments to the soil via various nutrients ultimately obtaining higher yield of cabbage. It is compulsory to ensure availability of crucial nutrient components for getting higher production and quality yield in any crop [12]. Nitrogen, phosphorus and potassium have profound effect on crop productivity and quality. Nitrogen is an essential plant nutrient, which is involved in physiological processes and enzyme activity [13] [14]. It plays a significant role in the building up of protoplasm and protein which induce cell division and initiate meristematic activities of plant when applied in optimum quantity. On the other hand, shortage of nitrogen during early growth may lead to the condition known as "buttoning" in Chinese cabbage when plant becomes stunted with reduce leaf development [15]. Phosphorus is concerned in energy transfer and nutrient movement within the plant. Adequate availability of phosphorus stimulates root development, increases stalk and stem strength and improves flower formation, fruiting and seed production. It also enhances uniform and early crop maturity, increases the nitrogen fixing ability of legumes, improves crop quality and increases resistance to plant diseases [16] [17] [18]. Potassium also increases better yield and improves quality during translocation of carbohydrate within crops. It exerts balancing role on the effects of both nitrogen and phosphorus, consequently it is especially important in multi-nutrient fertilizer application [19].

Leaf plucking in cabbage is an important factor for higher growth and yield of cabbage production. It means the removal of unfolded leaves or basal leaves without affecting the source-sink balance for proper head development. It is recommended that the successful cabbage production is possible by the application of basal 4-leaves plucking [20]. The plucked leaves may be positive either as vegetables or fodder as increased total biomass production. As cabbage heading begins, leaves become broader and sessile, and more erect in their posture [21]. After the formation and development of cabbage heads, the basal leaves are occasionally harbor of pathogen and insect which is normally induced decay before the time of harvest. Older or unfolded leaves are also competitive in nutrients requirement with younger leaves in cabbage head. Those impacts on slowly head development and maturation or sometimes cabbage head also indiscriminately infested with insect or fungal infection. The assemblage of layers of leaves over the growing point requires the maintenance of a short stem during the heading period [22]. But the possibility of leaf plucking of cabbage has not been explored earlier in Bangladesh. The present exploration was undertaken to evaluate the performance of nutrients and leaf plucking on growth, yield and economic return of cabbage.

## 2. MATERIAL AND METHODS

### 2.1 Experimental site

The experiment was conducted at the Horticulture Research Farm of Sher-e-Bangla Agricultural University (SAU), Dhaka, Bangladesh during the period from September 2016 to December 2016. Experimental site situated an elevation of 8 meters above the sea level in Agro-ecological zone of "Madhupur Tract" (AEZ-28) [23]. The soil was sandy loam and medium high land in texture having pH 5.46- 5.62.

### 2.2 Experiment Frame Work

The research was consisted of two factors: Factor A: NPK nutrients (four levels) as-  $N_0$  = control,  $N_1 = N_{120} P_{30} K_{100} \text{ kg ha}^{-1}$ ,  $N_2 = N_{140} P_{40} K_{120} \text{ kg ha}^{-1}$ ,  $N_3 = N_{160} P_{50} K_{140} \text{ kg ha}^{-1}$  and Factor B: leaf plucking (three levels) as-  $L_0$  = No leaf plucked,  $L_1$  = 4-leaves plucked and  $L_2$  = 6-leaves plucked. The two factors experiment was laid out following Randomized Complete Block Design (RCBD) with three replications. The experiment was divided into three equal blocks where each block was divided into 12 plots. Then 12 treatment combinations were allotted at randomly in each block. The size of the each unit plot was 1.8 m  $\times$  1.6 m. The distance maintained between two blocks and two plots were 0.5 m and 0.5 m, respectively. The seedlings were transplanted with maintaining distance row to row 60 cm and plant to plant 40 cm.

### 2.3 Application of manure and fertilizers

About 10 t  $\text{ha}^{-1}$  well decomposed cow dung was applied only control (as  $N_0$  treatment) plot and properly incorporated to the soil during final land preparation whereas others plot were applied with inorganic fertilizer as per treatment. Doses of inorganic fertilizers (Urea, TSP and MoP) were applied in the experimental plot according to the treatments (Table 1). Whole amount of TSP and half amount of MoP were also applied as basal dose before seedlings transplanting in the main field. 1st top dressing of urea was applied when seedlings established in the main field about 10days after seedling transplanting. 2nd top dressing of urea and rest amount of MoP was applied about 25days after 1st top dressing. Then rest amount of urea was applied as 3rd installment about 40 days after transplanting. Each top dressing was followed by manual irrigation.

**Table 1. Doses of nutrients application in the main field as per treatment**

Treatments	Available nutrients ( $\text{kg ha}^{-1}$ )			Fertilizers ( $\text{kg ha}^{-1}$ )			Doses ( $\text{g plot}^{-1}$ )		
	N	P	K	Urea	TSP	MoP	Urea	TSP	MoP
$N_1$	120	30	100	260.87	150	200	75.13	43.2	57.6
$N_2$	140	40	120	304.35	200	240	87.65	57.6	70.0
$N_3$	160	50	140	347.83	250	280	100.00	72.0	80.6

### 2.4 Application of Leaf Plucking

Leaf plucking was stunted only when the head grew 12-15 cm in diameter. Initially, two bottom leaves were plucked normally which was followed by further leaf plucking as per treatment. Older and unfolded leaves were plucked at 30 DAT and 40 DAT as per treatment of leaf plucking. The plucked leaves may be useful either as vegetables or fodder as increased total biomass production.

### 3. RESULTS AND DISCUSSION

#### 3.1 Plant height (cm)

Considerable variation was found among the different levels of nutrients in respect of plant height of cabbage (Table 2). At 40 DAT and 50 DAT, the tallest plant (34.42cm and 39.86cm, respectively) was recorded from N<sub>3</sub> while the shortest plant (25.76cm and 30.48cm, respectively) from N<sub>0</sub>. At harvest, the tallest plant (45.68cm) was observed in N<sub>3</sub> while the shortest plant (34.95cm) was in N<sub>0</sub>. Tekasangla *et al.* [24], Mankar *et al.* [25], Kumar *et al.* [26] and Farooque and Mondal [27] also noticed the related findings of the present study. At 50 DAT, the tallest plant (37.79cm) was observed in L<sub>2</sub> while the shortest plant (33.20cm) was in L<sub>0</sub> (Table 3). At harvest, the tallest plant (43.50cm) was in L<sub>2</sub> while the shortest plant (37.06cm) was in L<sub>0</sub> (Table 3).

**Table 2.** Effect of nutrients on growth parameters at different growth stages of cabbage

Treatments	Plant height (cm)			Number of loose leaves			Days required for head maturity
	40 DAT	50 DAT	At harvest	40 DAT	50 DAT	At harvest	
N <sub>0</sub>	25.76	30.48	34.95	9.20	11.20	12.68	60.32
N <sub>1</sub>	29.27	34.77	41.03	10.26	12.18	13.51	62.45
N <sub>2</sub>	32.45	37.95	42.42	11.21	12.94	14.43	63.75
N <sub>3</sub>	34.42	39.86	45.68	11.87	14.00	15.74	65.29
CV %	5.41	7.67	7.25	11.60	10.58	7.45	4.88
LSD (0.05)	1.91	1.87	2.82	0.64	0.83	0.69	1.47

The interaction between nutrients and leaf plucking treatments affects significantly on plant height (Table 4). At 40 DAT, the tallest plant (34.67cm) was obtained from N<sub>3</sub>L<sub>2</sub> which was statistically identical to N<sub>3</sub>L<sub>1</sub> while the shortest plant (25.53cm) was in N<sub>0</sub>L<sub>0</sub>. At 50 DAT, the tallest plant (41.70cm) was observed in N<sub>3</sub>L<sub>2</sub> while the shortest plant (27.90cm) was in N<sub>0</sub>L<sub>0</sub>. At harvest, the tallest plant (48.47cm) was obtained from N<sub>3</sub>L<sub>2</sub> whereas the shortest plant (31.60 cm) was in N<sub>0</sub>L<sub>0</sub>.

**Table 3.** Effect of leaf plucking on growth parameters at different growth stages of cabbage

Treatments	Plant height (cm)			Number of loose leaves			Days required for head maturity
	40 DAT	50 DAT	At harvest	40 DAT	50 DAT	At harvest	
L <sub>0</sub>	30.38	33.20	37.06	10.05	11.93	13.48	61.92
L <sub>1</sub>	30.65	36.30	42.50	11.16	13.26	14.93	63.50
L <sub>2</sub>	30.40	37.79	43.50	10.68	12.55	13.85	63.43
CV %	5.41	7.67	7.25	11.68	10.58	7.45	4.88
LSD (0.05)	NS	2.11	2.44	0.55	0.59	0.60	1.22

#### 3.2 Number of loose leaves plant<sup>-1</sup>

At 40, 50 DAT, the maximum number of loose leaves plant<sup>-1</sup> (11.87 and 14.00, respectively) was recorded from N<sub>3</sub> while the lowest (9.20 and 11.20 respectively) was in N<sub>0</sub> (Table 2). At harvest, the maximum (15.74) was recorded from N<sub>3</sub> while the minimum (12.68) was in N<sub>0</sub>. At 40 DAT and 50 DAT, the maximum (11.16 and 13.26, respectively) was observed in L<sub>1</sub> while the minimum (10.05 and 11.93) was in L<sub>0</sub> (Table 3). At harvest, the maximum (14.93) was recorded from L<sub>1</sub> while the minimum (13.48) was in L<sub>0</sub>.

**Table 4. Combined effect of nutrients and leaf plucking on growth parameters at different stages of cabbage**

Treatments	Plant height (cm)			Number of loose leaves			Days required for head maturity
	40 DAT	50 DAT	At harvest	40 DAT	50 DAT	At harvest	
N <sub>0</sub> L <sub>0</sub>	25.53	27.90	31.60	8.76	10.43	12.07	59.25
N <sub>0</sub> L <sub>1</sub>	25.96	30.73	35.86	9.76	11.86	13.30	60.91
N <sub>0</sub> L <sub>2</sub>	25.80	32.80	37.40	9.06	11.30	12.67	60.80
N <sub>1</sub> L <sub>0</sub>	28.80	31.97	36.73	9.86	11.80	13.03	61.96
N <sub>1</sub> L <sub>1</sub>	29.66	35.50	42.73	10.70	12.73	14.10	62.93
N <sub>1</sub> L <sub>2</sub>	29.33	36.86	43.63	10.20	12.03	13.40	62.47
N <sub>2</sub> L <sub>0</sub>	32.26	35.60	39.16	10.40	12.23	13.93	62.91
N <sub>2</sub> L <sub>1</sub>	32.56	38.46	43.56	11.83	13.53	15.30	64.25
N <sub>2</sub> L <sub>2</sub>	32.53	39.80	44.53	11.40	13.06	14.07	64.08
N <sub>3</sub> L <sub>0</sub>	34.20	37.36	40.76	11.16	13.26	14.90	63.57
N <sub>3</sub> L <sub>1</sub>	34.40	40.53	47.83	12.36	14.93	17.03	65.90
N <sub>3</sub> L <sub>2</sub>	34.67	41.70	48.47	12.06	13.80	15.30	66.40
CV %	5.41	7.67	7.25	11.68	10.58	7.45	4.88
LSD (0.05)	3.44	2.23	3.88	1.11	1.11	1.20	1.11

Different levels of nutrients and leaf plucking showed significant differences due to their interaction effect on number of loose leaves per plant of cabbage at 40, 50 DAT and at harvest. At 40 DAT, the maximum (12.36) was recorded from N<sub>3</sub>L<sub>1</sub> while the minimum (8.76) was in N<sub>0</sub>L<sub>0</sub> (Table 4). At 50 DAT, the maximum (14.93) was counted in N<sub>3</sub>L<sub>1</sub> whereas the minimum (10.43) was in N<sub>0</sub>L<sub>0</sub>. At harvest, the maximum (17.03) was obtained from N<sub>3</sub>L<sub>1</sub> while the minimum (12.07) was in N<sub>0</sub>L<sub>0</sub>.



At 40 DAT



At 50 DAT

**Fig. 1. Plots after leaf plucking**

### 3.3 Days required to head maturity

The maximum days required to head maturity (65.29) was obtained from N<sub>3</sub> while the minimum (60.32) was in N<sub>0</sub> at harvest (Table 2). Different levels of leaf plucking showed significant variation on days required to head maturity. The maximum days required to head maturity (63.50) was observed in L<sub>1</sub> while the minimum (61.92cm) was in L<sub>0</sub> at harvest (Table 3). Combined effect of nutrients and leaf plucking gives the maximum (66.40) days required to head maturity was recorded from N<sub>3</sub>L<sub>2</sub> which was statistically similar to N<sub>3</sub>L<sub>1</sub>, while N<sub>0</sub>L<sub>0</sub> showed the minimum (59.25) days required to head maturity (Table 4).

### 3.4 Diameter of stem (cm)

The maximum diameter of stem of cabbage (3.57cm) was observed in N<sub>3</sub> while the minimum (2.56cm) was in N<sub>0</sub> (Table 8). Different levels of leaf plucking showed significant influence on diameter of stem of cabbage. The maximum (3.47cm) was observed in L<sub>1</sub> whereas the minimum (2.93cm) was in L<sub>0</sub> (Table 9). Combined effect of nutrients and leaf plucking showed significant effect on diameter of stem of cabbage. The maximum diameter of stem (3.86cm) was recorded from N<sub>3</sub>L<sub>1</sub> while N<sub>0</sub>L<sub>0</sub> gave the minimum (2.36cm) diameter of stem (Table 10).

**Table 8.** Effect of nutrients on growth and yield contributing parameters at harvest stage

Treatments	Diameter of stem (cm)	Head Characteristics				
		Diameter of head (cm)	Head height (cm)	Head weight (kg)	Head thickness (cm)	%Dry matter
N <sub>0</sub>	2.56	16.60	10.65	1.20	7.30	6.33
N <sub>1</sub>	3.17	18.24	12.26	1.46	9.34	6.76
N <sub>2</sub>	3.42	19.47	13.40	1.69	10.35	7.33
N <sub>3</sub>	3.57	20.38	14.17	1.92	10.75	8.02
CV %	8.13	9.56	5.39	6.31	8.87	10.93
LSD (0.05)	0.19	0.19	0.32	0.151	0.39	0.37

### 3.5 Diameter of head (cm)

The highest diameter of head (20.38cm) was recorded from N<sub>3</sub> while the lowest (16.60cm) was in N<sub>0</sub> (Table 8). Similar findings on diameter of head are reported by Hossain *et al.* [11], Mankar *et al.* [25] and Naher *et al.* [28]. The highest (19.05cm) was observed in L<sub>1</sub> whereas the lowest (18.13cm) was recorded from L<sub>0</sub> (Table 9). Combined effect of different levels of nutrients and leaf plucking showed significant effect on diameter of head of cabbage (Table 10). The highest diameter of head (20.93cm) was observed in N<sub>3</sub>L<sub>1</sub> while the lowest (15.98cm) was in N<sub>0</sub>L<sub>0</sub>.

**Table 9.** Effect of leaf plucking on growth and yield contributing parameters at harvest stage

Treatments	Diameter of stem (cm)	Head Characteristics				
		Diameter of head (cm)	Head height (cm)	Head weight (kg)	Head thickness (cm)	%Dry matter
L <sub>0</sub>	2.93	18.13	11.96	1.44	8.60	6.86
L <sub>1</sub>	3.47	19.05	13.05	1.68	9.89	7.28

L <sub>2</sub>	3.13	18.85	12.85	1.58	9.80	7.19
CV %	8.13	9.56	5.39	6.31	8.87	10.93
LSD (0.05)	0.16	0.17	0.28	0.081	0.34	0.32

### 3.6 Head height (cm)

The maximum head height of cabbage (14.17cm) was obtained from N<sub>3</sub> while the minimum (10.65cm) was recorded in N<sub>0</sub> (Table 8). Hossain *et al.* [11] and Singh *et al.* [29] reported that the similar views on head height of cabbage of the present experiment. The maximum (13.05cm) was observed in L<sub>1</sub> while the minimum (11.96cm) was found in L<sub>0</sub> (Table 9). Combined effect of nutrients and leaf plucking showed significant effect on head height of cabbage (Table 10). The maximum (14.73cm) was recorded from N<sub>3</sub>L<sub>1</sub> which was statistically similar to N<sub>3</sub>L<sub>2</sub> (14.40cm), while N<sub>0</sub>L<sub>0</sub> gave the minimum (10.16cm).

**Table 10. Combined effect of nutrients and leaf plucking on growth and yield contributing parameters at harvest of cabbage**

Treatments	Diameter of stem (cm)	Head Characteristics				
		Diameter of head (cm)	Head height (cm)	Head weight (kg)	Head thickness (cm)	%Dry matter
N <sub>0</sub> L <sub>0</sub>	2.36	15.98	10.16	1.09	6.80	6.13
N <sub>0</sub> L <sub>1</sub>	2.80	16.91	10.96	1.26	7.63	6.53
N <sub>0</sub> L <sub>2</sub>	2.50	16.92	10.83	1.24	7.46	6.33
N <sub>1</sub> L <sub>0</sub>	2.96	17.63	11.73	1.32	8.43	6.43
N <sub>1</sub> L <sub>1</sub>	3.40	18.59	12.86	1.56	9.83b	6.97
N <sub>1</sub> L <sub>2</sub>	3.13	18.52	12.20	1.49	9.76b	6.90
N <sub>2</sub> L <sub>0</sub>	3.06	18.94	12.56	1.56	9.23	7.00
N <sub>2</sub> L <sub>1</sub>	3.80	19.77	13.66	1.80	10.96	7.53
N <sub>2</sub> L <sub>2</sub>	3.40	19.72	13.96	1.71	10.86	7.47
N <sub>3</sub> L <sub>0</sub>	3.33	19.96b	13.40	1.09	9.96	7.90
N <sub>3</sub> L <sub>1</sub>	3.86	20.93	14.73	2.08	11.15	8.10
N <sub>3</sub> L <sub>2</sub>	3.50	20.25	14.40	1.89	11.13	8.06
CV %	8.13	9.56	5.39	6.31	8.87	10.93
LSD (0.05)	0.33	0.34	0.56	0.138	0.69	0.69

### 3.7 Head weight (kg)

The highest head weight of cabbage (1.92 kg) was recorded from N<sub>3</sub> whereas the lowest (1.20 kg) was in N<sub>0</sub> (Table 8). Similar findings of head weight were observed with Mankar *et al.* [25]. The results under the present experiment were also fairly supported by Hasan and Solaiman [30]. The highest (1.68 kg) was observed in L<sub>1</sub> while the lowest (1.44 kg) was in L<sub>0</sub> (Table 9). The findings obtained from the experiment were partially conformed to Begum [20]. Due to combined effect of different levels of nutrients and leaf plucking, N<sub>3</sub>L<sub>1</sub> produced the highest head weight (2.08 kg) which was statistically similar to N<sub>3</sub>L<sub>1</sub> while the lowest (1.09 kg) was in N<sub>0</sub>L<sub>0</sub> (Table 10).

### 3.8 Head thickness (cm)

The highest head thickness (10.75cm) was found from N<sub>3</sub> while the lowest (7.30cm) Naher *et al.* [28] suggested that the similar results on head thickness of the present study (Table 8).

The highest head thickness (9.89cm) was attained from  $L_1$  whereas the lowest (8.60cm) was in  $L_0$  (Table 9). Interaction effect of different levels of nutrients and leaf plucking showed significant differences on thickness of head of cabbage. The highest head thickness (11.15cm) was recorded from  $N_3L_1$  which was statistically identical to  $N_3L_2$  (11.13cm) while the lowest (6.80 cm) was found from  $N_0L_0$  (Table 10).

### 3.9 Dry matter content of head (%)

The maximum dry matter content (8.02%) was observed from  $N_3$  while the minimum (6.33 %) was in  $N_0$  (Table 8). The maximum dry matter content (7.28 %) was observed in  $L_1$  which was statistically similar to  $L_2$  (7.19 %) whereas the minimum (6.86 %) was found from  $L_0$  (Table 9). Combined effect of nutrients and leaf plucking showed significant effect on % dry matter content of head. The maximum dry matter content (8.10 %) was recorded from  $N_3L_1$  which was statistically identical to  $N_3L_2$  (8.10 %) while the minimum (6.13 %) was in  $N_0L_0$  (Table 10).

### 3.10 Weight of whole plant (kg)

Different levels of nutrients showed significant effect on weight of whole plant of cabbage under the present study (Table 11). The highest (2.46 kg) was obtained from  $N_3$  while the lowest (1.39 kg) was in  $N_0$ . This result of the present study was partially supported by Mankar *et al.* [25] and Bojokalfa *et al.* [31]. The highest (2.09 kg) was recorded from  $L_1$  whereas the lowest (1.88 kg) was in  $L_0$  (Table 12). Combined effect of different levels of nutrients and leaf plucking showed significant variation on weight of whole plant of cabbage (Table 13). The highest (2.60 kg) was observed from  $N_3L_1$  which was statistically similar to  $N_3L_2$  while the lowest (1.30 kg) was in  $N_0L_0$ .

**Table 11. Effect of nutrients on yield parameters at harvest stage of cabbage**

Treatments	Weight of whole plant (kg plant <sup>-1</sup> )	Gross yield (t ha <sup>-1</sup> )	Marketable yield (t ha <sup>-1</sup> )	Economic production (kg plant <sup>-1</sup> )
$N_0$	1.39	60.26	44.24	1.34
$N_1$	1.91	75.66	60.79	1.52
$N_2$	2.20	83.07	64.35	1.74
$N_3$	2.46	90.53	68.95	1.90
CV %	9.34	9.63	10.27	11.43
LSD (0.05)	0.141	3.56	3.79	0.116

### 3.11 Gross yield (t ha<sup>-1</sup>)

The highest gross yield (90.53 t ha<sup>-1</sup>) was obtained from  $N_3$  while the lowest (60.26 t ha<sup>-1</sup>) was in  $N_0$  (Table 11). Jothi *et al.* [32] and Rahman [33] stated same views of the present study. It is evident that the highest gross yield (80.64 t ha<sup>-1</sup>) was observed in  $L_1$  while the lowest (74.13 t ha<sup>-1</sup>) was in  $L_0$  (Table 12). Begum [20] observed that 4-leaves plucking of cabbage was performed the highest gross yield which is similar to this result of the present study. Combined effect of nutrients and leaf plucking had significant effect on gross yield of cabbage. The highest gross yield (94.38 t ha<sup>-1</sup>) was observed in  $N_3L_1$  (94.38 t ha<sup>-1</sup>) while the lowest (58.75 t ha<sup>-1</sup>) was in  $N_0L_0$  (Table 13).



284 **Table 12. Effect of leaf plucking on yield parameters at harvest stage of cabbage**  
 285

Treatments	Weight of whole plant (kg plant <sup>-1</sup> )	Gross yield (t ha <sup>-1</sup> )	Marketable yield (t ha <sup>-1</sup> )	Economic production (kg plant <sup>-1</sup> )
L <sub>0</sub>	1.88c	74.13 c	56.96c	1.53c
L <sub>1</sub>	2.09a	80.64 a	62.08a	1.72a
L <sub>2</sub>	2.01b	77.36 b	59.71b	1.62b
CV %	9.34	9.63	10.27	11.43
LSD (0.05)	0.072	3.08	2.18	0.082

286

### 287 **3.12 Marketable yield (t ha<sup>-1</sup>)**

288

289 The highest marketable yield (68.95 t ha<sup>-1</sup>) was observed in N<sub>3</sub> while the lowest (44.24 t ha<sup>-1</sup>)  
 290 was in N<sub>0</sub> (Table 11). The results of the present study were partially supported by Singh [29].  
 291 The highest (62.08 t ha<sup>-1</sup>) was obtained from L<sub>1</sub> while the lowest (56.96 t ha<sup>-1</sup>) was in L<sub>0</sub>  
 292 (Table 12). The highest (71.91 t ha<sup>-1</sup>) was observed in N<sub>3</sub>L<sub>1</sub> while the lowest (41.15 t ha<sup>-1</sup>)  
 293 was in N<sub>0</sub>L<sub>0</sub> (Table 13).

294

295 **Table 13. Combined effect of nutrients and leaf plucking on yield parameters at**  
 296 **harvest stage**  
 297

Treatments	Weight of whole plant (kg plant <sup>-1</sup> )	Gross yield (t ha <sup>-1</sup> )	Marketable yield (t ha <sup>-1</sup> )	Economic production (kg plant <sup>-1</sup> )
N <sub>0</sub> L <sub>0</sub>	1.30	58.75	41.15	1.25
N <sub>0</sub> L <sub>1</sub>	1.43	62.43	46.98	1.42
N <sub>0</sub> L <sub>2</sub>	1.44	60.07	44.58	1.35
N <sub>1</sub> L <sub>0</sub>	1.76	71.32	58.78	1.42
N <sub>1</sub> L <sub>1</sub>	2.02	79.44	63.06	1.63
N <sub>1</sub> L <sub>2</sub>	1.96	76.18	60.49	1.50
N <sub>2</sub> L <sub>0</sub>	2.08	79.27	62.33	1.66
N <sub>2</sub> L <sub>1</sub>	2.31	86.77	66.32	1.84
N <sub>2</sub> L <sub>2</sub>	2.21	83.16	64.76	1.73
N <sub>3</sub> L <sub>0</sub>	2.36	87.15	65.90	1.80
N <sub>3</sub> L <sub>1</sub>	2.60	94.38	71.91	2.00
N <sub>3</sub> L <sub>2</sub>	2.42	90.03	68.99	1.90
CV %	9.34	9.63	10.27	11.43
LSD (0.05)	0.049	6.16	6.57	0.038

298

### 299 **3.13 Economic production (kg plant<sup>-1</sup>)**

300

301 The highest economic production (1.90 kg plant<sup>-1</sup>) was recorded from N<sub>3</sub> whereas the lowest  
 302 (1.34 kg plant<sup>-1</sup>) was in N<sub>0</sub> (Table 11). The findings of the present study are partially  
 303 supported with Sharma [34]. The highest economic production (1.72 kg plant<sup>-1</sup>) was  
 304 observed in L<sub>1</sub> whereas the lowest (1.53 kg plant<sup>-1</sup>) was in L<sub>0</sub> (Table 12). Combined effect of  
 305 different levels of nutrients & leaf plucking had significant effect on economic production of  
 306 cabbage. The highest economic production (2.00 kg plant<sup>-1</sup>) was observed in N<sub>3</sub>L<sub>1</sub> which was  
 307 statistically similar to N<sub>3</sub>L<sub>2</sub>, while the lowest (1.25 kg plant<sup>-1</sup>) was in N<sub>0</sub>L<sub>0</sub> (Table 13).

308  
309




Fig. 2. Some pictorial view of cabbage head as per treatment at harvest

#### 4. CONCLUSION

Crop yield and economic benefit of crop are both important for a crop production. Leaf plucking represents higher yield in cabbage plant than without no leaf plucking. According to the results of the present experiment, it may be concluded that efficient production of cabbage is increased by the application of nutrients and leaf plucking. Thus, the combined application of nutrients and leaf plucking may be helpful for higher & better qualitative cabbage production in considering crop productivity and economic return of cabbage. On the basis of benefit cost ratio, it may be suggested that  $N_{160} P_{50} K_{140} \text{ kg ha}^{-1}$  nutrients with 4-leaves plucked gave maximum and profitable yield of cabbage head.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Ryder EJ. Leafy salad vegetables. AVI publishing company. Inc., Westport, Connecticut; 1979.
2. Pierce LC. Vegetables; Characteristics, production and marketing. John Wiley and sons. Toronto, Canada; 1987.

- 335 3. Kibar B, Karaağaço, Hayati K. Correlation and Path Coefficient Analysis of Yield and  
336 Yield Components in Cabbage (*Brassica oleracea* Var. *capitata* L.) Acta Sci. Pol.  
337 2014:13 (6): 87-97.
- 338 4. Chauhan OVS. Vegetable production in India. Ram Prasad and sons, India; 1986.
- 339 5. Rashid MM. বঙ্গদেশের সবজি উৎপাদন. Bangla Academy, Dhaka, Bangladesh; 1993.  
340 Bangla.
- 341 6. FAOSTAT. Food and Agriculture Organization, United Nations. 2007.  
342 <http://faostat.fao.org>.
- 343 7. BBS. Bangladesh Bureau of Statistics of plan 10. Yearbook of Agricultural Statistics of  
344 Bangladesh. Bangladesh Bureau of Statistics of Plan., Govt. of the People's Republic of  
345 Bangladesh, Dhaka; 2010.
- 346 8. NHB. Final Area and Production Estimates for Horticulture Crops for 2013-2014.  
347 National Horticulture Board, Gurgaon, India; 2015.
- 348 9. Thompson HC, Kelly WC. Vegetable crops. 5<sup>th</sup> edition, New York, Tornado, London:  
349 McGraw Hill Book Co.; 1985.
- 350 10. BBS. Monthly Statistical Bulletin of Bangladesh Bureau of Statistics, Ministry of  
351 Planning, Govt. of the people's republic of Bangladesh, Dhaka; 2015.
- 352 11. Hossain D, Abuyusuf MAHM, Riad MM, Hussain AI. Response of cabbage to different  
353 levels of fertilizer application in salna silty clay loam soil. Bangladesh Res. Publication J.  
354 2011:6(2): 155-166.
- 355 12. Alam MS, Iqbal TMT, Amin M, Gaffer MA. কৃষিতাত্ত্বিক ফসলের উৎপাদন উন্নয়ন .  
356 Sirajgonj, Bangladesh; 1989. Bangla
- 357 13. Kodithuwakku DP, Kirthisinghe JP. The effect of different rates of nitrogen fertilizer on  
358 the growth, yield and postharvest life of cauliflower. Tropical Agric. Res. and Exten.  
359 2009:21 (1):110-114.
- 360 14. Neethu MT, Tripathi SM, Narwad AV, Sreeganesh S. Effect of N and P levels on growth  
361 and yield parameters of broccoli (*Brassica oleracea* var. *italica*) under South Gujarat soil  
362 conditions. Int. J. Trop. Agric. 2015:33 (2): 913-917.
- 363 15. Tindall HD. Vegetable in the Tropics. Macmillan Education Ltd. Houndmills. Basingstoke  
364 Hampshire. RhRI 2 \* S and London; 1983.[Cited from Hort. Abstr., 50 (6): 3791-1984]
- 365 16. Mitchell RL. Crop growth and culture. Ames, USA: Iowa State University Press; 1970.
- 366 17. Plaster EJ. Soil science and management. New York, USA: Delmar Publishers Inc.;  
367 1985
- 368 18. Plaster EJ. Soil science and management. 4<sup>th</sup> ed. New York, USA: Thomson Delmar  
369 Learning; 2003
- 370 19. Bardy NC. The Nature and properties of Soils. 10<sup>th</sup> edition, New York, USA: McMillan  
371 Publishing Company; 1990.
- 372 20. Begum HA. Effect of manuring and leaf plucking on growth and yield of cabbage. MS  
373 thesis, Department of Horticulture, Bangladesh Agricultural University, Mymensingh,  
374 Bangladesh; 2005.
- 375 21. Kato T, Sooen A. Physiological studies of head formation on cabbage. J. Jap.Soc. Hort.  
376 Sci. 1978:48 (4): 426-434.
- 377 22. North C. Studies in morphogenesis of *Brassica oleracea* L. Growth and development of  
378 cabbage during the vegetative phase. J. Exp. Bot., 1957:8 : 304-312.
- 379 23. UNDP. Land resource appraisal of bangladesh for agricultural development report 2:  
380 Agro-ecological Regions of Bangladesh, FAO, Rome, Italy; 1988.
- 381 24. Tekasangla SP, Kanaujia, Singh PK. Integrated nutrient management for quality  
382 production of cauliflower in acid alfisol of Nagaland. Karnataka J. Agric. Sci. 2015:28 (2):  
383 244-247.
- 384 25. Mankar A, Kumari C, Karuna K. Effect of nitrogen levels and microbial inoculants on  
385 growth, yield and quality of cabbage. Prog. Hort. 2015: 47(2): 296-299.

- 386 26. Kumar S, Singh JP, Rajbeer RN, Mohan B, Kaushik H, Kumar D. Influence of integrated  
387 nutrient management on growth and yield of cauliflower (*Brassica oleracea* var. *botrytis*  
388 L.) Int. J. Agric. Sci. 2013;9 (2):747-749.
- 389 27. Farooque, A. M. and Mondal, F. Effect of spacing and levels of nitrogen on growth and  
390 yield of cabbage. Bangladesh Hort. 1987;15 (2):1-6.
- 391 28. Naher MNA, Alam MN, Jahan N. Effect of nutrient management on the growth and yield  
392 of cabbage (*Brassica oleracea* var. *capitata* L.) in calcareous soils of bangladesh. The  
393 Agriculturists. 2014;12 (2): 24-33.
- 394 29. Singh KM, Chand T, Kumar M, Singh KV, Lodhi SK, Singh VP, Sirohi SV. Response of  
395 different doses of NPK and boron on growth and yield of broccoli (*Brassica oleracea* L.  
396 var. *italica*). IJBSM. 2015;6 (1): 108-112.
- 397 30. Hasan MR, Solaiman AHM. Efficacy of organic and organic fertilizer on the growth of  
398 *Brassica oleracea* L. (Cabbage). Intl. J. Agri. Crop Sci. 2012;4 (3): 128-138.
- 399 31. Bojokalfa MK, Kavak S, Ugur A, Yagmur B. Effect of phosphorus fertilizer application on  
400 yield and quality characteristics in savoy cabbage (*Brassica Oleracea* var. *sabudab* L.)  
401 arkeyEge/Universiteesi. 2003;40 (I): 17-24.
- 402 32. Jothi LJ, Papiiah CM, Ryagopalati R. Influence of NPK and *Azospirillum* on the yield of  
403 cabbage. South Indian Hort. 1993; 41 (5): 270-272.
- 404 33. Rahman MM. Effects of cowdung and NPK fertilizers on growth and yield of cabbage.  
405 MS thesis, Department of Horticulture, Bangladesh Agricultural University, Mymensingh-  
406 2202; 2005.
- 407 34. Sharma V. Effect of nutrient management on growth and yield of cauliflower (*Brassica*  
408 *oleracea* var. *botrytis*) inside low cost polyhouse. Himachal J. Agric. Res. 2016; 42 (1):  
409 88-92.