

CHALLENGES AFFECTING THE ADOPTION OF FARM FORESTRY LAND USE AROUND CHEPALUNGU FOREST

ABSTRACT

Anthropogenic activities around Chepalungu forest has lead to its reduction in size and has resulted in its degradation. Continued dependency on this forest may result in its total depletion despite its high biological diversity and natural resource conservation value. There have been low levels of adoption of farm forestry practice (less than 33% of farmers) around Chepalungu. This study identified challenges affecting the adoption of farm forestry practices around Chepalungu forest. The researchers study adopted ex-post facto research design in this study. The study was done in 2016. Study sites were four locations (Bing'wa, Siongiroi, Ndanai, and Abosi) selected around the forest using simple random sampling method. A total of 377 household questionnaires were administered in the four locations proportionately. Chi-Square and Mann Whitney U tests were used in the analysis of the results. Significant levels were expressed at $P < 0.05$ using SPSS version 17 software. The results showed that there was a significant association ($\chi^2 = 530.8$; $P < 0.01$) between types of farm forestry practices and challenges affecting farm forestry adoption. Notable challenges were: damage by animals, damage by man, tree nursery problems, inadequate capital, natural calamities, competing land uses, managerial problems and seed acquisition problems.

Key words: farm forestry, challenges, Chepalungu, adoption

1. INTRODUCTION

Over 60% of forest dependent people in the world are farmers [1]. A significant number of these farmers depend heavily on forest resources for their livelihood [2]. Due to this high dependence on existing forest resources, natural forests are being depleted and the supply of future forest products is becoming uncertain [3].

Natural forests in Kenya stand at about 1.2 million hectares, mostly being in high potential areas where they are facing intense competition with other land use practices [4, 5]. The growing population is exerting immense pressure on the forest resources with about 80% of forest dependent people in these areas being farmers [6, 7]. Forest ecosystems are fast dwindling in the high potential ecological zones, forcing people to move into forests and other areas which are less endowed in biodiversity [8].

The role of forests in the livelihood of the forest adjacent communities is diverse. Rural forest adjacent communities derive food, medicine and fuel-wood in addition they enjoy non-wood benefits such as spiritual, aesthetic and environmental services provided by forests. Therefore, continued degradation of forests is likely to reduce forest resources capacity to support environmental conservation and people's socio-economic livelihood [5, 9, 10].

Chepalungu forest is very important to people living near and far from it by providing both timber and non-timber forest products, but it is highly degraded due to grazing, settlement and farming. In

37 addition, forest excision at Chepalungu cleared important tree species such as *Olea capensis* (Olea)
38 and *Juniperus procera* (Cedar) [11]. Farm forestry, which is the growing of trees on farms for
39 subsistence and commercial purposes, is an important land use option that has the potential to take
40 over a substantial part of the functions of indigenous and plantation forests and to reduce pressure on
41 natural forests [12].

42 Farm forestry offers many benefits to both landholders and to the wider community. In the short term,
43 it can provide many private benefits, such as increasing the visual amenity of the farm, improved soil
44 stability and in some cases improved productivity of other farm activities [12]. Over the longer term,
45 farm forestry provides landholders with a source of income and a means of diversifying the farm
46 business [5, 13]. Farm forestry also offers many potential benefits to the wider community by
47 improving catchment water quality, reducing stream-bank erosion, and increasing landscape
48 biodiversity. A well managed farm forestry can also improve the economic and social sustainability of
49 a region [14].

50 In early 2000s, Green Belt Movement initiated conservation programmes that included farm forestry to
51 reduce continued encroachment and destruction of the remnant forest. However, [15] reported a low
52 level of adoption of farm forestry practice (less than 33% of farmers) around Chepalungu Forest but
53 they did not provide detailed account of the challenges affecting farm forestry development in the
54 area. Moreover, communities adjacent to Chepalungu forest are still dependent on this forest as
55 source of products accessed through destructive activities [15]. This study aimed at identifying
56 challenges affecting the adoption of farm forestry land use around Chepalungu forest.


57 **2. MATERIALS AND METHODS**

58 **2.1 Research Design**

59 The study employed the use of ex-post facto research design in establishing the farm forestry
60 challenges affecting farm forestry development among the communities adjacent to Chepalungu
61 forest.

62 **2.2 Study area**

63 Chepalungu Forest lies on latitude 00° 53' 00" S and longitude 35° 10' 00" E. The study was carried
64 out around Kapchumbe and Siongiroi blocks of Chepalungu forest, Bomet County, Kenya.
65 Chepalungu forest is administratively divided into two management blocks, Kapchumbe (in the South-
66 West) and Siongiroi (in the North-East) (Figure 1).



67
68 **Figure 1: Chepalungu forest and adjacent sampled settlement Locations [8]**

69 The area has medium to long cropping season followed by a medium to short and intermediate rains.
70 The mean annual rainfall is 1200 mm – 1350 mm per year with an altitude range of 1550m – 2000m
71 above the sea level. The mean annual temperature ranges from 17.9°C – 20.5°C.

72 The soils are predominantly loamy black cotton soils. Maize and marginal coffee crops are the main
73 crops in the area which occupies almost 18.72 Km² of the agricultural land [16].

74 **2.3 Target Population**

75 Kapchumbe and Siongiroi blocks of Chepalungu forest are adjoined by six settlement Locations.
 76 These locations are Abosi, Bingwa, Siongiroi, Makimenyi, Ndanai and Kongasis. These locations
 77 have a total of 80,673 persons occupying approximately 15,849 households [11, 17].

78 **2.4 Sample Size and Sampling Procedures**

79 Simple random sampling technique based on random numbers generated on a scientific calculator,
 80 was used to select four Locations and households adjacent to Chepalungu forest for the study. The
 81 Locations selected were Bing'wa, Siongiroi, Ndanai, and Abosi.

82 The household sample size in each location was calculated based on formula equation 1 at 0.1
 83 margin error [18]:

$$84 \quad n = \left[\frac{N}{(1+Ne^2)} \right] \dots\dots\dots 1$$

85 Where

86 N= population size

87 n = sample size

88 e = margin error

89 Therefore the sample size in each Location was calculated based on the Location's available
 90 households. According to [19], there were 2010, 1820, 2003, and 1157 households in Bing'wa,
 91 Siongiroi, Ndanai and Abosi Locations respectively.

92 Therefore, using equation 1, the sample size in:

$$93 \quad \text{Bing'wa} = n = \left[\frac{N}{(1+Ne^2)} \right] = \left[\frac{2010}{(1+(2010 \times 0.1^2))} \right] = 95.67 \text{ households} = 96 \text{ households}$$

$$94 \quad \text{Siongiroi} = n = \left[\frac{N}{(1+Ne^2)} \right] = \left[\frac{1820}{(1+(1820 \times 0.1^2))} \right] = 94.79 \text{ households} = 95 \text{ households}$$

$$95 \quad \text{Ndanai} = n = \left[\frac{N}{(1+Ne^2)} \right] = \left[\frac{1459}{(1+(142003 \times 0.1^2))} \right] = 93.58 \text{ households} = 94 \text{ households}$$

$$96 \quad \text{Abosi} = \left[\frac{N}{(1+Ne^2)} \right] = \left[\frac{1143}{(1+(1143 \times 0.1^2))} \right] = 91.95 \text{ households} = 92 \text{ households}$$

97 **2.5 Data Collection Procedures**

98 The household questionnaires were administered to the sampled household samples.

99 **2.6 Data Analysis and Presentation**

100 The responses from household questionnaires were coded and analyzed by identifying relevant
 101 qualitative activities and outcomes. The quantitative data was cleaned, coded and analyzed with

the help of SPSS version 21 software and using both descriptive and inferential statistics as described below.

The test for variations in challenges hindering farm forestry development was carried out using Chi-Square test of association. Chi-Square Test of association was used to identify factors that are significantly associated with the various farm forestry practices. The null hypothesis was rejected if the computed P was less than or equal to 0.05.

3. RESULTS AND DISCUSSION

3.1 Results

It was found that inadequate capital and competing land uses are the most notable challenges facing adoption of farm forestry among farmers living around Chepalungu Forest. Damages by animals affect home-gardens, riparian planting, wind breaks, scattered trees and shade trees. Destruction by human man was associated with home-gardens and wind breaks (Table 1).

Table 1: Challenges affecting adoption of farm forestry practices in Chepalungu

Types of Farm Forestry	Response (%) on Major Challenges								
	Damaged by animals	Damaged by man	Seed acquisition	Tree nursery problems	Managerial problem	Inadequate capital	Competing land use	Natural calamities	
Home-garden	52.9	5.9	0.0	0.0	0.0	23.5	5.9	11.8	
Riparian	50	0.0	0.0	0.0	0.0	25.0	25.0	25.0	
Planting	0.0	40.0	0.0	0.0	0.0	0.0	21.4	40.0	
Wind breaks	0.0	0.0	0.0	42.9	0.0	28.7	0.0	0.0	
Hedgerowing	64.3	0.0	0.0	21.4	28.6	7.1	0.0	7.1	
Scattered Trees	66.7	0.0	0.0	0.0	0.0	0.0	33.3	0.0	
Shade Trees	0.0	0.0	28.6	0.0	0.0	21.4	28.6	0.0	
Boundary planting	0.0	0.0	0.0	0.0	87.5	0.0	12.5	0.0	
Woodlot									

Chi-Square test of association indicated that there was a significant association between types of farm forest practices and potential challenges affecting adoption of farm forest practices (Table 2).

120 **Table 2: Chi-Square tests of association**

	Value	df	Exact Sig. (2-sided)
Pearson Chi-Square	5.308E2 ^a	63	0.000

121

122 It was also found that seed acquisition had no significant association with any type of farm forestry
 123 practices. Tree nursery problems were significantly associated with hedgerawing, scattered trees on
 124 farms and woodlot practices (Table 3).


125 **Table 3: Chi-Square tests on individual challenges in Chepalungu**

Types of Farm Forestry	Response on Major Challenges							
	Animal damage	Mandam age	Seed acquisition	Nursery	Managerial	Capital	Competing land use	Natural calamities
Home-garden	X ² = 23.54, P<.001	X ² = 18.097 P=.045	X ² = 4.042, P=.050	X ² = 12.762, P=.077	X ² = 45.500, P=.094	X ² = 17.225, P=.002	X ² = 3.752, P=.453	X ² = 24.798, P=.<.001
Riparian Planting	X ² = 41.017, P=.001	X ² =9.007, P=.134	X ² = 5.223, P=.091	X ² = 8.073, P=.067	X ² = .318, P=.980	X ² = 41.017, P=.001	X ² = 17.717, P<.001	X ² = 15.517, P=.037
Wind breaks	X ² = 21.401, P<.001	X ² = 22.500, P<.001	X ² = 1.755, P=.670	X ² = 8.276, P=.292	X ² = 45.500, P=.094	X ² = .947, P=.824	X ² = 17.342, P=.040	X ² = 27.423, P=.020
Hedgerawing	X ² = 6.802 P=.301	X ² = 4.544 P=.395	X ² = 8.229 P=.327	X ² = 45.401, P=<.001	X ² = 6.766, P=.330	X ² = 10.541 P=.001	X ² = .545, P=.946	X ² = 9.171, P=.411
Scattered Trees	X ² = 44.870 P=.031	X ² = 9.155 P=.467	X ² = 1.053 P=.818	X ² = 10.111 P=.011	X ² = 67.300 P=.025	X ² = 16.855, P=.770	X ² = 3.333, P=.102	X ² = 11.001, P=.067
Shade Trees	X ² = 9.870 P=.007	X ² = 1.041 P=.820	X ² = 1.261 P=1.000	X ² = 3.709 P=.308	X ² = 91.500 P=1.000	X ² = .705, P=.970	X ² = 132.900 P=.001	X ² = 8.760, P=.094
Boundary planting	X ² = 2.238 P=.556	X ² = 3.709 P=.900	X ² = 13.350 P=.029	X ² = 4.012 P=.206	X ² = 2.715, P=.096	X ² =26.981 P=.009	X ² = 51.500 P=<.001	X ² = 4.981, P=.100
Woodlot	X ² = 8.602 P=.547	X ² = 1.494E2 P=.390	X ² = 73.018 P=.623	X ² = 3.720 P=.717	X ² = 10.955, P=.360	X ² = 14.900 P=.290	X ² = 12.271 P=<.001	X ² =11.759, P=.075

126

127 The majority of farmers practicing different types of farm forestry in the four locations obtain seedlings
128 from tree nurseries (Figure 2).

129



130

131 **Figure 2: Respondents obtaining seedlings from tree nurseries**

132 Tree nurseries around Chepalungu forest were found to be small in size (Plate 1) due to unavailability
133 of essential production factors like water supply, quality soils, poor road network, and distance to the
134 planting site. The experience of seed collectors is also low, eventually jeopardizing the quality of
135 seedlings supplied to farmers.


136



137

138 **Plate 1: A tree nursery in Bing'wa Location in April, 2016**

139 **Managerial Problems;** Tree Managerial problems were significantly associated with woodlot and
 140 scattered trees on farm forestry practices (Table 4). Over 36% of respondents in the four locations
 141 were unaware of silvicultural practices like pruning, thinning and pollarding (Figure 3).



142

143 **Figure 3: Unawareness level of silvicultural practices in Chepalungu as at 2016**

144 **Inadequate Capital:** Inadequate capital was significantly associated with home-garden, riparian
 145 Planting, wind breaks, shade trees, and boundary planting.

146 **Competing Land Use:** Competing land use was significantly associated with riparian planting, wind
 147 breaks, shade trees, boundary planting, and woodlot practices.

148 **Natural Calamities:** Natural calamities were significantly associated with home-garden, riparian
 149 planting, and wind breaks.

150 **3.2 Discussion**

151 The most notable challenges facing farmers practicing farm forestry could be categorized into
 152 Damages by Animals and Human beings, Seed Acquisition and Tree Nursery Problems, Managerial
 153 Problems and Inadequate Capital, Competing Land Use, Natural Calamities.

154 **3.2.1 Damages by animals and human beings**

155 Domestic animals including sheep, goats and cattle destroy farm forest trees through mechanical
 156 means like stepping on them and or feeding on shoots. On the other hand, human beings engage in
 157 various activities that either affects the on-farm trees either directly or indirectly. Direct destructions
 158 include cutting of trees for various purposes, and use of mechanical cultivation like tractors and
 159 combined harvesters. Indirect destruction of farm-trees entails making the soils unsupportive to trees

through soil pollution by using chemical fertilizers, herbicides, and insecticides. The results concur with [20] that some chemical fertilizers make it difficult for micro-organisms in the soil to produce nutrients naturally; hence making soils unsupportive to plant growth. Globally, farmers are clearing scattered trees on their farms to ensure easy mechanized farming [21].

3.2.2 Seed acquisition and tree nursery problems

Seed acquisition is not a significant challenge because above 82% of farmers obtain planting materials from locally established private nurseries. The significant challenge emanates from tree nursery production problems like inadequate water supply, poor road network and inexperienced seed collectors. Such challenges emerge because privately established tree nurseries in Chepalungu Forest neighborhoods are small in size, implying that owners have inadequate capital to institute all necessary conditions that will ensure production of quality seedlings. According to [22], farm forestry in developing countries experience the supply of low quality seeds, seedlings, cuttings or propagules, resulting to unsatisfactory benefits in terms of fruit, timber and shade quality. [23] adds that inadequate experience among local seed collector in Africa is alarming, and eventual results is low quality planting materials that lowers the survival rates of trees and effectiveness of agroforestry initiatives.

3.2.3 Managerial problems and inadequate capital

Inadequate managerial knowledge on silvicultural practices like pruning, pollarding, and thinning is a significant challenge facing farm forestry in Chepalungu Forest neighborhood. This is coupled with inadequate capital to undertake adequate land preparation, and disease control among other tree management practices. The results are coincides with [24] that the level of practicing silvicultural practices varies from one location to the other as some farmers practice better silviculture than others. This is because of variations in the understanding of silvicultural practices. Poorly managed on-farm forests, affects the quality of output products and services that eventually discourages farmers from engaging in farm forestry [23].

3.2.4 Competing land use

Farm forestry faces significant competition from other profitable land uses like crops. Therefore, since farm forestry practices like scattered trees on-farm competes with crops for limited resources water and nutrients, farmers prefer cutting trees to maximize their yields on farm crops. These findings

concur with that of [25] who found that on-farm forestry was reducing in Trans-Nzoia County as farmers preferred maize production that was perceived to be more profitable. Also [26, 25, 28] found that adoption of agroforestry was hampered by preference of other profitable land use activities like crop farming.

3.2.5 Natural calamities

Natural calamities including strong winds and pests like aphids which attack Cypress are significantly challenging the existence of farm forestry in the study area. This is because they destroy already established trees on farms resulting in huge losses. These losses therefore discourage farmers from establishing new farm forest practices. According to [29], natural disasters including fires, droughts, cyclones & typhoons, and diseases have both direct and indirect impacts on farm forestry. Floods make land unsuitable for vegetation growth, while diseases and hurricanes destroy trees on-farms.

In general, the findings of this study revealed that education, capacity building and training is very important in enhancing the success of farm forestry. This is because some of the challenges like competing and use, managerial problems and inadequate capital can be curbed through education, capacity building and training on the proper arrangement of trees on farm and cheapest practices available.

4. CONCLUSIONS

The main challenges affecting the farm forestry by communities living adjacent to Chepalungu Forest include: damages by animals and human beings, seed acquisition and tree nursery problems, managerial problems and inadequate capital, competing land use, natural calamities. If these challenges are left unaddressed, they will threaten the conservation and regeneration efforts directed to Chepalungu Forest. This is because such challenges will discourage adoption of farm forestry, leading to a situation where the community will entirely depend on Chepalungu Forest. Total dependence on the forest will lead to unsustainable harvesting in the forest that will eventually lead to forest degradation.

REFERENCES

1. Chao S. Forest Peoples: Numbers across the world. Baltimore: Johns Hopkins University. 2012.

- 217 2. Food and Agriculture Organization. The State of Food and Agriculture. *Social protection and*
218 *agriculture: Breaking the cycle of rural poverty*. Rome: FAO. 2012
- 219 3. Foley G, Bernard G. *Farm and Community Forestry*. Earth scan Energy Information
220 Programme, International Institute for Environment and Development, London. 1984.
- 221 4. Government of Kenya. *Forest Policy, 2014*. Nairobi: Government Printer. 2014.
- 222 5. Muchiri JM, Ascolillo L, Mugambi M, Mutwiri T, Ward HD, Naumova EN, Egorov AI, Cohen S,
223 Else JG, Griffiths JK. Seasonality of *Cryptosporidium* oocyst detection in surface waters of
224 Meru, Kenya as determined by two isolation methods followed by PCR, Journal of Water and
225 Health. 2009; 7, 67-75.
- 226 6. Wass P.(Ed.). *Kenyas indigenous forests; Status, Management and conservation*, IUCN,
227 Glands, Switzerland, and Cambridge, U.K. 1995; Xii + 205pp.
- 228 7. Graeuba B, Chappellb C, Wittmand H, Ledermanne S, Bezner K, Gemmill-Herrena B. The
229 state of family farms in the world. *World Development*, 20(2016), 1-15.
- 230 8. Kenya Forest Service. *The state of Chepalungu natural forest*. 2003.
- 231 9. Langat CK, Koech E, Kiplagat A, Owino YO. Threats and conservation incentives for
232 sustainable management of South West Mau Forest, Kenya. *Asian Journal of Environment &*
233 *Ecology*. 2018; 7(1): 1 -12. DOI: 10.9734/AJEE/2018/42130
- 234 10. Owino YO, Sirmah PK, Hitimana J. The prediction of leaf biomass production from *Faidherbia*
235 *albida* in semi arid land, Pokot County, Kenya. *Asian Journal of Research in Agriculture and*
236 *Forestry*. 2018; 1(2): 1 -10. DOI: 10.9734/AJRAF/2018/ 40867
- 237 11. Langat D, Sigu G, Wanyondu J, Bii W. *State of Chepalungu Forest*, KFRI. 2002.
- 238 12. Harrison S, Herbohn J. *Agroforestry and farm forestry support systems to assess the viability*
239 *of whole-farm and regional agroforestry enterprises*. Rural Industries Research and
240 Development Corporation. 2008
- 241 13. Sunderlin WD, Angelsen A, Wunder S. *Forests and poverty alleviation. In State of the world's*
242 *forests 2003. Rome, Italy: Food and Agriculture Organization of the United Nations*. 2003
- 243 14. Dewes PA, Saxena NC. *Tree planting and household land and labour allocation: case studies*
244 *from Kenya and India. In Arnold, J.E.M.,Dewes, P.A. (eds) Tree Management in Farmer*

- 245 *Strategies. Response to agricultural intensification*. Oxford Science Publication. 1995; pp 242-
246 267.
- 247 15. Langat D, Maranga E, Cheboiwo J, Aboud A. Forest use and dependence by forest adjacent
248 households on East Mau forest ecosystem, Kenya. *Journal of Agricultural Economics*. 2015;
249 3(10): 326-337
- 250 16. Jaetzold R, Hornetz B, Shisanya C. (Eds). *Farm management handbook*. Ministry of
251 Agriculture, Kenya. 2010.
- 252 17. Ministry of Agriculture. *Economic Review of Agriculture 2010*. Kenya. 2010.
- 253 18. Guilford JP, Frucher B. Fundamental statistics in psychology and education. New York: MC
254 Graw-Hill.1973.
- 255 19. Kenya Bureau of statistics. 2009 *Population by district*. Retrieved 2 October 2012, from
256 <http://www.google.co.ke/search?q=2009 Population By District>. 2010.
- 257 20. Maynard DG, Paré D, Thiffault E, Lafleur B, Hogg KE, Kishchuk B. How do natural
258 disturbances and human activities affect soils and tree nutrition and growth in the Canadian
259 boreal forest? *Environmental Reviews*. 2016. 22(2):161-178.
- 260 21. Zubair M, Garforth C. *Farm level tree planting in Pakistan: The role of farmers' perceptions*
261 *and attitudes*. AFTA 2005 Conference Proceedings. 2005.
- 262 22. Nyoka BI, Roshetko J, Jamnadass R, Muriuki J, Kalinganire A, Lillesø B, Beedy T, Odhiambo
263 K. *Positive deviance in the adoption of agroforestry technologies within Lower Nyando Basin*,
264 Kenya. Nairobi: Kenyatta University. 2014.
- 265 23. Roshetko J, Mulawarman M, Dianarto A. Tree seed procurement-diffusion pathways in
266 Wonogiri and Ponorogo, Java. *Small-scale Forestry*. 2008. 7(4): 333-352.
- 267 24. Walters B, Sabogal C, Snook LK, Almeida E. Constraints and opportunities for better
268 silvicultural practice in tropical forestry: an interdisciplinary approach. *Forest Ecology and*
269 *Management*, 209 (2005): 3–18.
- 270 25. Wafuke S. *Adoption of agroforestry technologies among small scale farmers in Nzoia*
271 *Location, Lugari District, Kenya*. ICRAF. Nairobi, Kenya. 2014
- 272 26. Mandila B, Hitimana J, Kiplagat A, Mengich E, Wekesa T. Prevalence and adoption of
273 agroforestry technologies and practices in Semi-Arid Regions of West-Pokot County, Kenya.
274 *Research Journal of Agriculture and Forestry Sciences*. 2015; 3(6): 6-15.

- 275 27. Namaswa WT, Mbego J, Muisu F, Mandila B. *Firewood accessibility among rural and urban*
276 *households in Trans-Nzoia and West Pokot Counties, Kenya*. International Research Journal
277 of Environmental Sciences. 2016; 5: 1-11.
- 278 28. Nkamleu G, Manyong V. Factors affecting the adoption of agroforestry practices by farmers in
279 Cameroon. *Small-scale Forestry*. 2005; 4(2):135-148.
- 280 29. Sivakumar K. *Impacts of natural disasters in agriculture: An overview*. Geneva: World
281 Meteorological Organization. 2016.