

# Assessment of off-season water supply situation: The case of Abetifi in the Kwahu-East District of Ghana

John Adu Kumi<sup>1\*</sup>, Philemon Obiri Afari<sup>1</sup> and Beatrice Nyarko<sup>2</sup>

<sup>1</sup>Department of Environmental and Natural Resources Management, Presbyterian University College  
Ghana, Akropong-Akuapem, Ghana

<sup>2</sup>Korle Bu Teaching Hospital, Korle Bu, Accra, Ghana

## ABSTRACT

The main aim of this study was to investigate the characterizing features of off-season water supplies in the Kwahu-Abetifi Township situated in the Kwahu-East District of Ghana.

A descriptive cross sectional survey was employed for the research.

The study was conducted in the Kwahu East district of Ghana between August 2017 and December 2017.

The stratified sampling technique was employed to select 200 respondents for the study. Respondents were interviewed using prepared and validated questionnaires. Data obtained from the completed questionnaires, were analyzed using the statistical Package for the Social Sciences (SPSS).

The community people including school-going children in the study area, trek long distances and stay in long queues to access water. They however spend long hours in accessing water for use in their homes during the dry season. Durations respondents used in accessing water and returning to their homes were as follows: Approximately 15 minutes (10%, n=200); 20 minutes (16%, n=200); 35 minutes (22%, n=200); and more than 40 minutes (52%, n=200). Nevertheless, the accessed water were perceived to be good (68%, n=200) and were used for several domestic purposes including drinking (19%, n=200), cooking (47%, n=200), bathing and washing (34%, n=200). The study underscored the need for the revamp of broken down water supply systems in the study area and expansion of existing ones by Government to ease and improve accessibility to adequate and potable water supplies mostly during the dry season.

**Keywords:** Potable water supply, off-season, accessibility, proximity, affordability

## 1. INTRODUCTION

Ensuring the provision of adequate and potable water supplies to communities across all facets of life around the world is very critical. Good health, livelihood enhancement, sustainable development among other key aspects of life all hinge on the supply of clean and sufficient quantities of water that can be reliable at all times [1,2]. Worldwide, it is estimated that, the total cost of water insecurity affecting the global economy is US\$500 billion [3]. Over the years, there have been incessant concerns at the international, regional, sub-regional and grass root levels concerning issues of potable water supplies to people. Aside the concerns, water supplies in most parts of the world face challenges including but not limited to scarcity. It therefore becomes very prudent to address water related issues in the context of scarcity and security, sanitation and health crisis with infrastructure deterioration and destruction to reverse the threatening supply and quality trends.

Despite the enormous efforts to improve the supply of potable water to people worldwide, 663 million people suffer inaccessibility to 'improved' drinking water sources [4]. Moreover, the 'improved' water is perceived by many as either not safe, unreliable or not affordable. In respect of water safety, levels of arsenic above World Health Organization (WHO) standards in drinking water consumed by 45 million people in Bangladesh have been documented [5].

In Ghana, scarcity situations have not been different as some communities continue to suffer stern water shortages usually during the lean season. Several community water connectivity exist, but equally an appreciable number of communities continue to receive their share of supply challenges, perhaps as a result of frequent system break down and other factors.

## **1.1 Objectives of the study**

The main objectives of the study were:

To determine the challenges that face water supply accessibility in Abetifi-Kwahu in the Eastern region of Ghana during the dry season.

To ascertain the perceived quality of water supplied in Abetifi-Kwahu in the Eastern region of Ghana during the dry season.

## **2. The global water cycle and situation**

Water, being an important element of life, cycles between land, the oceans and the atmosphere through solid, liquid and gaseous phases. Precipitation from the atmosphere gets to land, underground aquifers and oceans. Rivers get their supplies from groundwater as water moves through sediments and rocks. Rivers intend provide freshwater discharge into seas. Water finally evaporates or transpires from the ocean and land into the atmosphere to complete the cycle.

Earth's water resources including rivers, lakes, oceans and underground aquifers are under stress in many regions worldwide. Of the 70% portion of the Earth's water cover, only 2.5% is freshwater. This limited resource is to support an estimated 9.7 billion of the world's population by 2050 [6]. In sub-Saharan Africa, high population growth triggers high consumption rate of this limited resource. Worldwide, some regions are endowed with freshwater than others as a result of global circulation patterns which translate to distinct wet and dry phases of multi-annual climate cycles.

Despite the fact that developed countries have more water available than most of the countries in Africa, Asia and the Pacific, some areas are water-stressed because the available water sources are being depleted at very high rates [7]. It is envisaged that, global water demand would increase by 50% by the year 2030 [8]. Water supply situation, moreover is estimated to be worsened with an approximately 3.9 billion (over 40%) of the world's population, to dwell in severely water-stressed basins [6]. Vicinities with water-stressed situations, consequently makes people seek water from very distant sources. With water scarcity affecting over 40% of the world's population, there is the need for urgent interventions and implementation of highly workable strategies [9].

The United Nations and the international community however, set targets to reduce the number of persons with inadequate potable water and sanitation by 2015. Meeting this target was to provide an extra 260,000 people per day with clean drinking water and 370,000 people per day with improved sanitation through the year 2014 due to the escalating global water demand [10].

## **2.1 Community water supplies in Ghana**

In most rural communities of Ghana, people rely on all sorts of water sources during the wet season including streams, rivers, dug wells, ponds, boreholes, rainfall, etc but face the challenge of dryness or severe drought getting to the end of the year. Most commonly, people who dwell in these areas resort to these available water sources, which might not be potable but use them unknowingly at the highest health risk [11]. Provision of clean water, is thus imperative in ensuring that people do not suffer likely detrimental diseases and other health risks.

79 Again, personal hygiene is very essential, and must not receive less attention because of water  
80 inaccessibility [8]. Per capita water consumption is reliance on accessibility, which could primarily be  
81 defined by distance, time, reliability and cost [12]. There is a higher health benefit when water is  
82 rather at the household level. Interventions by Governments, International organizations, and Non-  
83 Governmental Organizations have made it possible for several rural communities in Ghana to boast of  
84 pipe-borne water for which their accessibility, reliability, permanence, and cost effectiveness are also  
85 matters of concern.

86 Potable water accessibility challenges limit personal and household hygiene, which intend affect  
87 public health. A supply system that would ensure the everlasting needs of people and also accessible  
88 to all without denying the living poor and the healthy living is key for the development of human lives.  
89 There is the utmost need for an affordable and efficient rural water supply system that is sustainable  
90 and hinges on an effective logical management framework [13].

91 Sustainable water management is very fundamental in realizing most of the sustainable development  
92 goals most especially Sustainable Development Goal (SDG) 6. Thus ensuring the sustainability of  
93 rural water supplies through effective management systems is worth considering. This will make  
94 people and their immediate environments continue to obtain the right quantities of potable water for  
95 growth and development. In most instances, rural water systems unreliability occur due to system  
96 breakdown, financial constraints, lack of sense of responsibility for service payment leading to  
97 systems failure, poor or no system evaluation and the failure in implementing evaluation  
98 recommendations. These factors limit the efficient performance of rural water supply systems.  
99

### 100 **3.1 DESCRIPTION OF STUDY AREA**

101 The study area, Abetifi-Kwahu is located in the Kwahu-East District of the Eastern Region of Ghana. It  
102 is geographically located on (6° 40' 0" N, 0° 45' 0" W). The over 15,000 populace of the study area  
103 are engaged in several economic activities including farming, trading and public service work. Abetifi-  
104 Kwahu as the name goes has been and still known for its severe perennial dry season water crises  
105 making life difficult for its inhabitants during this portion of the year.

106 The town used to be connected to the water lines of the Kotoso water treatment station making it  
107 easier for people who lived mostly in bungalows accessed water in their homes at the time. For the  
108 past two decades, the supply lines connected to this water station are damaged and have not been  
109 fixed, making the people access quantities of water that are described as scarce and frustrating to  
110 find during the dry season. Towns like Nkwatia and Mpraeso are connected to the Ghana Water  
111 Company Limited lines but Abetifi has not yet seen its share of this connectivity. During the dry  
112 season, rivers, streams as shown in Figure 1 below dry up leaving only the few constructed bore  
113 holes with standing pipes and dug wells for the use of the over 15,000 inhabitants of the community.




Fig.1. Sketch map of study area showing water sources

### 3.1.1 Climate and rainfall

Lying within the semi-equatorial region, Abetifi is characterized with two rainy seasons in a year, the major and minor rainy seasons. The major rainy season starts from April and ends in July whilst the minor starts from September and ends in October. Annual rainfall averages between 1580mm and 1780mm. The town experiences warmth and dryness between November and March as it is hit by the Tropical Continental (cT) air mass [14]. Within the month of February, the highest form of dryness with 23mm of rainfall is experienced. Known as the highest inhabited point in Ghana, the town is located about 2080ft above sea level and prides itself with a very serene weather condition with monthly temperature averaging 30°C in the dry season which falls to 26°C during the wet season. Relative humidity ranges from 75% to 80%.

### 3.1.2 Vegetation

The town lies within the Semi-Deciduous forest zone of Ghana [14]. The vegetation is characterized by a dense tree cover with trees shedding off their leaves mostly during the dry season. There exist the lower, middle and upper layers in terms of the tree coverage in the forest. Specifically, these layers include the canopy, understory layer, shrub layer, herbaceous layer and the forest floor. The canopy, the uppermost layer of the vegetation, consists of the upper leaves and branches of the tallest trees. Beneath the canopy, the understory layer consists of small, shorter trees. Beneath the understory layer is the shrub layer consisting of woody vegetation tending to be less than 10 feet high. The herbaceous layer is found beneath the shrub layer and consists of small, soft-stemmed plants such as wildflowers and ferns, as well as grasses. The forest floor, the lowest layer of this forest vegetation, consists of fallen leaves, twigs, and branches, as well as small plants such as mosses and lichens.

The forest vegetation is rich with several flora most of which, are of immense commercial and scientific values. Trees of economic importance which can be identified in the area include Odum, Wawa, Mahogany, etc [14]. Moreover, trees of commercial and scientific values, which abound in the forest, include Onyina, Emire, Ofra, Bompaghya, Kyenkyen, etc. Until a few centuries ago, the impact of human populations on both the vegetation and cohabiting species was minimal because of small population. In recent times, destruction of this vegetation in the study area caused by human population and economic pressures is considered a major threat to the natural environment. The natural vegetation cover continues to deteriorate as a result of activities including farming, logging of wood for timber, indiscriminate bush fires mostly during the dry season. Such activities change the natural vegetation which in turn affect the dynamics of biodiversity in the area.

148


**3.1.3 Soil**

149 Described as deeply weathered soils, forest ochrosols as the name goes are the type of soils found in  
150 the study area [14]. They consist of thin (about 20cm), dark greyish brown, humus-stained, sandy  
151 loam and silt loam topsoil which are usually moderate fine granular in structure and friable in  
152 consistency. The subsoil are thick, often more than 120cm thick over the weathered substratum. The  
153 texture of the subsoil varies and may be sandy clay loam, silty clay loam, sandy clay or silty clay with  
154 common to many (10-40%) quartz gravels and stones and hard iron and manganese dioxide  
155 concretions [15]. The forest ochrosols are generally slightly to moderately acid in the topsoil (pH 6.5-  
156 5.1 in 1:1 soil: water ratio). With good chemical properties, the soils are fertile and support the  
157 production of food and cash crops including cassava, yam, plantain, cocoa, coffee, etc.

**4. METHODOLOGY**

**4.1 Sampling and sample technique**

160 The stratified sampling technique was used to select the study sample from the study population [16,  
161 17]. This technique was chosen in order to evenly spread the sample across the entire population.  
162 Stratified sampling ensures a greater degree of representativeness, thereby reducing the probable  
163 sampling error. It ensures that different groups of population are adequately represented in the  
164 sample. It combines homogeneity and heterogeneity at different levels. With this technique, the study  
165 area was divided into five (5) separate groups called strata with residential location as criterion (Figure  
166 2). In each stratum, forty (40) respondents were selected by simple random sampling employing the  
167 lottery method. The lottery method was done by preparing a list of names and contact details of 150  
168 persons which constituted the sample frame for a particular stratum. Each person's name was then  
169 transferred from the sample frame (that is, the list of 150 persons) and then put onto separate pieces  
170 of papers. The pieces of papers were placed in a container and thoroughly mixed. The required  
171 number of respondents for a particular stratum (that is, 40), were selected without looking. The names  
172 selected constituted the simple random sample. The same random sampling procedure, was done for  
173 the four (4) other strata.



174

175 **Fig.2. Layout of Abetifi Township showing various strata where respondents were**  
176 **randomly selected** **Source: The Authors, 2018**

177



## 4.2 Primary data collection

The study employed a descriptive cross-sectional survey to collect primary data. In all, 200 respondents from individual households including institutional bungalows, flats, compound houses, semi-detached houses, etc were contacted and interviewed. Both qualitative and quantitative data were collected from the respondents by the use of a set of validated semi-structured questionnaires.

## 4.3 Data analysis and presentation

The data was keyed into the Statistical Package for the Social Sciences (SPSS) and analyzed using the interpretative technique to analyze observation and interview information obtained. The descriptive statistical method was used in summarizing and presenting data in the form of tables and numerical measure (percentages) in an informative way.

# 5. RESULTS AND DISCUSSION

## 5.1 Demographic characteristics of respondents

### 5.1.1 Gender of respondents

Majority of the respondents interviewed were females represented by a proportion of 54% of the total number of the interviewees as shown in Table 1 below. On the other hand, there was 46% male representation for the study sample. Commonly, women and girls are known to be involved in the handling and management of household water for several domestic purposes. Addressing the needs of females in respect of the quantity and quality of water supply is worth it.

**Table 1. Gender distribution of respondents**

Gender of respondents	Number of respondents	Percentage of respondents
Male	92	46
Female	108	54
Total	200	100

*Source: Field survey, 2017*

### 5.1.2 Age of respondents

Majority of the respondents fell under the age group of 31-40 years represented by 34% of the study sample as shown in Table 2 below. This showed that majority of the respondents reached were in the parenting and working ages. Persons in this age group use water for domestic activities at the household level and other socio-economic activities. Moreover, 26% and 22% of the respondents were in the age groups of 21-30 years and 11-20 years respectively. The least of respondents' representation in terms of age was the over 40 years age group representing 18% of the study sample. In all, there was a fair representation of all age groups for the study.

**Table 2. Age distribution of respondents**

Age of respondents (years)	Number of respondents	Percentage of respondents
11-20	44	22
21-30	52	26
31-40	68	34
>40	36	18
Total	200	100

*Source: Field survey, 2017*

### 5.1.3 Duration of residency of respondents

Table 3 below indicates that majority of the respondents (24%) have stayed in the study area between 31-40 years. Twenty-three percent (23%) and 21% of the respondents have lived in the study area within the periods, less than 10 years and 21-30 years respectively. Eighteen percent (18%) of the respondents have lived in the area within 10-20 years with the least of the respondents (14%) having been residents of the study more than 40 years. The number of years respondents have stayed in the study area was very important in order to obtain information that tells the real situation in terms of water supply during the dry season.

**Table 3. Duration of residency of respondents**

Duration of residency of respondents (years)	Number of respondents	Percentage of respondents
<10	46	23
10-20	36	18
21-30	42	21
31-40	48	24
>40	28	14
Total	200	100

*Source: Field survey, 2017*

### 5.2 Proximity to water supply points

Public Stand posts made up of several taps connected to a service line (a borehole) as described by one author [18] and dug wells are primarily patronized by the folks in the study area during the dry season as the available streams are dried up. Some authors [19] describe a reliable service as the one which is easy to access without one going through unnecessary stresses. Results of the study refuted this assertion; water accessibility at supply points in the study area has always been faced with challenges during the off-season period. Proximity to stand posts and dug wells have invariably been a major challenge in rural communities for which the study area was not an exemption. In most times, the community people trek long distances before accessing the available water.

The study revealed that 16% and 22% of the respondents spent approximately 20 minutes and 35 minutes respectively to access water and return to their homes (Table 4). A few (10%) claimed they spent approximately 15 minutes doing same. A relatively larger proportion of the sampled population (52%) rather spent more than 40 minutes to access water from the available supply points and return to their homes. The longer duration could have possibly been attributed to longer distances one has to pursue in accessing water amid queuing times. Those who may find queuing to access water frustrating leave their containers in the custody of water sales personnel who man the stand pipes so they could be served in their absence. Despite the fact that such people do not waste time waiting at the water supply point, there is always a longer time before they are able to access the water again in such a water-stress period. The implication is that, households have to manage quantities of water for a period before they could access the water supplies again. Households with larger family sizes are the ones that suffer from this challenge the most.

**Table 4. Duration used in accessing water and returning to homes**

Duration for accessing water	Number of respondents	Percentage of respondents
Appx. 15 minutes	20	10
Appx. 20 minutes	32	16
Appx. 35 minutes	44	22
> 40 minutes	104	52
Total	200	100

*Source: Field survey, 2017*

### 5.3 Wholesomeness of accessed water during the dry season

As shown in Table 5 below, there were different forms by which people in the study area use accessed water. When asked, respondents gave several uses of accessed water. Respondents used accessed water for drinking (19%), cooking (47%), bathing and washing (34%).

**Table 5. Uses of water accessed by respondents**

Uses of accessed water	Number of respondents	Percentage of respondents
Drinking	38	19
Cooking	94	47
Bathing and washing	68	34
Total	200	100

*Source: Field survey, 2017*

The various uses of accessed water indicated by the respondents were not surprising as the majority of them (68%) claimed the supplied water was good with only a few (32%) making mention of its particulate nature (Table 6). One author [20] attested to the fact that water from dug wells in rural areas have over the time been challenged with the lack of quality tests and as such stands a higher risk of compromising people's health.

It is imperative for community water supplies to be of good quality to ensure the safety of consumers [19]. Consequently, researchers [21, 22, 23] have attributed many health challenges and diseases to unhygienic rural water supplies. Very importantly, ensuring supply of good quality water in their right quantities safeguards primary health of people as well as socio-economic development [24]. Compromising universal water, sanitation and hygiene is thus not what one would ever wish for.

**Table 6. Perceived quality of available water supply in the study area**

Perceived water quality	Number of respondents	Percentage of respondents
Good	136	68
Salty	0	0
Bad odour	0	0
Tainted water	0	0
Particulate water	64	32
Total	200	100

*Source: Field survey, 2017*

Globally, approximately 3.5 million deaths and a daily record of 1,000 child mortality occur as a result of unsafe water, poor sanitation and hygiene [25,26]. When the basic water accessibility level is not attained, achieving a proper personal hygiene as well becomes very difficult to envisage. Time and distance thus come into play in determining the volumes of water that people are able to access to meet their personal hygiene needs [12]. According to these authors, the levels at which people are able to access water for use can be linked to the assurance of their health (Table 7).

**Table 7. Summary of requirement for water service level (accessibility) to promote health**

Service level	Access measure	Needs met	Level of health concern
No access (quantity collected often below 5l/c/d)	More than 1000 m or 30 minutes total collection time	Consumption: cannot be assured. Hygiene: not possible (unless practised at source).	Very high



Basic access (average quantity unlikely to exceed 20l/c/d)	Between 100 and 1000 m or 5 to 30 minutes total collection time	Consumption: should be assured. Hygiene: hand washing and basic food hygiene possible; laundry/bathing difficult to assure unless carried out at source	High
Intermediate access (average quantity about 50l/c/d)	Water delivered through one tap on-plot (or within 100 m or 5 minutes total collection time)	Consumption: assured. Hygiene: all basic personal and food hygiene assured; bathing and laundry should also be assured	Low
Optimal access (average quantity 100l/c/d and above)	Water supplied through multiple taps continuously	Consumption: all needs met. Hygiene: all needs should be met.	Very low


**Source:** Howard and Bartram, 2003 [12]

## 5.4 Community water costing and reliability

Most often, water supply systems have broken down primarily as a result of managerial as well as financial challenges. Researchers have over the years explained and made it clear on the essence of developing a more efficient system. A key factor in ensuring the sustainability of rural water supply systems is the ability and willingness of community people to pay for the service provided. As indicated in figure 3 below, reliable payments for water service will ensure that community water supply management framework works effectively. In that case, such water supplies must be potable, reliable, and easily accessible without spending longer time in accessing them.

The challenge of rural water supply systems sustainability transcends local and regional boundaries. There is however the utmost need for full community involvement and support, in the project during its design, construction and management in order to ensure its sustainability. These when done, would go a long way in ensuring the financial as well as the managerial strength of the system. Good financial management certainly needs to be strictly adhered to in order to cater for the operational and maintenance needs (O&M) of water supply systems and as well ensure their robustness and sustainability (Figure 3).

At the time of the study, a 25 litre container full of water cost 40 Ghana pesewas (US\$0.08) with a 30 litre container going for 50 Ghana pesewas (US\$0.10) at water stand posts. The respondents however described these prices as fair and were willing to pay without hesitation. Apparently, the volumes of water accessed by the respondents were not affected by cost but perhaps other factors [27]. Some respondents had shown high interest in household connectivity to the Ghana Water Company lines and were ready to pay for bills they would receive from the company if connected to its lines whilst others called for the construction of more mechanized boreholes at vantage points for easy accessibility of the greater majority during the dry season. Apparently, this was in line with [28] who had argued the need to reduce water accessibility distance and encouraged household connections.



**Fig.3. A proposed structure of a sustainable rural water supply system (Source: The Authors, 2018)**

## 6. SUMMARY OF FINDINGS

(1) Majority of the people (52%, n=200) in the study area spend longer period (more than 40 minutes) in accessing water from available supply points before they return to their homes during the dry season.

(2) The quality of accessed water in the study area, were perceived to be good (68%, n=200) during the dry season.

## 7. CONCLUSION

During the dry season, people in the study area use a lot of time to obtain water from the available supplies for use. Sanitation and health aspects of the community folks could be jeopardized because the per capita water consumption at the household level would not be met with this stemming from insufficient water supplies for homes. Moreover, water supply in Abetifi-Kwahu during the dry season is a challenge to the extent that inhabitants who can only afford resort to purchasing water from neighbouring towns that are connected to the Ghana Water Company lines increasing cost of accessing water and consequently living standards during this period. On the other hand, accessed water during this period of the year in the study area, were perceived to be good.

## 8. RECOMMENDATION

Water should easily be accessible to all in their sufficient quantities and quality. There is the need for more stand posts and mechanized boreholes to be constructed at vantage points in the study community. Again, household connectivity to water supply lines should be considered, critically assessed and implemented by Government through the Ghana Water Company Limited and all relevant stakeholders. Water quality is very critical in ensuring proper public health and the prevention of water related diseases. It is therefore prudent that water resources used mostly for domestic purposes are highly accessible in their significant quantities and free from all forms of pathogenic materials. The authors of this study consequently recommend a study on the assessment of the water quality of the several standpipes, dug wells and ponds which the community people resort to as there is little or no work done in that regard.

## ACKNOWLEDGEMENTS

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors. The authors are however thankful to all the enumerators who assisted immensely in the study area stratification and data collection processes. A special thank you goes to Dr. Gifty Seiwaa Nyarko for her immense assistance in the manuscript preparation. Last but not least,

the authors are grateful to Mr. Kwaku Karikari Amoah, Head of Institutional Advancement and Consultancy Unit of the Presbyterian University College Ghana for proofreading the manuscript.

## COMPETING INTERESTS

Authors have declared that no competing interests exist. The company name used for this research is commonly and predominantly selected in our area of research and country. There is absolutely no conflict of interest between the authors and company because we do not intend to use this company as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the company rather it was funded by personal efforts of the authors.

## AUTHORS' CONTRIBUTIONS

All authors collaborated in carrying out this study. Authors JAK and POA designed the study and wrote the protocol. Authors POA and BN managed the data collection and statistical analysis. Author POA managed the literature searches and wrote the first draft of the manuscript. All authors read and approved the final manuscript.

## REFERENCES

1. IDLO. Water Tenure Reform and Public Access to Water as a Basic Need, International Development Law Organization. Voice of Development Jurists Series; 2006.
2. Quagraine EK, Adokoh CK. Assessment of dry season surface, ground and treated quality water in the Cape Coast Municipality of Ghana. *Environ. Monit. Assess.* 2010; 160:521-539.
3. WWAP (United Nations World Water Assessment Programme). The United Nations World Water Development Report 2016: Water and Jobs. Paris, UNESCO; 2016. Accessed 18 April 2017. Available: <http://unesdoc.unesco.org/images/0024/002439/243938e.pdf>.
4. Guha-Sapir D, Hoyois P, Below R. Annual Disaster Statistical Review 2013: The Numbers and Trends. Centre for Research on the Epidemiology of Disasters (CRED), Institute of Health and Society (IRSS), Université Catholique de Louvain, Brussels; 2014. Accessed 18 April 2017. Available: [www.cred.be/sites/default/files/ADSR\\_2013.pdf](http://www.cred.be/sites/default/files/ADSR_2013.pdf).
5. Flanagan SV, Johnston RB, Zheng Y. Arsenic in tube well water in Bangladesh: health and economic impacts and implications for arsenic mitigation. *Bull World Health Organ.* 2012; 90:839-846. Accessed 18 April 2017. Available: <https://www.scienceopen.com/document?vid=ea6da984-9275-45e6-8c85-efd78fb8adba>.
6. OECD. Environmental Outlook to 2050: the consequences of inaction. OECD; 2012. Accessed 15 May 2017. Available: <http://www.oecd.org/env/indicators-modelling-outlooks/oecd-environmental-outlook-1999155x.htm>.
7. UNEP (United Nations Environment Programme) GEO. Global Environment Outlook 3, London: Earthscan. 2002; pp. 150–177. Accessed 15 May 2017. Available: <http://www.unep.org/geo/geo3/english/pdf.htm>.
8. WWAP (World Water Assessment Programme). The United Nations World Water Development Report 4: Managing Water Under Uncertainty and Risk. UNESCO, Paris; 2012. Accessed 15 May 2017. Available: <http://www.unesco.org/new/en/naturalsciences/environment/water/wwap/wwdr/wwdr4-2012/>.
9. United Nations (n.d.). Water. Accessed 21 May 2017. Available: <http://www.un.org/en/sections/issues-depth/water/>.
10. World Health Organization. "Water, Sanitation, and Hygiene Links to Health: Facts and Figures"; 2005. Accessed 18 April 2017. Available: [http://www.who.int/water\\_sanitation\\_health/factsfigures2005.pdf](http://www.who.int/water_sanitation_health/factsfigures2005.pdf).
11. UNESCO. UNESCO water portal newsletter no. 161: Water-related diseases; 2006.
12. Howard G, Bartram J. Domestic Water Quantity, Service Level and Health. WHO; 2003. Accessed 20 April 2017. Available: [http://www.who.int/water\\_sanitation\\_health/](http://www.who.int/water_sanitation_health/).

13. Nguyen TN. An Evaluation of Evaluation Systems for Rural Water Supply and Sanitation Systems. International Journal of Environmental and Rural Development, IJERD. 2013; 4-2.
14. Kwahu East – Ministry of Food & Agriculture (n.d.). Accessed 21 May 2017.  
Available: [http://mofa.gov.gh/site/?page\\_id=1518](http://mofa.gov.gh/site/?page_id=1518)
15. Adjei-Gyapong T, Asiamah, RD. The interim Ghana soil classification system and its relation with the World Reference Base for Soil Resources. Rapport sur les Ressources en Sols du Monde (FAO); 2002. Accessed 15 May 2017.
16. Nketiah-Amponsah E, Aidam PW, Senadza B. Socio-economic determinants of sources of drinking water: some insight from Ghana. In Conference on International Research on Food Security, Natural Resource Management and Rural Development, University of Hamburg; 2009.
17. Wedgwood A, Sansom K. Willingness-to-pay surveys: a streamlined approach: guidance notes for small town water services. WEDC, Loughborough University; 2003.
18. Jordan TD. A handbook of gravity-flow water systems for small communities. London, Intermediate Technology Publications; 1984.
19. IRC, WHO. Management of operation and maintenance of rural water and sanitation programmes—a training package for managers and planners. Geneva, World Health Organization; 2000.
20. Charrois JWA. Private drinking water supplies: challenges for public health; 2010.
21. Esrey SA, Feachem RG, Hughes JM. Interventions for the control of diarrhoeal diseases among young children: improving water supplies and excreta disposal facilities. Bulletin of the World Health Organization.1985; 63(4): 757-772.
22. Esrey SA, Potash JB, Roberts L, Schiff C. Effects of improved water supply and sanitation on ascariasis, diarrhoea, dracunculiasis, hookworm infection, schistosomiasis, and trachoma. Bulletin of the World Health Organization.1991; 69(5): 609-621.
23. WHO. Health systems: improving performance, World Health Report, 2000, WHO, Geneva, Switzerland; 2000.
24. Cairncross AM. Health impacts in developing countries: new evidence and new prospects. Journal of the Institution of Water and Environmental Management.1990; 4(6): 571-577.
25. UNEP (United Nations Environment Programme). Healthy Environment, Healthy People. Thematic Report for the Ministerial policy review session of the Second Session of the United Nations Environment Assembly of the United Nations Environment Programme, Nairobi.2016; 23–27. Accessed 21 May 2017.  
Available:<http://wedocs.unep.org/bitstream/handle/20.500.11822/17602/K1602727%20INF%205%20Eng.pdf?sequence=1&isAllowed=y>.
26. United Nations. Sustainable Development Goals. Goal 6: Ensure access to water and sanitation for all, facts and figures; 2016. Accessed 21 May 2017.  
Available: <http://www.un.org/sustainabledevelopment/water-and-sanitation/>.
27. Howard G (ed).Water supply surveillance - a reference manual. WEDC, Loughborough University, UK; 2002.
28. WELL. Guidance manual on water supply and sanitation programmes, WEDC, Loughborough, UK; 1998.

## APPENDIX

### QUESTIONNAIRE

#### RESPONDENTS: COMMUNITY FOLKS IN ABETIFI-KWAHU, GHANA

*This study is aimed at finding out the opinions of community folks on the characteristics of off-season water supplies in Abetifi-Kwahu situated in the Kwahu-East District of Ghana. The study requires your truthful and honest opinion as a participant. As such, this request is voluntary so do not feel forced in any way in stating your views.*

*The research is purely for academic and development purposes and so your responses are assuredly confidential and anonymity is safeguarded.*

*Thank you for your participation.*

**INSTRUCTION: Please tick (✓) the appropriate answer**

#	SECTIONS	RESPONSES
<b>Demographics</b>		
<b>1</b>	Gender	1. Male 2. Female
<b>2</b>	Age at latest birthday	1. 11-20 2. 21-30 3. 31-40 4. More than 40
<b>3</b>	What is your highest level of education?	1. None 2. Primary 1-6 3. Middle/JHS 4. SSS/Technical 5. University/College 6. Non-formal 7. Others (Specify.....)
<b>4</b>	For how many years have you lived in this community?	1. Less than 10 2. 10-20 3. 21-30 4. 31-40 5. More than 40
<b>Water supply issues</b>		
<b>5</b>	What are the available water sources you have in your community?	1. Borehole 2. Dug well 3. Stream 4. Pond 5. Others (Specify.....)
<b>6</b>	Are all available water sources in your community accessible during the dry season?	1. Yes 2. No
<b>7</b>	What are the available water sources you have in your	1. Borehole

	community during the dry season period?	2. Dug well 3. Stream 4. Pond 5. Others (Specify.....)
<b>8</b>	In your view, do you feel there are challenges with accessing water sources in your community during the dry season period?	1. Yes 2. No
<b>9</b>	What is the proximity to available water sources in your community during the dry season?	1. Closer to my house 2. Far away from my house 3. Others (Specify.....)
<b>10</b>	Is the number of water sources available to you during the dry season period adequate?	1. Yes 2. No
<b>11</b>	Do you queue to access available water sources in your community during the dry season?	1. Yes 2. No
<b>12</b>	In your view, approximately how many minutes do you use to access available water sources and return to your home?	1. 15 minutes 2. 20 minutes 3. 35 minutes 4. More than 40 minutes
<b>Perceived water quality and health issues</b>		
<b>13</b>	How do you perceive the quality of the available water sources during the dry season?	1. Good 2. Salty 3. Bad odour 4. Tainted water 5. Particulate water
<b>14</b>	What is your UTMOST use of the water you are able to access during the dry season period?	1. Drinking 2. Cooking 3. Bathing and washing



15	Have you ever been taken ill after ingesting the available water during the dry season?	1. Yes 2. No
<b>Community water costing and reliability</b>		
16	In your view, do you think it is affordable to access water sources during the dry season?	1. Yes 2. No
17	In your view, do you think it is affordable to access water at stand pipes in your community during the dry season?	1. Yes 2. No
18	What is the cost of water at community stand pipes during the dry season?	.....
19	Do the community stand pipes frequently break down during the dry season?	1. Yes 2. No
20	In your view, do you think the management of the operations and maintenance of stand pipes in your community is good?	1. Yes 2. No
21	Are you always willing to pay for the water supply service in your community?	1. Yes 2. No
22	Would you prefer household water connectivity in your community?	1. Yes 2. No
23	Will you be willing to pay for bills you will receive from the Ghana Water Company after you have been connected to household connection?	1. Yes 2. No

**\*Kindly write any other comments and contributions you have concerning the questionnaire/Interview schedule at the back of the last page.**

487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500

501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
  
516  
  
517  
  
518  
  
519  
  
520