

1 *Original Research Article*

2

3 ***Impact of papaya seed soaking in different BA,***

4 ***colchicine and EMS solutions on germination,***

5 ***growth and chromosomal behavior***

---

6

7

8

9 **ABSTRACT (ARIAL, BOLD, 11 FONT, LEFT ALIGNED, CAPS)**

10

The present investigation was carried out during two consecutive seasons 2015 and 2016 in fruit nursery of faculty of Agriculture at Moshtohor, Benha University, in order to throw some spotlight on the impact of some chemical substances (Ethyl Methane Sulphonate – EMS10,20 and 30 ppm); (colchicine at 1,2and 3%) and (benzyl adenine BA at 1,2 and 3%) on seed germination %, seed germination rate, some seedling growth measurements and cytological examination of root tip of *Carica papaya* cv. Solo. The treatments were arranged in complete randomized block design with nine replicates (polyethylene bags), however, each replicate was represented by two papaya seedlings. The seedlings were divided into three categories according to their growth vigor, each category represented by three replicates for each treatment and subsequently each category sampled by 60 seedlings for all studied treatments. Seedling growth and chromosomal behavior as imported by the three studied chemical substances were evaluated on the 1st week of December. Data obtained revealed that both BA 2 % and BA 3 % increased significantly germination %, germination rate and growth measurements. On the contrary, the least significant increase was always in concomitant to EMS at 3 % and colchicine at 3 % during both experimental seasons. Moreover, EMS was more inhibitor of cell division followed by BA than Colchicine. This may be due to more damage resulted by BA and EMS affected on DNA replication during mitosis.

11

12 *Keywords: [Carica papaya, germination %, seed germination rate, growth*

13 *measurements, cytological examination, BA, Colchicine and EMS.]*

14

15

16 **1. INTRODUCTION (ARIAL, BOLD, 11 FONT, LEFT ALIGNED, CAPS)**

17

18 The papaya (*Carica papaya* L.) is cultivated for is ripe fruits, favored for people in the tropical region as breakfast fruit, and as ingredient in juice, jellies and preserves or cooked with young leaves and shoots as a vegetable plant. The fruit contains high level of papain; the proteolytic enzyme used for medical purposes and as a tenderizer for meat. The fruit, also, contains considerable quantities of vitamin A, B and C and about 10 % sugar. Fruits and seeds extract have pronounced bactericidal activity against *Staphylococcus aureus*, *Bacillus cereus* and *Escherischia coli* and the latex is used to remove freckles. Other parts such as bark is used for making rope while leaves are also used as a soap substitute supposed to remove stains.

26

27 Cytokinins can alter flower sex ratio in species with imperfect flowers. Cytokinins generally, increase the ratio of female flowers to male flowers which has implications for fruit production[1]. BA has also been used in the vegetable crop industry to alter flower sex ratios of monoecius and dioecious plants to increase the number of female flowers available to produce fruit [2].Exogenous cytokinins can promote an accumulation of chlorophyll and

28

29

30

31

32 promote the conversion of etioplasts into chloroplasts [3] even in dark grown seedlings. This  
33 may appear as a greening effect on ornamental crops which may be perceived as an  
34 increase in quality in green leaved crops and a decrease in quality in crops with other leaf  
35 colors. There is also some evidence that cytokinins can help increase the flower size of  
36 some plants. Cytokinins increased the size of petunia flowers [4]. In ferns however,  
37 cytokinins appear to induce maleness in the gametophytes [5]. The reduction in percentage  
38 of seed germination and survival was due to the disturbances caused at the physiological  
39 level coupled with chromosomal damage. Disturbance in the formation of enzymes involved  
40 in the germination process may be one of the physiological effects caused by mutagenic  
41 treatments particularly chemical mutagens [6].

42 Colchicine ( $C_{22}H_{25}NO_6$ ), originally extracted from *Colchicum autumnale*, may induce some  
43 morphological, cytological and histological changes, and even changes in the gene  
44 expression level [7]. Chemical mutagens such as ethyl methane sulfonate (EMS), a  
45 compound of the alkaline sulfonate series, is most frequently used for chemical mutagenesis  
46 in higher plants due to its potency and the ease with which it can be used [8]. It usually  
47 causes high frequency of gene mutations and low frequency of chromosome aberrations [9].  
48 The present investigation was planned and carried out to study the influence of some  
49 chemical substances i.e., (BA, colchicine, ethyl methane sulphonate) at different  
50 concentrations on some seed germination parameters, some vegetative growth  
51 measurements, as well as root till chromosomal behavior of papaya cultivar "Solo" through  
52 the cytological examination of papaya seedling.

53

## 54 **2. MATERIAL AND METHODS / EXPERIMENTAL DETAILS / METHODOLOGY** 55 **(ARIAL, BOLD, 11 FONT, LEFT ALIGNED, CAPS)**

56

57 The present investigation was carried out during two consecutive seasons 2015 and  
58 2016 in fruit nursery of faculty of Agriculture at Moshtohor, Benha University, in order to  
59 throw some spotlight on the impact of some chemical substances (Ethyl Methane  
60 Sulphonate – EMS; colchicine and benzyl adenine "(BA) on seed germination %, seed  
61 germination rate, some seedling growth measurements and cytological examination of root  
62 tip of *Carica papaya* cv. Solo.

63 In this regard, mature papaya fruits were collected from the trees which grown at  
64 fruit farm of Faculty of Agriculture, Moshtohor, Benha Univ., seed were extracted when the  
65 fruits have been ripened, and washed three times with tap water to get rid of fruit pulp  
66 residual. Finally, seeds were kept in shading place to be dried and stored in small coped  
67 glass contain calcium chloride to be ready for carrying out the investigation.

68 On the first week of March of both seasons, dried stored papaya seeds were soaked  
69 in tap water for 24h then taken out and placed in shade for 10 minutes to dry. those seeds  
70 were divided into ten groups. Each group represented by two hundred seeds and subjected  
71 to one of the following treatments:

72 1- Soaking in tap water for 12 hours (control).

73 2- Soaking for 12 hours in benzyl adenine (BA) at 1 %.

- 74 3- Soaking for 12 hours in benzyl adenine (BA) at 2 %.
- 75 4- Soaking for 12 hours in benzyl adenine (BA) at 3 %.
- 76 5- Soaking for 12 hours in colchicine at 1 %.
- 77 6- Soaking for 12 hours in colchicine at 2 %.
- 78 7- Soaking for 12 hours in colchicine at 3 %.
- 79 8- Soaking for 12 hours in ethyl methane sulphonate (EMS) at 10 ppm.
- 80 9- Soaking for 12 hours in ethyl methane sulphonate (EMS) at 20 ppm.
- 81 10- Soaking for 12 hours in ethyl methane sulphonate (EMS) at 30 ppm.

82 The dried seeds were soaked in aqueous solutions of the three investigation  
 83 chemical substances as well as control seed were soaked in tap water for 12 hours. Those  
 84 seeds were redried for 10 minutes in shade after soaking in the investigation chemical  
 85 substances and immediately sown on March 9th and 21st during 2015 and 2016 seasons,  
 86 respectively, in black polyethylene bags (30 cm in diameter) filled with a mixture of sandy  
 87 and clay soil (1:1 v/v) and kept under greenhouse conditions. The seeds were watered every  
 88 other day in the morning till the appearance of plumule. Furthermore, fungicide was applied  
 89 at the time of seed sowing as a tool protection against the fungal attack of *Rhizoctonia*  
 90 *solani* and *Fusarium* species, as well as weeds were completely removed along with their  
 91 roots as soon as they appear. The first appearance of plumule was recorded in the 1st week  
 92 of April during both seasons of study.

93 The abovementioned ten investigated treatments were arranged in complete  
 94 randomized design, where each treatment was replicated ten times (10 polyethylene bags)  
 95 and each replicate represented by an individual polyethylene bag which contains twenty  
 96 papaya seeds. Furthermore, the number of emerged seedlings was counted as soon as the  
 97 appearance of first true leaves on the 4th week at April of three days' intervals until seed  
 98 germination was completely ceased, then the following seed germination parameters were  
 99 calculated:

100

$$101 \quad \text{Total number of emerged seedling}$$

$$102 \quad 1\text{- Germination percentage} = \frac{\text{Total number of emerged seedling}}{\text{Total number of planted seeds}} \times 100$$

$$103 \quad \text{Total number of planted seeds}$$

104

105 2- Germination rate according to equation [10] :

106

107 
$$A_1 T_1 + A_2 R_2 + A_3 T_3 + \dots + A_n T_n$$
108 Germination rate = 
$$\frac{\text{A1 T1 + A2 R2 + A3 T3 + .....An Tn}}{\text{A1 + A2 + A3 ..... An}}$$

109

110

111 T1 = Number of days passed from soaking till first count 1.

112 T2 = Number of days passed from soaking till second count to Tn.

113 A1 = Number of germinated seeds at first count.

114 A2 = Number of germinated seeds at second count to An.

115 3- Number of days required for germination completion.

116 In order to study the impact of the three investigated chemical substances on some  
 117 seedling growth measurements and chromosomal behavior of sprouted papaya seedlings,  
 118 thin out of un-desirable seedlings (the weakest and the strongest ones) was done on the first  
 119 week of July, while the nearly uniform seedlings in their growth vigor were remained in the  
 120 polyethylene bags.

121 The treatments were arranged in complete randomized blocks design with nine  
 122 replicates (polyethylene bags), however, each replicate was represented by two papaya  
 123 seedlings. The seedlings were divided into three categories according to their growth vigor,  
 124 each category represented by three replicates for each treatment and subsequently each  
 125 category sampled by 30 seedlings for all studied treatments.

126 Seedling growth and chromosomal behavior as impacted by the three studied  
 127 chemical substances were evaluated on the 1st week of December through studding the  
 128 following parameters:

129 **A- Growth parameters: -**

130 1- Seedling height.

131 2- Stem diameter (cm).

132 3- root length

133 4- Number of leaves/seedling.

134 **B -cytological studies: -**

135 Papaya (*Carica papaya* L.) seedling roots were used for bioassay. Papaya seeds  
 136 were kindly supplemented from the research farm of Faculty of Agriculture, Moshtohor,

137 Benha University to be used in this study. Seeds were soaked in three different  
138 concentrations of Benzyl adenine, EMS and Colchicine. Root meristem raised in water were  
139 fixed in a fixative solution (3:1) and kept in alcohol 70 % in refrigerator until used for  
140 cytological examination.

141 About 100 cleaned papaya seeds were set up in petri dishes and soaked for 24  
142 hours here in tap water here in ,10 seeds were re-soaked in tab water and used as a control  
143 while the other 90 reads were picked out and divided into 3 groups, each one contain thirty  
144 reads and subjected to 1,2 and 3 of Benzyl adenine (BA), Ethyl methane sulphonate (EMS)  
145 and Colchicine for 12 h.

#### 146 **Mutagenic agents:**

147 Ethyl methane sulfonate (EMS): The linear formula of EMS is  $\text{CH}_3\text{SO}_3\text{C}_2\text{H}_5$ . This  
148 formula was referred to the free chemical database: (ChemSpider ID: 5887). Seeds before  
149 germination were subjected to the following concentrations; 1 %, 2 % and 3% for twelve  
150 hours. Benzyl adenine (BA) or 6-Benzylaminopurine (BA) is  $\text{C}_{12}\text{H}_{11}\text{N}_5$ . Cyclophosphamide  
151 (Colchicine) at the concentrations of 1 %, 2 % and 3%. The linear formula of colchicine is  
152  $\text{C}_7\text{H}_{15}\text{Cl}_2\text{N}_2\text{O}_2\text{P}$ .

#### 153 **Fixation and storage solutions:**

154 Root tips of the germinated Papaya seeds in the different investigated substances  
155 and ab water as control were excised and fixed in 1: 3 acidic alcohol consisted of a mixture  
156 of glacial acetic acid and ethanol respectively and later preserved in 70 % ethyl alcohol.

157 Staining agent (acetocarmine).

158 A carmine stain was prepared at the concentration of 1% by dissolving it in 45%  
159 acetic acid. Before adding the stain, root tips were put in a boiling acetocarmine for one  
160 minute for losing the tissue.

#### 161 **Root collection and slide preparation**

162 Papaya seeds were germinated at lab temperature using petri dishes filled with  
163 enough tap water to top. four to five weeks for root tips to grow. Seeds subjected to  
164 treatments were transferred to each concentration of BA, EMS and Colchicine after the

165 length of the roots reached to 1- 1.5 cm maximum. Roots were harvested at the morning.  
166 Root tips excised from treated and controlled materials were fixed in 1: 3 acidic alcohols and  
167 preserved in 70% ethyl alcohol. Root tips squashed were conducted using 1% Acetocarmine  
168 stain.

#### 169 **Mitotic index (MI) determination:**

170 The slides were viewed under the light microscope using 40 objective lens. On one  
171 slide for each treatment dividing cells (prophase, metaphase, anaphase and telophase) were  
172 counted to determine MI. MI was expressed as the number of dividing cells per 1000 cells  
173 scored.

174 Chromosomal aberrations were characterized and classified in the following types: large  
175 chromosomal deletion or losing a hole chromosome, sticky chromosomes, anaphase bridge  
176 chromosomes, lagging chromosomes, disrupted chromosome segregation, star cluster  
177 chromosomes, clumped chromosomes in metaphase. These aberrations were saved in  
178 photographic pictures.

#### 179 **Statistical analysis: -**

180 All the obtained data during each season of this study were subjected to statistical  
181 analysis of variance according to the method described by [11]. However, the differences  
182 means were differentiated by using Duncan's multiple range test [12].

183 - fourth level heading.]

184

### 185 **3. RESULTS AND DISCUSSION**

186

#### 187 **Effect of seeds pre-sowing soaking in different BA, colchicine and EMS solutions on** 188 **some germination measurements.**

189 In this regard some germination measurements germination percentage and  
190 germination rate of papaya Solo cv. in response to pre-sowing soak in some BA, colchicine  
191 and EMS solutions were investigated during 2015 and 2016 experimental seasons are  
192 presented in **Table (1)**.

#### 193 **- Seeds germination percentage:**

194 Data presented in **Table (1)**, indicate that the seeds germination percentage of  
195 papaya "Solo" cv. after 4 weeks from planting as influenced by their soaking for 12 hours in  
196 different BA, colchicine and EMS solutions significantly increased during both experimental  
197 seasons. However, pre-sowing soak in the highest BA concentration surpassed significantly  
198 than investigated treatments. On the other side, the least concentration of colchicine and

199 EMS solutions at (1 %) showed significantly the highest increase over control during two  
200 experimental seasons. In addition, other pre-sowing soak solutions (1 & 2 %) of BA ranked  
201 statistically the second one. Moreover, BA as a growth promoter explain the function for  
202 activating growth and germination particularly cell division.

203 - **Seeds germination rate:**

204 **Table (1)** reveal obviously that germination rate followed typically the same trend  
205 previously discussed with germination percentage. Herein, all BA, colchicine and EMS  
206 solutions resulted in a significant increase over the tap water soaked seeds (control) during  
207 both experimental seasons. The highest BA solution were statistically the superior, while  
208 their lowest concentration (1 & 2 %) ranked statistically second. In addition, tap water  
209 soaked seeds (control) was the inferior such trend was true during 2015 and 2016  
210 experimental seasons.

211 These results are in accordance with the findings of [13] reported that freshly extracted  
212 seeds of acid lime (*Citrus aurantifolia* swingle) were shade dried and were soaked in 15, 30,  
213 45 or 60 mM EMS solution for 12h caused decrease of percentage seed germination (36%)  
214 with increasing of EMS concentrations to 60 mM. Despite, seeds of *L. esculentum* cv. Roma,  
215 were treated with 0.1, 0.5 and 1% ethyl methane sulphonate (EMS) and exposed for 3 and  
216 6h, decrease in seed germination was observed with increasing EMS% [14]. Papaya seeds  
217 treated with colchicine at 0.5 or 1.0% and EMS at 200 ppm and 100 ppm improved  
218 germination parameters compared with untreated seeds (control) [15]. A clear effect of  
219 different EMS-treated on seeds germination percentage of *L. esculentum* (cv. Pusa – Early-  
220 Dwarf) showed that germination percentage increased with increasing EMS concentrations  
221 from 0.0150 to 0.1205%. Thereafter, decrease in germination percentage was observed at  
222 the highest concentration (0.2410%) [16]. Addition colchicine to cultured medium of *Solidago*  
223 *altissima* at 125 mg/l had an inhibition, while the other treatments (low concentration of  
224 colchicines) possessed the most promotion influences on survival capacity of explants (75-  
225 100 %) [17].

226 seeds of water melon without coat when seed nicking at radicle end with colchicine-  
227 treated showed high germination rates 84.3 and 77.1%, respectively [18]. The effect EMS  
228 and colchicine-treated seeds of Papaya at 0.1% and 0.5%, they found the stimulatory effects  
229 of low-dose colchicine treatment on seedling emergence and seed germination decreased  
230 with the increasing doses of colchicine [19]. [Reduced seed germination due to the effect of](#)  
231 [increasing doses of chemical mutagens on the meristematic tissues of the seeds may be](#)  
232 [causing damage of cell constituents at a molecular level or](#) to disturbance in the formation of  
233 enzymes involved in the germination process caused by EMS and colchicine. Impact of

234 mutagenic treatments i.e., EMS-treated seeds at 0.25- 0.30% of rice causing the reduction in  
 235 percentage of seed germination and survival was due to the chromosomal damage and  
 236 disturbance in the formation of enzymes involved in the germination process [20] and [6].

237

238 **Table (1): Impact of papaya seed soaking in different BA, colchicine and EMS solutions on**  
 239 **seed germination percentage and germination rate during 2015 & 2016 experimental**  
 240 **seasons.**

Parameters Treatments	Germination percentage %		Germination rate	
	First season	Second season	First season	Second season
1. control	55.67 g	54.33 h	3.68 i	3.42 i
2. BA at 1 %.	77.00 b	79.33 b	5.10 c	5.04 c
3. BA at 2 %.	80.67 a	81.00 b	5.32 b	5.24 b
4. BA at 3 %.	81.67 a	83.33 a	5.43 a	5.33 a
5. colchi at 1 %.	68.67 d	68.00 e	4.09 f	3.96 f
6. colchi at 2 %.	73.33 c	71.67 d	4.24 d	4.11 e
7. colchi at 3 %.	75.00 c	74.33 c	4.28 d	4.13 d
8. EMS at 10 ppm	61.00 f	63.67 g	3.75 h	3.57 h
9. EMS at 20 ppm	65.33 e	65.67 f	3.89 g	3.78 g
10. EMS at 30 ppm	65.67 e	67.33 ef	4.15 e	4.10 e

241 Means followed by the same letter/s within each column during every season are not significantly at 5 % level.

242

243 - **Impact of papaya seed soaking in different BA, colchicine and EMS solutions on some growth**  
 244 **measurements during 2015 & 2016 experimental seasons.**

245 In this concern average seedling height, stem diameter, root growth and average  
 246 number of leaves/seedling in response to various treatments were investigated during two  
 247 2015 and 2016 experimental seasons are presented in **Tables (2)**.

248 - **Average seedling height (cm):**



249           Concerning the response of average seedling height to the differential treatments, it  
250 is quite clear as shown in **Table (2)** , that all investigated treatments with various solutions  
251 from BA, colchicine and EMS. resulted in an increase in average seedling height of papaya  
252 "Solo" cv. translocated seedlings during both experimental seasons. Anyhow, the increase  
253 was more pronounced with (BA at 3 %) treated seeds, descendly followed by BA at 2 %, BA  
254 at 1 %, colchicine at 2 % and colchicine at 3 %. However, such increase was too few to  
255 reach level of significance either the investigated treatments were compared each other's or  
256 to tap water soaked seeds (control) only with few exceptions particularly with colchicine at 3  
257 % in the second season. Such trend of response was true during both 2015 and 2016  
258 experimental seasons.

259

#### 260 -Seedling diameter (cm):

261           Regarding the effect of different investigated treatments on stem diameter of papaya  
262 "Solo" cv. translocated seedlings **Table (2)** displays obviously that both (T3 & T4) treatments  
263 of BA 2 % and BA 3 % solutions induced significantly the thickest stem. Such trend was true  
264 during two seasons of study. Moreover, (T10 and T2) treatments of (EMS at 3 % and BA at 1  
265 %), respectively, ranked statistically second as their effect on stem diameter was concerned  
266 for papaya Solo cv. translocated seedlings during two experimental seasons. On the other  
267 side other investigated treatments increased significantly the average stem thickness during  
268 both seasons of study but T8 (EMS 1 %) showed statistically the least significant increase in  
269 stem diameter during 2015 and 2016 experimental seasons. In addition, other investigated  
270 treatments were statistically in between the aforesaid two extremes during two experimental  
271 seasons.

272           Moreover, BA as a growth promoter explain the function for activating growth  
273 specially stem diameter by increase cell division which gave more thickness for the stem.

#### 274 - Root length (cm):

275           This is The response of root length to various investigated treatments during both  
276 2015 and 2016 experimental seasons. and data obtained during both seasons for papaya  
277 Solo cv. translocated seedlings are presented in **Table (2)**. It is quite evident as shown from  
278 tabulated data that a noticeable grade of variance in trend of response could be observed  
279 between investigated treatments in this concern. Anyhow, the greatest length of root was  
280 significantly in closed relationship to BA at 3 % during two seasons of study. Moreover, BA  
281 at 2 % came statistically second. On the contrary, the least significant increase in root length  
282 was always in concomitant to EMS at 3 % and colchicine at 3 % during 2015 and 2016

283 experimental seasons of study. In addition, other treatments were statistically in between the  
284 aforesaid two extremes. Such trend was true during both seasons.

285 Moreover, the trend of response of root length of translocated seedlings may be  
286 attributed to the variance in biological and physiological roles could be played by BA  
287 pertaining shoot growth and root length and development.

#### 288 - Number of leaves/seedling:

289 With regard to the response of leaves number of per seedling an individual seedling  
290 to the differential investigated treatments, obtained data are presented in **Table (2)**. It is  
291 quite evident that the greatest leaves number of per seedling was significant in closed  
292 relationship to such seedling was subjected to BA at 3 % during 2015 and 2016  
293 experimental seasons. Moreover, BA at 2 % ranked statistically second. Anyhow, pre-sowing  
294 soaked in BA at 1 % solution ranked statistically 3<sup>rd</sup>, descendingly followed by soaking in  
295 EMS 1 %, EMS 2 % and EMS 3 % during both 2015 and 2016 experimental seasons. On  
296 the contrary, the least significant leaves number per seedling that exhibited by three  
297 investigated treatments (colchicine at 3 %, control and colchicine at 2 %), respectively. Such  
298 trend was true during 2015 and 2016 experimental seasons. Treated seeds of two pea  
299 cultivars with EMS concentrations of 0.5, 0.75 and 1.0 %. In M1-generation, number of  
300 branches decreased with EMS at 0.75 and 1.0 % [21].

301 The cytokines promote shoot development through increased cell division, regulation of the  
302 cell cycle and the number of cycles that cells in the meristems [22]. Adding, 20 mg/l  
303 colchicine into the medium for one week inducing tetraploidy plants. Morphological  
304 observations showed that the stems and the leaves of tetraploid plants were thicker and  
305 larger than in diploid ones [23] (1999). Also, BA treatment at 10 ppm increased growth  
306 characters i.e., plant height, total root length fresh and dry weights of shoots and roots of  
307 maize plants [24]. Foliar spray of soybean plants with benzyl adenine at 75 ppm  
308 significantly increased plant height, leaves number and branches per plant and dry matter of  
309 plant [25]. The effect beneficial of foliar application of soybean plants with benzyl adenine at  
310 50 ppm significantly increased stem length, diameter, leaf area surface, branches number,  
311 leaves number per plant and fresh and dry weights of plant [26]. Similarity, the foliar  
312 application of pelargonium (Geranium) plants with BA at 20 and 40 mg/L significantly  
313 increased plant height and number of branches/plant finding by [27]. Egyptian lupine plants  
314 exposed to salt stress, observed that foliar application of benzyl adenine (BA) (1 & 100 ppm)  
315 has stimulating effect on all growth characters, i.e., plant height and number of  
316 branches/plant grown under normal and saline conditions [28]. In *Nigella sativa* plants  
317 which benzyl adenine (5 & 25 ppm) treatments as seed soaking increased root length and

318 diameter, plant height stem diameter, number of leaves, total leaf area/plant and net  
 319 assimilation rate [29]. Foliar spray of snap bean plants with benzyl adenine (BA) at 20 & 40  
 320 ppm and putrescine (Put) at 200 ppm significantly increased plant height, leaves number  
 321 /plant and branches and fresh and dry weights of shoots [30]. The increased values of  
 322 vegetative parameters due to the lower dose of colchicine might be due to enhanced the  
 323 action of auxin (indole-3-acetic acid) and the cells divided more actively in *Helianthus*  
 324 *tuberosus* [31]. Higher doses of colchicine led to increased leaf size and number of leaves  
 325 per plant in colchicine-treated plants over control in *Gossypium arboreum* L [32] . EMS-  
 326 treated plants was also reported in papaya increased cell division as well as activation of  
 327 growth hormones such as auxin [33]. The effect of colchicine-treated seeds of *Phlox*  
 328 *drummondii* increasing seed germination and morphological characteristics at low  
 329 concentrations [34]. The effect of EMS-treatments on induced micro mutations and obtained  
 330 on dwarf plant types. The minimum plant height in dwarf mutant was below 90 cm. The  
 331 maximum frequency of dwarf mutants was observed in 30kr + 0.1% EMS followed by 40kr +  
 332 0.25% EMS treatment. The tallest mutant (155cm) was observed in 0.25 % EMS treatment  
 333 followed by a mutant with 131 cm in 30kr+0.25% EMS while the parent of rice Akshaya cv.  
 334 possess 100-110cm height [35].

335 **Table (2): Impact of papaya seed soaking in different BA, colchicine and EMS solutions on**  
 336 **some growth measurements during 2015 & 2016 experimental seasons.**

Parameter s Treatment s	No. leaves /seedling		Seedling height (cm)		Seedling diameter (cm)		Root length (cm)	
	First seaso n	Secon d seaso n	First seaso n	Second season	First seaso n	Secon d seaso n	First seaso n	Secon d seaso n
1.control	9.33 f	7.67 f	52.33e	58.67f	2.53e	2.45de	14.73d	14.85d
2. BA at 1 %.	14.00 c	11.67 cd	99.00a	101.00c d	2.77d	2.83c	18.53c	18.63c
3. BA at 2 %.	15.33 b	13.00 b	97.00a b	103.00b c	3.13b	3.20b	21.38b	21.40b
4. BA at 3 %.	17.67 a	16.33 a	96.83a b	100.00d	3.37a	3.40a	23.80a	23.87a
5. colchi at 1 %.	10.33 e	13.00 b	75.00c	101.33c d	2.93c	3.13b	13.63e f	13.50f
6. colchi	10.67	8.00 ef	95.07b	105.00b	2.65de	2.62d	13.32f	13.30f

at 2 %.	e						g	
7. colchi at 3 %.	7.67 g	8.67 e	97.00a b	113.33a	2.37f	2.27e	13.02g	13.07f
8. EMS at 10 ppm	12.67 d	11.00 d	70.00d	78.67e	2.50ef	2.45de	13.45f g	13.50f
9. EMS at 20 ppm	12.67 d	12.00 c	76.33c	80.00e	2.65de	2.57d	13.93e	13.92e
10. EMS at 30 ppm	12.67 d	13.33 b	69.00d	80.67e	2.97c	2.87c	14.10e	14.23e

337 Means followed by the same letters within each column during every season are not significantly at 5 % level.

338


339

340

341

342

343




344

345

346

347

**Photo (1) Impact of papaya seed soaking in different BA, solutions on vegetative growth during 2015 & 2016 experimental seasons.**  
**Con= control, BA1= BA solution at 1% , BA2=BA solution at 2% and BA3=BA solution at 3%.**



348

349

350 **Photo (2) Impact of papaya seed soaking in different colchicine solutions on vegetative growth**  
351 **during 2015 & 2016 experimental seasons.**

352 **C1= control , C2= Colchicine solution at 1% , C3= Colchicine solution at 2% and C4=**

353

**Colchicine solution at 3%.**

354



355

356 **Photo (3) Impact of papaya seed soaking in different EMS solutions on vegetative growth during**  
 357 **2015 & 2016 experimental seasons.**

358 **Con= control, E1= EMS solution at 10ppm, E2= EMS solution at 20ppm and E3= EMS**  
 359 **solution at 30ppm.**

360 **Mitotic Index:**

361 Means of mitotic index (MI %) resulted by BA, EMS and Colchicine are shown in  
 362 Table 3. The means of mitotic index at three levels of Colchicine were close to each other  
 363 and the same trend was also obtained by EMS. These results appeared that the differences  
 364 between different levels of each agent were insignificant.

365 The means of dividing cells treated with Colchicine were significantly higher than of  
 366 BA and EMS. This indicated that Colchicine did not interfere with mitosis and did not prevent  
 367 cell division if compared with of BA and EMS which decreased the mitotic index and  
 368 interfered with mitosis to greater extent.

369 Therefore, it can be concluded that EMS was more inhibitor of cell division followed  
 370 by BA than Colchicine. This may be due to more damage resulted by BA and EMS affected  
 371 on DNA replication during mitosis.

372 **The figure shows the different chromosomal aberration as follows:**

373 Sticky chromosomes at metaphase, Laggards and lagging chromosomes and polyploidy are  
 374 the main chromosomal aberrations or abnormalities during the cell division of papaya after  
 375 treatment with the three mutagens. with different ratio and different appearance.

376 Colchicine and EMS showed disrupted type of chromosomal aberrations which  
377 appeared during metaphase stage. It appeared that disrupted metaphase varied from  
378 Colchicine to EMS. In addition, EMS caused disrupted chromosomes in metaphase followed  
379 by anaphase which did not occur with Benzyl adenine.

380 Both Colchicine and EMS caused abnormal mitosis which appeared as sticky  
381 chromosomes. Colchicine caused sticky chromosomes in during metaphase and telophase.  
382 Similarly, EMS showed sticky with polyploidy chromosomes during metaphase, anaphase  
383 and telophase. These results indicated that colchicine had strongest effect on chromosomal  
384 behavior during mitosis and exerted more chromosomal damage. Indeed, sticky  
385 chromosomes would cause the death of those cells. Similar results were obtained by authors  
386 among them.

387 A chromatid bridge would occur as a result of the weakness of the spindle fiber.  
388 Bridge as an aberration occurs due to treatment by both EMS and Colchicine.

389 During abnormal chromosomal behavior of mitosis, spindle fiber can not to attract  
390 one chromosome, this chromosome remains near the middle of the cells. This phenomenon  
391 called lagging chromosome and resulted genome aneuploidy  $2n-1$ . This kind of aberration  
392 did not occur by Among the chromosomal aberrations caused by Colchicine or EMS. the  
393 formation of star type of chromosomes was shown. Both Colchicine and EMS caused this  
394 type of aberration.

395 In conclusion, the treatments by colchicine and EMS caused different types of  
396 chromosomal aberrations with variable percentages than the normal cells in control  
397 experiment the same time there were differences of the percentage ratio of each. This  
398 indicated that both chemical agents are dangerous. Although, EMS was more dangerous  
399 than Colchicine because of cytotoxicity delaying mitosis and inducing mass chromosomal  
400 aberrations.

401  
402 Sex determination in papaya (*C. papaya* L.) is due to a single gene with three allelic  
403 forms: *m*, *M1* and *M2*. The *mm*, *M1m*, and *M2m* genotypes represent gynoecious, and  
404 roecious and hermaphrodite individuals, respectively. The *M1M1*, *M2M2* and *M1M2*  
405 genotypes are not found due to the zygotc lethality. The *m* homologous region is normal and  
406 the viable genotypes are *M1m* (male plant), *M2m* (hermaphrodite plant) and *mm* (female  
407 plant). A large concentration of genes for femaleness is in the sex chromosomes but genes  
408 for maleness are in the autosomes. Therefore, the *mm* genotype is distillated and its  
409 homozygote condition confers phenotypic stability [36] and [37] . Small doses of colchicine  
410 enhanced the action of auxin (indole-3-acetic acid) because the cells divided more actively;

411 instead, at higher doses, colchicine led to C-mitoses and inhibited cell Multiplication In  
412 *Helianthus tuberosus* [38].

413

414 The karyotype of *Carica papaya* L. consisted of eight medians (metacentric) four  
415 submedian, four sub terminal and two terminal-centromeric chromosomes, formed that the  
416 arm ratio value of eight median centromeric chromosomes range from 1.0 to 1.3 while the  
417 arm ratio value of four submedian centromeric chromosomes were very close to 3.1 the  
418 lowest extreme of the arm ratio range of the sub terminal centromeric chromosome [15]. The  
419 cells with a larger complement of chromosomes grow larger to maintain a constant ratio of  
420 cytoplasmic to nuclear volume, and express more proteins with the presence of more genes.  
421 This increase in size may translate to an increase in the plant and its organs [32]. Also,  
422 using several BAC clones that were explaining mapped to th papaya X/Y chromosomes,  
423 found that the presumed sex chromosomes of *J. spinosa* are homomorphic and pair  
424 completely. In other species, chromosomes had been counted with traditional means, and all  
425 were reported to have a diploid number of  $2n = 18$ . The remaining three genera have never  
426 been studied, yet are disproportionally important because, respectively, they represent the  
427 deepest divergence in the Caricaceae ( *Cylicomorpha* ) and the sister clade to *Carica* [39].  
428 Gamma radiation, EMS, and their combinations are potent mutagens, well known for their  
429 action causing point mutations, enzyme inhibitions and chromosomal aberrations [40]. Sister  
430 to all New World Caricaceae is an African genus ( *Cylicomorpha* ) with two species. A draft of  
431 the papaya genome became available in 2008, and since then, considerable effort has gone  
432 into understanding the sex chromosomes of *C. papaya* [41]. All Caricaceae species are  
433 classified as diploids ( $2n=2x=18$  chromosomes) and dioecious, except for *C. papaya*, *V.*  
434 *monoica* e *V. cundinarmacensis*. The plant sexual determination in papaya is due to one  
435 gene with three alleles. It was not observed sexual chromosome in their study. Thus, if there  
436 are sexual chromosomes in *C. papaya*, they are probably homomorphic [42].

437

438







439

440

**photo (4): Normal metaphase without any treatment in the mitotic cell of papaya.**






441


442 **photo (6): The effect of Benzyl adenine with three different concentrations on the mitotic**  
 443 **cells of papaya.**

444 photo (6): The effect of Benzyl adenine with three different concentrations on the  
 445 mitotic cells of papaya. photo 6-A and B anaphase with irregular distribution of  
 446 chromosomes between the two poles. photo 6-C three star groups of scattering of  
 447 chromosomes in a dividing cell of a root tip at the beginning of telophase. photo 6-D one  
 448 fragment at the equator of the metaphase. photo 6-E irregular distribution of chromosomes  
 449 at the metaphase. photo 6-F Two laggards at metaphase.

450

451







452 **photo (7): The effect of EMS with three different concentrations on the mitotic cells of papaya.**

453 photo (7): The effect of EMS with three different concentrations on the mitotic cells  
 454 of papaya. photo 7-A and B metaphase with one lagging chromosome. photo 7-C Scattering  
 455 of chromosomes in a dividing cell of a root tip at metaphase. photo 7-D one lagging  
 456 chromosome at metaphase. photo 7-E irregular distribution of chromosomes at the  
 457 beginning of anaphase. photo 7-F clear polyploidy in metaphase with tetraploid number of  
 458 chromosomes and C-metaphase.

459  
 460  
 461





462

463 **photo (8): The effect of Colchicine at three different concentrations on the mitotic cells of papaya.**

464 photo (8): The effect of Colchicine with three different concentrations on the mitotic  
 465 cells of papaya. photo 8-A metaphase with one lagging chromosome. photo 8-B metaphase  
 466 with two lagging chromosomes. photo 8-C Unequal distribution of chromosomes in  
 467 anaphase with polyploidy. photo 8-D sticky chromosomes at metaphase. photo 8-E  
 468 metaphase with tetraploid number of chromosomes. photo 8-F scattering of chromosomes in  
 469 a dividing cell of a root-tip exposed to 3% of colchicine.

470 **Table (3): Type and percentage of mitotic abnormalities in the root tips of**  
 471 **papaya exposed to the Benzyl adenine, Ethyl methane sulphonate and colchicine with**  
 472 **three different concentrations.**

Conc. Ppm of mutagen	Total cells scor s	No.of Divid . cells	MI %	Number of cells in the different phases of the cell cycle				
				Interpha se	proph ase	Metapha se	Anaph ase	Telopha se.
Control	500	92	18.4 %	15.9 %	2.20 %	0.12	0.5	0.13
BA 1%	500	47	9.40 %	8.02 %	1.09 %	0,10	0.04	0.15
BA 2%	500	32	6.40 %	5.10 %	0.98 %	0.09	0.11	0.12
BA 3%	500	18	3.60 %	2.11 %	1.20 %	0.07	0.02	0.20
Control	500	87	17.4 %	14.8 %	2.00 %	0.22	0.08	0.0.8
EMS 1%	500	40	8.00 %	6.01 %	1.35 %	0.16	0.04	0.28
EMS 2%	500	24	4.80 %	3.00 %	0.80 %	0.25	0.49	0.26
EMS 3%	500	20	4.00 %	2.90 %	1.10 %	0.00	0.00	0.00
Control	500	97	19.4 %	17.95 %	1.06 %	0.20	0.09	0.11
Colchicine1 %	500	81	16.2 %	15.05 %	0.85 %	0.40	0.05	0.30

<b>Colchicine2</b> %	<b>500</b>	<b>69</b>	<b>13.8 %</b>	<b>12.00 %</b>	<b>1.12 %</b>	<b>0.23</b>	<b>0.30</b>	<b>0.15</b>
<b>Colchicine3</b> %	<b>500</b>	<b>53</b>	<b>10.6 %</b>	<b>9.00 %</b>	<b>0.95 %</b>	<b>0.16</b>	<b>0.30</b>	<b>0.19</b>

473

474

475

476

#### 4. CONCLUSION

477

478 [It can be recommended from the results of this study that both BA 2 % and BA 3 % increased  
479 significantly germination %, germination rate and growth measurements. Moreover, EMS  
480 was more inhibitor of cell division followed by BA than Colchicine. This may be due to more  
481 damage resulted by BA and EMS affected on DNA replication during mitosis. option]

482

483

484

485

#### COMPETING INTERESTS

486

486 Authors have declared that no competing interests exist.

487

488

489

#### REFERENCES

490

491 [1] Halmann, M. (1990). Synthetic plant growth regulators. *Advances in Agronomy* 43:47-  
492 105.

493 [2] Khryanin, V.N. 2002. Role of phytohormones in sex differentiation in plants. *Russian*  
494 *Journal of Plant Physiology* 49 (4):545-551.

495 [3] Davies, P.J. (2004a). The plant hormones : Their nature, occurrence and function. In  
496 *Plant hormones biosynthesis, signal transduction, action!*, ed. P. J. Davies, 750. Dordrecht ;  
497 Norwell, MA: Kluwer Academic Publishers.

498 [4] Nishijima, T., M. Hideari, K. Sasaki, and T. Okazawa. 2006. Cultivar and anatomical  
499 analysis of corolla enlargement of petunia (*Petunia hybrida* Vilm.) by cytokinin application.  
500 *Scientia Horticulturae* 111:49-55.

501

502 [5] Menendez, V., M.A. Revilla, and H. Fernandez. (2006). Growth and gender in the  
503 gametophyte of *Blechnum spicant* L. *Plant Cell Tissue and Organ Culture* 86 (1):47-53.

504 [6] Kulkarni GB. (2011). Effect of mutagen on pollen fertility and other parameters in  
505 horsegram (*Macrotyloma uniflorum* (Lam.) Verdc). *Bio. Sci. discovery*. 2 (1): 146-150.

506 [7] Murali K.M., Jeevanandam V., Shuye J. and Srinivasan R. (2013). Impact of colchicine  
507 treatment on *Sorghum bicolor* BTx 623, *Mol. Plant Breed*. 4(15) 128–135.

508 [8] Wattoo J.I., Aslam K., Shah S.M., Shabir G., Sabar M., Naveed S.A., Waheed R.,  
509 Samiullah, Muqaddasi Q.H.,

510 [9] Mohamed Z., Ho W.S., Pang, S.L., Ahmad F.B. (2014). EMS-induced mutagenesis and  
511 DNA polymorphism assessment through ISSR markers in *Neolamarckia cadamba*  
512 (kelampayan) and *Leucaena leucocephala* (petai belalang), *Eur. J. Exp. Biol*. 4(4):156–163.

513 [10] Chacko, E.K. and R.N. Singh. (1966). The effect of gibberellic acid on the germination of  
514 papaya seeds and subsequent seedling growth. *Trop. Agr*. 43:341–346.

515 [11] Snedecor, G.W. and Cochran, W.G. (1989). *Statistical methods*, pp 177-195. 8th  
516 edition. Iowa state university press.

517

- 518 [12]Duncan, D.B. (1955): Multiple range and multiple F tests. *Biometrics*, II: 1-42.
- 519 [13] Jawaharlal , M.; Sambandamoorthy, S. and Irulappan, L. (1991). Effect of gamma ray  
520 and EMS on seed germination and seedling growth in acid lime (*Citrus aurantifoliaswingle*).  
521 *South Indian Horticulture* .39 (6). 332 – 336.
- 522 [14] Nusrat, S. and Mirza, B. (2002). Ethyl methane sulfonate induced genetic variability in  
523 *Lycopersiconesculentum*. *International J. of Agric. and biology*. 4:1.89-92.
- 524 [15] Bakry, KH. A. and Ismaeil, F. H. (2002): Pre-sowing treatments of papaya seeds as  
525 influenced by some chemicals and irradiation on germination, growth, flowering, sex  
526 expression and fruit quality. 2ndIntre. Conf. Hort. Sci., 10- 12 sept., Kafr El-sheikh, Tanta  
527 Univ. Egypt.
- 528 [16] Padma, K. and Chauhan, P. (2005). Effect of EMS on germination, plant height and  
529 fruiting of *Lycopersiconesculentum*. *Flora and fauna ( Jhansi)*, 11. 39-41.
- 530 [17] Sayed, S. Sawsan; Yousef, Hanan, M. A. and Yousef, E.M.A. (2007). Influence of  
531 colchicines and sodium azide treatments on micropropagability and biochemical constituents  
532 of *Solidago altissima* Gray var "Tara" explants in vitro. *J. Biol. Chem. Environ. Sci. Vol. 2 (2):*  
533 257-276.
- 534 [18] Jaskani, M.J.; Kwon, S. W.; Kim, E. and Bokrae, K. (2004): Polypoidy affects fruit  
535 characteristics, seed morphology and germination in watermelon (*Citrulluslanatus*). *J. of the*  
536 *Korean society for Horti. Science*. 45 (5) 233 – 237.
- 537 [19] Pranath, B., Rekha, A. and Pandey, A.K. (2015). Effect of pre-sowing treatments with  
538 chemical mutagens on seed germination and growth performance of jamun (*Syzygium*  
539 *cumini* L. Skeels) under different potting substrates. *Fruits*. vol. 70(4): 239-248.
- 540 [20] Chakraborty N. R. and Kole P.C. 2009.Gamma ray induced morphological mutations in  
541 non-basmati aromatic rice. *Oryza* 46 (3): 181-187.
- 542
- 543 [21]El-Kobisy, O. S. A. (1988). Ethylmethane sulphonate morphological .Ms.D. Thesis of  
544 Agric.cairo Univ .
- 545 [22] Arigita,.L.; Fernandez, B.; Gonzalez, A. and Sanchez Tames, R. (2005): Effect of the  
546 application of benzyladenine pulse on organogenesis, acclimatisation and endogenous  
547 phytohormone content in kiwi explants cultured under autotrophic conditions.*Plant*  
548 *Physiology and Biochemistry*, 43:161-167.
- 549 [23]Wang-Honghe; X.U. Gexin; Q. (1999). In vitro induction of applied plants in colchicine-  
550 treated *Sinningia speciosa*. *J. of Tropical and Subtropical Botany*, 7: 237-242.
- 551 [24] Shadi, A. I.; Sarwat, M. I.; El-Din, M. A. T. and Abou Deif, M, H. (2001): Effect of  
552 benzyladenine treatment on chemical composition and salt tolerance of some maize in  
553 breeds under salt a stress. *J. of Agric., Sci.*, 9(1):95-108.
- 554 [25] atil, R. R.; Deotale, R. D.; Hatmode, C. N. and Band, P. E.; Basole, V. D. and  
555 Khobragade,T. R. (2002 b).Effect of 6-benzyladenine on biochemical and yield contributing  
556 parameters and yield of soybean. *India. J. of Soils and Crops*, 12(2): 270-273.
- 557 [26] Gad, M. S. H. (2005): Physiological studies on the effect of some growth regulators on  
558 soybean plant. M.Sc. Thesis, Fac. of Agric., of Moshtohor; Zagazig Univ.
- 559 [27] Youssef, A. A. (2004). Influence of foliar spray with brassinosteroid and benzyladenine  
560 on the growth, yield and chemical composition of {*Pelargonium graueolens* L} plants. *Annals*  
561 *Agric. Sci, Ain Shams Univ Cairo*.49(1)313-326.
- 562 [28]Medani, R. A. (2006): Effect of salinity, benzyladenine and their interaction on botanical  
563 characters and chemical constituents of Egyptian lupine plant (*Lupinus termis*, L.) *Annals of*  
564 *Agric. Sci.*, Moshtohor, 44(4): 1609-1628.
- 565 [29] Abd El-Gawad, H. A. (2006): Growth performance of black cumun (*Nigella sativa* L.)  
566 plants using certain growth conditions. Ph.D. Thesis, Fac. of Agric., Moshtohor Benha Univ.
- 567 [30] Abo El-Saoud, M. S. (2005): Physiological studies on the role of some bioregulators in  
568 growth, flowering and yield of snap bean. Ph.D. Thesis, Fac. of Agric., Moshtohor, Benha  
569 Univ.

- 570 [31] Bennici A., Silvia S. and Bruno M. (2006). Morphogenic effect of colchicine in *Cichorium*  
571 *intybus* L. Root explants cultured in vitro, *Caryologia* 59(3) 284–290.  
572
- 573 [32] Raufe S., Khan I.A. and Khan F.A., (2006). Colchicine-induced tetraploidy and changes  
574 in allele frequencies in colchicine-treated populations of diploids assessed with RAPD  
575 markers in *Gossypium arboreum* L, *Turk. J. Biol.*( 30) 93–100.  
576 [33] Singh S.V., Singh D.B., Yadav M., Roshan R.K. and Pebam N.(2010). Effect of EMS on  
577 germination, growth and sensitivity of papaya (*Carica papaya* L.) cv. Farm Selection-1, *Acta*  
578 *Hort.* 851 113–116.  
579 [34] Tiwari A.K. and Mishra S.K. (2012). Effect of colchicine on mitotic polyploidization and  
580 morphological characteristics of *Phlox drummondii*, *Afr. J. Biotechnol.* 11(39) (2012) 9336–  
581 9342.  
582
- 583 [35] BOLBHAT SADASHIV N. BHOGE VIKRAM D. AND DHUMAL KONDDIRAM N. (2012).  
584 Effect of mutagens on seed germination, plant survival and quantitative characters of  
585 horsegram (*Macrotyloma uniflorum* (Lam.) Verdc). *Research Article. Vol 2(4). Oct-Dec.* 129-  
586 136.  
587 [36] Hofmeyr J. D. J. (1938). Genetical studies of *Carica papaya* L. the inheritance and  
588 relation of sex and certain plant characteristics. *South African Department of Agri. and*  
589 *Science Bulletin*, n. 187:123-155.  
590 [37] Storey W. B., (1953). Genetics of the papaya. *Journal of Heredity*, 44 (2): 70-78.  
591 [38] Martin G., (1945). Action de la colchicine sur les tissus de topinambour cultivé in vitro.  
592 *Rev. Cytol. Cytophysiol. Veg.* 8. 1–34. Mooney, P.A. and Van Staden, J. 1986. *J. Plant*  
593 *physiology.* 123, 1-21.  
594 [39] Alexander Rockinger 2 , Aretuza Sousa , Fernanda A. Carvalho , and Susanne S.  
595 Renner 2016. Chromosome number reduction in the sister clade of *Carica papaya* with  
596 concomitant genome size doubling. *AMERICAN J. OF BOTANY.* 103 (6):1082 – 1088.  
597
- 598 [40] Auti SG. 2005. Mutational Studies in mung (*Vigna radiata* (L.) Wilczek). Ph.D.  
599 Thesis. University of Pune, Pune (MS), India.  
600 [41] Ming , R. , S. Hou , Y. Feng , Q. Yu , A. Dionne-Laporte , et al. 2008 . The draft genome  
601 of the transgenic tropical fruit tree papaya ( *Carica papaya* Linnaeus). *Nature* 452 : 991 –  
602 996 .  
603 [42] Corrêa, D.J.P., Fabiane, R.C., Telma, N.S., Pereira, M.F.N. and Messias, G.P.(2009).  
604 Karyotype determination in three *Caricaceae* species emphasizing the cultivated form (*C.*  
605 *papaya* L.). *CARYOLOGIA* Vol. 62, no. 1: 10-15.  
606  
607  
608  
609  
610  
611

612

**APPENDIX**