

Hypoglycaemic and Hypolipidemic Effects of *Treculia Africana* aqueous Leaves Extract in Alloxan Induced Diabetic Rats

Abstract

This study investigated the hypoglycaemic and hypolipidaemic effects of *Treculia Africana* plant used in Nigeria as medicinal plant. Diabetes mellitus was induced by a single dose intraperitoneal injection of alloxan 150mg/kg body weight. Twenty five (25) male albino rats were divided into five groups, five (5) rats per group; normal control, diabetic control and diabetic groups treated with aqueous leaves extract of 200,400 and 800mg/Kg body weight respectively for 21 days orally. The effects of the extract on some biochemical parameters were evaluated; fasting blood glucose level was assayed using glucose oxidase method, total cholesterol and HDL –cholesterol were assayed using enzymatic method while LDL-cholesterol was determined by Friedewald equation. The results showed that, extract significantly ($p<0.05$) decrease the elevated fasting blood glucose levels, total cholesterol, triglyceride and LDL- cholesterol when compared with the diabetic control rats. The extract also caused significant ($p<0.05$) increased in HDL –cholesterol and body weight when compared with diabetic control rats. Aqueous leave extract of *Treculia Africana* possess hypoglycemic effect and the most effective dose was 800mg/Kg body weight in amelioration of hyperglycaemia and most all toxicity effects of alloxan on lipid profile.

Key words: Hypoglycaemic, Hypolipidaemic, *Treculia Africana* plant

1.0 Introduction

Diabetes mellitus (DM) is a metabolic disease characterized by a persistent elevation of fasting blood glucose level (FBGL) above 200 mg/dl, due to insufficient or complete cessation of insulin synthesis or secretion and/or peripheral resistance to insulin action (Ortiz-Andrade *et al.*, 2005). It is characterized by persistent hyperglycemia and disturbances in the metabolism of fuel compounds (Saidu *et al.*, 2010). It occurs when pancreas cannot produce insulin (type 1 DM) or cannot produce enough of insulin or the body cannot effectively use the produced insulin for effective uptake of glucose to the target cells (type II DM) (Hajera *et al.*, 2011). Lipid abnormalities, anemia, alteration of liver and kidney functional indices has been implemented as major risk factors to the progression of both microvascular and

32 macrovascular diabetes complications (Canargo and Gross, 2014). Globally, as of 2013, an
33 estimated 382 million people have diabetes worldwide, with type 2 diabetes making up about
34 90% of the cases. This is equal to 3.3% of the population, with equal rates in both women and
35 men (Vos, *et al.*, 2012). Ethnomedicinal plants might provide new oral hypoglycemic
36 compounds, which can counter the high cost of the current hypoglycemic medicines for many
37 rural populations in developing countries (Noor *et al.*, 2008).

38 African breadfruit is a large, slow-growing, evergreen tree with a dense, spreading crown;
39 usually growing 15 - 30 metres tall but with some specimens up to 50 metres (Ruffo *et al.*,
40 2001). *Treculia africana* (African breadfruit) is a genus of the evergreen flowering plant and
41 belongs to the family of moraceae. There are three (3) varieties of *Treculia africana*: *Treculia*
42 *africana* var. *africana*, *Treculia africana* var. *inverssa* and *Treculia african mollis* ((Orwa,
43 Mutua, Kindt, R., & Anthony, 2009)).It is a common forest tree called by various names
44 among different tribes in Nigeria, such as “Ukwa” (Igbo), “afon” (Yoruba), “barafuta”
45 (Hausa), “Ize” (Benin), “eyo” (Igala) and “edikang” (Efik) (Olapade & Umeonuorah, 2014).
46 *Treculia africana* are highly nutritious and constitute a cheap source of vitamins, minerals,
47 proteins, carbohydrate and fats (Ojokoh, Fayemi, & Ocloo, 2014). It also contained
48 appreciable amounts of flavonoid, polyphenols, anthraquinones, saponins and cardiac
49 glycosides. These secondary metabolites are known to have potential ethno-medicinal effects
50 as well as other physiological activity (Osabor V. N., Ogar, Okafor, & Egbung, 2009)
51 (Osabor *et al.*, 2009). Survey among herbalist and number of patients attending Diabetic
52 clinics in the University College hospital Ibadan revealed the ethno-medicinal hypoglycemic
53 effect of *Treucia Africana* (Oyelola *et al.*, 2007). In Ghana, the root decoction is used as an
54 anthelmintic and febrifuge. Ethnomedically, it is used as a verbrifuge, vermifuge,
55 galactagogue and laxative (Ogbonnia *et al.*,2008). The stem and bark of *Treculia africana* are
56 employed in the treatment of helminthiasis, malaria, leprosy and rheumatism (Sherifat,
57 Olusegon, & Oluwole, 2007). It also used for the treatment of inflammation, diabetes
58 mellitus, diarrhea, and tapeworm infection (Chukwuemeka, *et al.*, 2012). Herbal medicine is
59 in used by about 60% of the world population both in the developing and in the developed
60 countries where modern medicines are predominantly used (Richert *et al.*, 1999). Ethno-
61 medicinal anti-diabetic plants might provide new oral hypoglycemic compounds, which can
62 counter the high cost and poor availability of the current medicines for many rural
63 populations in developing countries (Noor *et al.*, 2008)

64 **2.0. METHODOLOGY**

65 **2.1. COLLECTION AND IDENTIFICATION OF PLANT MATERIALS**

66 *Treculia Africana* leaves was obtained from Umuidi community of Anambara State of
67 Nigeria. The sample was identified and authenticated by the Department of Science
68 Laboratory Technology, Federal Polytechnic Kaura Namoda, Zamfara State. Nigeria

69 **2.2. PREPARATION OF SAMPLES**

70 The samples were thoroughly washed to remove sand and the drained parts were air dried
71 later. The samples were grounded using wooden mortar and pestle until powder was obtained
72 to ensure homogeneity. 100g of the plant powder was soaked in 1000ml of boiled distilled
73 water and agitated intermittently for 24 hours. The solution was then filtered using filter
74 paper to obtained the aqueous extract which was then be allowed to dried in an oven dryer at
75 100°C to obtained the crude extract (Luka, 2013). The extract was stored in an air tight
76 container for further work, the required doses of 200mg, 400mg and 800mg/kg body weight
77 was obtained by reconstituting the stored extract using distilled water and administered
78 (Meraiyebu, 2014).

79 **2.3 Experimental Animals**

80 Twenty five (25) apparently healthy young male Wister albino rats weighing between 100 –
81 200 g were used for this study. The rats were kept at animals' house under normal
82 environmental conditions and maintained with free access to pelletized growers feed, and
83 access to water *ad libitum*. The animals were allowed to acclimatize for two weeks (14 days).
84 All animal procedures were performed according to the Guide for the Care and Use of
85 Laboratory Animals of the National Institute of Health as well as the guidelines of the
86 Animal Welfare Act. Experiments were carried out with prior permission from the
87 Institutional animal **Ethical committee**, Federal Polytechnic Kaura Namoda, Zamfara State,
88 Nigeria.

89 **2.4 Induction of Diabetes**

90 All rats, except the Normal Control Group were intraperitoneally injected with 150 mg/kg
91 body weight of the prepared alloxan. Seventy two hours (three days) after alloxan
92 administration, the animals were fasted overnight and diabetes was confirmed from the rats

by measuring their fasting blood glucose level with the aid of a single touch glucometer. Rats that have fasting blood glucose level >7.0 mmol/l (126mg/dl) were considered diabetic and included in the study (Kandur and Goyal, 2005).

2.5. Experimental Design

By the end of the seven days acclimatization period, the animals were randomly assigned into five different groups of five rats each, designate as group A-E. Group A received water and feed only and serve as positive control. Diabetes was induced in group B, C, D, and E. Group B serve as negative control while group C to E corresponding to 200, 400 and 800 mg/kg doses of the aqueous leave extract of *Treculia Africana* were administered orally.

2.5. 0. Collection of Blood Sample

After four weeks of treatment all albino rats were fasted overnight, the rats where anesthetized by placing them in a seal cotton wool soaked in diethyl ether inhalation jar, the animals were sacrificed by decapitation at the end of three weeks of treatment and blood samples were obtained and centrifuged at 4000 ×g for 10 min at 4°C and supernatant kept at 37°C for further biochemical measurements. Different biochemical parameters were analyzed including estimation of fasting blood sugar level, Lipid profile, Heamatoligical and liver function indices.

2.5.1 Determination of Biochemical Parameters

2.5.2 Estimation of Serum Glucose Level

Serum glucose was estimated by glucose oxidase/ peroxidase method using Randox kit (Trinder, 1969).

Reagent composition (Randox Kit)

Contents	Concentration in the Test
R1. Buffer	
Pipes	100mmol/l, pH 7.6
ATP	4 mmol/l
NAD ⁺	3 mmol/l
Magnesium ions	15 mmol/l
R2. Enzyme Reagent	
Hexokinase	≥ 0.5 U/ml

G-6-PDH	≥ 1.5 U/ml
R3. PGI	
Phosphoglucose Isomerase	≥ 6.8 U/ml
CAL Standard	
Glucose	

115

116 2.5.3 Procedure

117 Test tubes were set up in triplicates and labeled blank, test and standard. 10µl of serum,
 118 standard (5.5 mmol/L) and distilled water were respectively pipetted in to the test tubes. Each
 119 test tube is then followed by 1000 µl of the reagent. The tubes were mixed properly,
 120 incubated at 37°C for 10 minutes and the absorbance of standard and tests read against the
 121 blank at 500nm using spectrophotometer.

122 **Calculation:** The glucose concentration was calculated using the relation:

$$123 \text{ Serum glucose (mmol/L)} = \frac{\text{Absorbance of Test}}{\text{Absorbance of Standard}} \times \text{conc of standard}$$

124 2.5.4 Estimation of Serum Total Cholesterol

125 Serum total cholesterol (TC) was estimated by enzymatic method using Enzopak kit (Allain
 126 *et al.*, 1974).

127 Reagent composition (Enzopak)

128

Reagent composition	Initial concentration
Active ingredients	
R1. Enzyme Reagent 1	
Cholesterol oxidase	≥ 500 U/l
Cholesterol Esterase	≥ 600 U/l
Peroxidase	≥ 6000 U/l
4-Amino Antipyrine	0.5 mmol/l
R2. Enzyme Reagent 2	
Buffer	100 mmol/l
Detergent	15 mmol/l

Phenol	20mmol/l
Surfactant	20mmol/l
pH 7.00±0.1 at 25°C	
Cholesterol standard(200mg/dl)	
Also contains non-reactive filters and stabilizers	

129

130 Three test tubes were set up and labeled blank, test and standard. In to test tubes labelled test,
 131 standard and blank, 10 µl of serum, standard (200 mg/dl) and distilled water were
 132 respectively pipetted in to the test tubes. Each test tube is then followed by 1000 µl of the
 133 reagent as shown above. The test tubes were mixed, incubated at 37°C for 5 minutes and the
 134 absorbance of the standard and test were read against the blank at 500 nm against the reagent
 135 blank.

136 **Calculation**

137 Cholesterol concentration was obtained using the relation:

$$138 \text{ Serum total cholesterol (mg/dl)} = \frac{\text{Absorbance of Test}}{\text{Absorbance of Standard}} \times \text{Conc of Standard}$$

139 **2.5.5 Estimation of Serum HDL – C**

140 This was done by enzymatic method of (Burstein *et al.*, 1970) using Randox Kit. Into
 141 centrifuge tubes, 200 µl of serum and 500 µl of precipitant (0.55 mmol/L phosphotungstic
 142 acid and 25 mmol/l Magnesium Chloride) were added, mixed and allowed to stand for 10
 143 minutes at room temperature. The tubes were centrifuged for 10 minutes at 4000 rpm. The
 144 supernatant was collected and used for the analysis. Three test tubes were set up and labeled
 145 blank, test and standard. In to test tubes labeled test, standard and blank, 10 µl of serum,
 146 standard and distilled water were respectively pipetted in to the test tubes. Each test tube is
 147 then followed by 1000 µl of the reagent. The test tubes were mixed, incubated for 5 minutes
 148 at 37°C and the absorbance of the samples and standard were measured against the reagent
 149 blank at 500nm.

150 **Reagent composition (Randox Kits)**

Contents	Initial
concentration	
R1. Enzyme Reagent 1	
N,N-Bis(2-hydroxyethyl)- 2-aminoethanesulfonic acid	100 mM, pH 6.6 (+25 °C)
N-(2-hydroxy-3-Sulfopropyl)- 3,5-dimethoxyaniline, sodium salt (HDAOS)	0.7 mM

Cholesterol Esterase [E.C.3.1.1.13. Microorganism]	≥800 U/L
Cholesterol Oxidase [E.C.1.1.3.6. <i>Streptomyces</i> sp]	≥500 U/L
Catalase [E.C.1.11.1.6. Microbial]	≥300 U/L
Ascorbate oxidase [EC.1.10.3.3. <i>Acremonium</i> sp.]	≥3000 U/L

R2. Enzyme Reagent 2

N,N-Bis(2-hydroxyethyl)- 2-aminoethanesulfonic acid	100 mM, pH 7.0 (+25 °C)
4-Aminoantipyrine	4.0 mM
Peroxidase [E.C.1.11.1.7, Horse Radish, +25 °C]	≥3500 U/L
Sodium Azide	0.05 w/v %
Surfactants	1.4 % w/v %

151

152 Calculation

153 The HDL-C concentration was obtained from the relation:

$$154 \quad \text{Serum HDL-C (mg/dl)} = \frac{\text{Absorbance of Test}}{\text{Absorbance of Standard}} \times \text{Conc. of Standard}$$

155 2.5.6 Estimation of Serum Triglyceride

156 This was assayed by the method of (Tietz, 1990) using Randox Kit.

157 Reagent composition (Randox Kit)

Contents	Concentration in the test
RIa. Buffer	
Pipes Buffer	40 mmol/l, pH 7.6
4-Chloro-phenol	5.5 mmol/l
Magnesium- ions	17.5 mmol/l
Rib. Enzyme Reagent	
4-aminophenazone	0.5 mmol/l
ATP	1.0 mmol/l
Lipase	≥150U/ml
Glycerol-kinase	≥ 0.4 U/ml
Glycerol-3-phosphate oxidase	≥ 1.5 U/ml
Peroxidase	≥ 0.5 U/ml
CAL Standard	11.95 mmol/l

158

The tubes were mixed and incubated at 37°C for 5 minutes and the absorbance of the standard and tests were read at 500nm against the blank.

Calculation: The TG levels were calculated using the relation:

$$\text{Serum TG (mg/dl)} = \frac{\text{Absorbance of Test}}{\text{Absorbance of Standard}} \times \text{Conc. of Standard}$$

2.5.7 Estimation of Serum LDL – C

This was calculated using Friedewald formula (Friedewald *et al.*, 1972).

$$\text{LDL - C (mg/dl)} = \text{TC} - (\text{HDL - C}) + \left(\frac{\text{TG}}{5} \right)$$

2.5.8 Estimation of Serum VLDL – C

This was calculated using Friedewald formula (Friedewald *et al.*, 1972)

$$\text{VLDL - C (mg/dl)} = \frac{\text{TG}}{5}$$

3.0 RESULT AND DISCUSSION

3.1 Table (1) Effect of alloxan on fasting blood sugar levels and lipid profile in alloxan induced diabetic albino rats

	Fasting Blood Sugar (FBS) (mg/dl)	Serum Triglyceride (TAG) (mg/dl)	Cholesterol (mg/dl)	High Density Lipoprotein (HDL)(mg/dl)	Low Density Lipoprotein (LDL) (mg/dl)
Non diabetes control	61.33±2.9a	1.58 ± 0.3a	3.11±0.6a	1.75±0.4a	1.04±0.2a
Diabetes control	268.84±0.3b	4.86±0.3b	5.98±0.3b	1.37±0.3b	3.31±0.1b

Values are expressed as mean ± SD, n= 5 for each group, Mean values having different superscript letter in the same column are significantly different at ($p<0.05$).

As shown in table 1, after induction of diabetes mellitus by alloxan, the serum glucose level raised to about 268.33 mg/dl with significant increase levels of serum triacylglyceride (TAG), cholesterol, and LDL- Cholesterol with significant decrease in HDL- Cholesterol level in all diabetes induced groups.

Fasting blood glucose concentration which is the most routine biochemical marker in the diagnosis of diabetes mellitus in clinical and experimental settings (Mayfield, 1998) was measured in this study. Table 1 revealed significant increased ($P < 0.05$) in fasting blood glucose concentration, triacylglyceride (TAG), total cholesterol and low density lipoprotein (LDL) cholesterol with significant ($P < 0.05$) decreased in HDL- cholesterol in all alloxan induced diabetes albino rats compared to non-diabetic control rats. Elevated value of fasting blood glucose concentration (268.33 mg/dl) observed in diabetic rat is due to the toxic effect of alloxan on islet beta cells of the pancreas (Ohno *et al.*, 2000) through its ability to induce reactive oxygen species (ROS) formation, resulting in the necrosis of the pancreas (Lenzen, 2008) and loss of capacity of the pancreas to secrete insulin (Zafar *et al.*, 2009) leading to hyperglycemia. Chronic exposure to hyperglycemia is the primary casual factor in the pathogenesis of diabetic complications and cause changes in vascular tissue that promote atherosclerosis (Rafiu and Lika, 2015). The elevated values for lipid profile parameters such as triglyceride, LDL-cholesterol, and total cholesterol observed in the alloxan induced diabetic rats could be partly due to increased intestinal biosynthesis of cholesterol (Luka *et al.*, 2013) because diabetes shifted the major site of cholesterologenesis from the liver to the small intestine (Young *et al.*, 1982) leading to hypercholesterolemia. Severe diabetes mellitus due to insulin deficiency might be accompanied with a reduced LDL-receptor (Swami *et al.*, 1996) resulting to high concentration of serum LDL cholesterol in diabetic subject.

3.2 Table (2) Effect of aqueous leaves extract of *Treulia africana* on fasting blood sugar levels in alloxan induced diabetic albino rats

Experimental groups	Fasting Blood Glucose levels (mg/dl)
Diabetic control	268.33 $\pm$ 16.8 ^{bcd}
Non-Diabetic control	61.43 $\pm$ 8.73 ^{acd}
Alloxan + 200mg/kg b.w of the <i>T. africana</i>	73.66 $\pm$ 5.77 ^{abd}
Alloxan + 400mg/kg b.w of the <i>T. africana</i>	66.01 $\pm$ 5.00 ^{abc}
Alloxan + 800mg/kg b.w of the <i>T. africana</i>	53.33 $\pm$ 5.50 ^{abcd}

Values are expressed as mean $\pm$ SD, n= 5 for each group, Mean values having different superscript letter are significantly different at ($p<0.05$).

Table 2, Revealed that oral administration of high dose of aqueous leaves extract of *Treculia africana* 800mg/kg body weight for twenty one (21) days to the diabetic rats significantly ($p<0.05$) decreased the elevated fasting blood sugar concentration compared with the diabetic control rats. Our result was in line with the finding of Ogbonnia et al (2008) which stated that, aqueous ethanolic extracts of *Treculia africana* Decne was the most powerful in amelioration hyperglycemia. According to researchers at the Departments of Physiology and Biochemistry, College of Medical Sciences, University of Calabar, who conducted a study using a *Treculia africana* (breadfruit) seed diet on rats, found that it significantly lowered blood lipid levels and blood glucose levels in diabetic rats compared to rats fed on normal diet (Efeooeye, 2013). Oral administration of high dose of aqueous ethanol extract of *Treculia africana* root bark (500mg/kg for 21days) to the diabetic rats ameliorated the diabetic complications by declined the glucose levels (Ogbonnia, et al., 2012) reflecting a restoration of the pancreatic β -cells activity (El-Desouki et al., 2015). Phytochemical screening revealed that, *Treculia Africana* was found to contain polyphenol and tannins (Ogbonnia et al., 2012; Osabor et al., 2009). Phenolic compounds might be useful medicinal food components and could contribute to manage both hyperglycemia and proper cellular redox status (Ali, 2013). Quercetin-3-0- glucoside, quercetin-3-0 (-6''- malonyl glucoside) and kaempferol-3-0 (-6''- malonyl glucoside) which are polyphenol compounds were reported to be responsible for hypoglycemic activity (Luangpiom et al., 2013) due to their inhibitory activity on α -glucosidase enzyme (Kim et al., 2008).

3.3 Table (3) Effect of aqueous leaves extract of *Treculia africana* on serum lipid profile in alloxan induced diabetic albino rats

Experimental groups	Total cholesterol (mg/dl)	TAG (mg/dl)	HDL- Cholesterol (mg/dl)	LDL- Cholesterol (mg/dl)
Diabetic control	5.86 $\pm$ 0.15 ^a	4.86 $\pm$ 0.15 ^b	1.37 $\pm$ 0.06 ^c	3.31 $\pm$ 0.72 ^d
Non- Diabetic control	3.11 $\pm$ 0.05 ^a	1.58 $\pm$ 0.50 ^b	1.75 $\pm$ 0.55 ^c	1.04 $\pm$ 0.55 ^d
Alloxan +	3.18 $\pm$ 0.65 ^b	1.55 $\pm$ 0.66 ^c	2.2 $\pm$ 0.70 ^d	0.67 $\pm$ 0.65 ^e

200mg/kg b.w. of the <i>T. africana</i>				
Alloxan + 400mg/kg b.w. of the <i>T. africana</i>	3.02 ± 0.60 ^{ab}	1.56 ± 0.51 ^{bc}	1.95 ± 0.61 ^{cd}	0.84 ± 0.55 ^{db}
Alloxan + 800mg/kg b.w. of the <i>T. africana</i>	3.13 ± 0.55 ^{ac}	1.4 ± 0.62 ^{bd}	1.9 ± 0.90 ^{ad}	0.84 ± 0.66 ^{dc}

Values are expressed as mean ± SD, n= 5 for each group, Mean values having different superscript letter in the same column are significantly different at ($p<0.05$).

Table (3) revealed that oral administration of aqueous extract of *Treculia africana* at a doses of 200, 400 800 mg/kg body weight for 21days to diabetic rats resulted in significant ($p<0.05$) reduction in total cholesterol, low density lipoprotein and triglyceride. These finding of the present study was in agreement with the finding of (Ogbonnia *et al.*, 2008; Ogbonnia *et al.*, 2011), this reduction in total plasma cholesterol, low density lipoprotein and triglyceride confirming the hypolipidaemic properties of *Treculia Africana*. This could be due to the presence of saponins that can inhibit taurocholate and deoxycholate absorption in a dose-dependent manner (Sheikhla, 2013). Phytochemical screening revealed that, *Treculia africana* was found to contain saponin (Ogbonnia *et al.*, 2008). Oral administration of *Treculia africana* leave extract to the diabetic rats ameliorated the diabetic complications by declined the LDL-Cholesterol, total cholesterol and triglyceride levels reflecting a restoration of the pancreatic β -cells activity (El-Desouki *et al.*, 2015) to secrete insulin. Insulin is a portent activator of lipoprotein lipase and inhibits VLDL production by the liver as well as to promote the reduction of LDL-cholestreol concentration (Galland, *et al.*, 2006). Thus, the effect of *Treculia africana* at reducing the LDL-cholesterol level of the diabetic rats may suggest the ability of the extract to increase the number of LDL receptors, with consequent reduction in LDL-cholesterol (Omega *et al.*, 2011), this confirming the hypoglycemic and hypolipidaemic properties of *Treculia africana* leaves.

3.3.1. Conclusion

Aqueous leave extract of *Treculia Africana* possess hypoglycemic effect and the most effective dose was 800mg/Kg body weight in amelioration of hyperglycaemia and most all toxicity effects of alloxan on lipid profile and some liver biomarker enzymes.

Consent: NA

References

- Abdel-Barry, J., Abdel-Hassan, I., & Al-Hakiem, M. (1997). Hypoglycemic and antihyperglycemic effects of *Trigonella foenum-graecum* leaf in normal and alloxan induced diabetic rats. *Journal of Ethnopharmacology* 58:, 149-155.
- Aja, P., Nwachuku, N., Ibiam, A., Igwenyi, I., & Onu, P. (2014). Comparative evaluation of transaminases and alkaline phosphatase activities in albino rats administered aqueous, ethanolic and methanolic extracts of *Moringa oleifera* seeds locally grown in Abakalaki, Nigeria. *J. Biol. Chem. Res.* 31 (1), 164-181.
- Ajibola, M., Eunice, O., & Stephanie, I. (2014). Effects of Aqueous Extract of *Moringa oleifera* Seeds on Alloxan Induced Hyperglycemia. *Basic Sciences of Medicine* 3 (3), 37-42.
- Ali, A. M. (2013). Anti-Diabetic Potential Of Phenolic Compounds: A Review. *International Journal of Food Properties*, 16, 91–103.
- Allain, C., Poon, L., Chan, C., Richmond, W., & Fu, P. (1974). Enzymatic Determination of Total Serum Cholesterol. *J. Clinical Chemistry*, 20: 470, 20, 470.
- Auwal, M., Tijjani, A., Sadiq, M., Saka, S., Mairiga, I., Shuaibu, A., et al. (2013). Antibacterial and Haematological Activity of *Moringa oleifera* Aqueous Seed extract in Wistar Albino Rats. *Sokoto Journal of Veterinary Sciences* 11 (1), 28-37.
- Bello, Aliero, A., Saidu, Y., & Muhammad, S. (2011). Phytochemical Screening, Polyphenolic Content and Alpha-Glucosidase Inhibitory Potential of *Leptadenia hastata* (Pers.) Decne. *Nigerian Journal of Basic and Applied Science* 19, 181-186.

- 277 Bello, Aliero, A., Saidu, Y., & Muhammad, S. (2011 a). Hypoglycaemic and Hypolipidaemic
278 Effects of *Leptadenia hastata* (Pers.) Decne in Alloxan. *Nigerian Journal of Basic and*
279 *Applied Science*, 19 (2), 187-192.
- 280 Bello, Aliero, A., Saidu, Y., & Muhammad, S. (2011 b). *Nigerian Journal of Basic and*
281 *Applied Science* 19:, 181-186.
- 282 Bennett, R., Mellon, F. A., Foild, N., Pratt, J., Dupont, M., Perkins, L., et al. (2013).
283 Profilling Glucosinolates and Phenolic in Vegetative and Reproductive Tissues of
284 Multipurpose Trees; *Moringa oleifera* L(Horseradish tree) and *Moringa stenopetala* L.
285 *Journal of Agriculture Food Chemistry* 51 (212), 3546-3556.
- 286 Berg, M., Tymoczko, L., & Stryer, L. (2002). *Biochemistry* (Fifth Edition ed.). Egland: W.H.
287 Freeman and Company.
- 288 Bukol, R. O., Adeleke, A. A., Omamuyovw, i. M., & Stephen, O. A. (2015). Lead-Induced
289 Damage on Hepatocytes and Hepatic Reticular Fibres in Rats; Protective Role of
290 Aqueous Extract of *Moringa oleifera* Leaves (Lam). *Journal of Biosciences and*
291 *Medicines*, 3, 25-37.
- 292 Burstein, M., Scholnick, H., & Morfin, R. (1970). Rapid Method for the Isolation of
293 Lipoproteins from Human Serum by Precipitation with Polyanions. *J. Lipid Res*, 11,
294 583-595.
- 295 Canargo, J., & Gross, J. (2014). Conditions Associated with Very low Values of
296 Glycohemoglobin Measured by an HPLC Method. *Journal of Clinical Pathology*, 57:
297 , 346-349.
- 298 Cennate, R., & Ebubekir, C. (2010). Effect of Streptozotocin on Biochemical Parameters in
299 Rats. *Asian Journal of Chemistry* 22 (3), 2375-2378.
- 300 Chukwuemeka, R. N., Danie, I. U., Michca, M., JeAnn, M., Rupika, D., Karen, T., et al.
301 (2012). Possible mechanisms of action of the aqueous extract of *Artocarpus altilis*
302 (breadfruit) leaves in producing hypotension in normotensive Sprague–Dawley rats.
303 *Pharmaceutical Biology*, 1096-1102.
- 304 David, K., Weber, K., Danielson, S., Wright, J., & Foley, E. (2002). Hematology and Serum
305 Biochemistry Values of rats. *Journal of Wild Life - Dis* 38 (3), 576-582.

306 Efeooeye. (2013). *African breadfruit (Gbeere): Nature's antidote to sleeplessness,*
307 *hypertension.* University of Calabar, Physiology and Biochemistry, College of
308 Medical Sciences . Calabar: tribune.com.ng.

309 Efiong, E., Igile, G., Mgbeje, B., Otu, E., & Ebong, P. (2013). Hepatoprotective and anti-
310 diabetic effect of combined extracts of Moringa oleifera and Vernonia amygdalina in
311 Streptozotocin-induced diabetic albino Wistar rats. 45-50.

312 Ekundina, V., Ebeye, O., Oladele, A., & Osham, G. O. (2015). Hepatotoxic and Nephrotoxic
313 Effects of Moringa oleifera Leaves Extract in Adult Wistar Rats. *Journal of Natural*
314 *Science Research vol.5 (3)*, 110-117.

315 El-Desouki, N. I., Aboulfotouh, B. M., Abdelmonaim, M. H., & El -Aama, M. S. (2015).
316 Moringa oleifera Leaf Extract Ameliorates Glucose, Insulin and Pancreatic Beta Cells
317 Disorder in Alloxan-Induced Diabetic Rats. *Research Journal of Pharmaceutical,*
318 *Biological and Chemical Sciences*, 6(3), 652.

319 Fahey, H. (2005). Moringa oleifera: A Review of the Nutritional Evidence for its Nutritional,
320 Therapeutic and Prophylactic Properties. Properties Part 1. *Trees of Life Journal*, 1
321 (3), 1-5.

322 Farooq, A., Sajid, L., Muhammad, A., & Anwarul, H. G. (2007). Moringa oleifera: A Food
323 Plant with Multiple Medicinal Uses . *Phytother. Res.* 21, , 17–25 .

324 Friedewald, W., Levy, R., & Fredrickson, D. (1972). Estimation of LDL-C in Plasma without
325 the use of the Preparative Ultracentrifuge. *Clinical Chemistry.*, 18(6), 499-502.

326 Galland, F., Duvillard, L., Petit, J., Lagrost, L., Vaillant, G., Brum, J., et al. (2006, July 12th).
327 Effect of insulin treatment on plasma oxidized LDL/LDL Cholesterol ration in types 2
328 diabetics patient. *Diabetes Metabolism*, 32, 625-631.

329 Gilani, H. A., Janbaz, H. K., & Shah, B. (1997). Quercetin exhibits hepatoprotective activity
330 in rats. *Biochem Soc Trans* 25, 85.

331 Gray, A., Abdel-Wahab, Y., & Flatt, P. (2000). The traditional plant treatment, Sabucus nigra
332 (Elder) exhibits insulin like and insulin releasing actions in vitro. *Journal of*
333 *Nutrition*. 130, 15-20.

334 Hajera, K., Ajijur, R., Mohitosh, B., & Anwar, U. (2011). Water-soluble Fraction of
 335 *Abelmoschus esculentus* L. Interacts with Glucose and Metformin Hydrochloride and
 336 Alters their Absorption Kinetics after Coadministration in Rats. *ISRN Pharmaceutics*,
 337 6, 260537.

338 Jain, P. J., Patil, S. D., Haswani, N., Girase, M. V., & Surana, S. (2010). Hypolipidemic
 339 activity of *Moringa oleifera* Lam., Moringaceae, on high fat diet induced
 340 hyperlipidemia in albino rats. *Braz. J.Pharmacog.*, 969-973.

341 Kamada, C., da Silva, E. L., Ohnishi-Kameyama, M., Moon, J. H., & Terao, J. (2005).
 342 Attenuation of lipid peroxidation and hyperlipidemia. *Free Radi. Res.*39: , 185–194.

343 Kankara, I., Wadatau, S., & Sado, M. (2016). Inhibitory Effect Of Aqueous Chloroform
 344 Leaves And Seed Extract Of *Moringa Oleifera* (Zogala) On The Activity Of Alpha
 345 Glucosidase. *International Journal of Scientific Research and Engineering Studies*
 346 3(5), 1-3.

347 Katzung, B. (1995). *Basic and Clinical Pharmacology, six Edition*. UK: Practice Hall
 348 International (UK) limited.

349 Kim KY, Nam KA, Kurihara H, Kim SM. Potent α -glucosidase inhibitors purified from the
 350 red alga *Grateloupia elliptica*. *Phytochem.* 2008;69:2820–5

351 Kothari, R., & Bokariya, p. (2012). A Comparative study of Haematological Parameters in
 352 Type 1 Diabete Mellitus Pateints And Healthy Young Adolescent. *Int. J.Biol. Med.*
 353 *Res.* 3 (4), 2429-2432.

354

355 Lange, A., & Momoh, J. (2015). Effects of Cinnamon Aqueous Extract on Blood Glucose,
 356 Liver Biomaker Enzymes, Hematological and Lipid Profile Parameters in Alloxan
 357 Induce Diabetic Albion Rats. *European Scientic Journal Vol.1*, 418-425.

358 Lenzen, S. (2008). Review: The mechanisms of alloxan- and streptozotocin-induced diabetes.
 359 *Diabetologia*, 51, 216–226.

360 Lin, X., Racette, S. B., Lefevre, M., Spearie, C. A., Most, M., Ma, L., et al. (2010). The
 361 effects of phytosterols present in natural food matrices on cholesterol metabolism and

LDL-cholesterol: a controlled feeding trial. *European Journal of Clinical Nutrition* 64,
1481–1487.

Luangpiom, A., Kourjampa, W., & Junaimaung, T. (2013). Anti-hyperglycemic Properties
of *Moringa oleifera* Lam. Aqueous Leaf Extract in Normal and Mildly Diabetic Mice.
British J. Pharmacol. Toxicology 4(3), 106-109.

Luka, C., Tijjani, H., Joel, E., Ezejiofor, U., & Onwukike, P. (2013). Hypoglycaemic
Properties of Aqueous Extracts of *Anacardium occidentale*, *Moringa oleifera*,
Vernonia amygdalina and *Helianthus annuus*: A Comparative Study on Some
Biochemical Parameters in Diabetic Rats . *International Journal of Pharmaceutical
Science Invention*, Volume 2 Issue 7, 16-22.

Magandala, D. S., Bellamakonda, R., & Kumari Dasireddy, S. (2012). Evaluation of
antidiabetic and antihyperlipidemic potential of aqueous extract of *moringa oleifera*
in fructose fed insulin resistant and stz induced diabetic wistar rats: a comparative
study. *Asian Journal of Pharmaceutical and Clinical Research* Vol 5, Issue 1., 67-72.

Magee, P. N. (1969). Nitroso compounds. *Br. Med. Bull.*, (25), 240-244.

Mayfeild, J. (1998). Diagnosis and classification of diabetes; New criteria. *Family of
physiology* 58 (6), 1355-1362.

Miikue-Yobe, T., Uwakwe, A. A., & Akaninwor, J. O. (2015). EFFECT OF AQUEOUS
LEAF EXTRACT OF *HEINSIA CRINATA* ON HAEMATOLOGICAL AND SOME
BIOCHEMICAL INDICES OF TOXICITY IN STREPTOZOTOCIN INDUCED
DIABETIC RATS. *International Journal of Scientific Research and Innovative
Technology* 2(1), 116-126.

Mostafa, M. A., Serwah, A. A., Mohammed, S. W., & Mohammed, M. K. (2013). EFFECTS
OF SOME ANTIDIABETIC MEDICINAL PLANTS ON PANCREAS AND LIVER
OF DIABETIC ALBINO RATS. *The Egyptian Journal of Hospital Medicine* Vol. 50,
156– 168.

Murray, K. R., Bender, D., Botha, M., Kennelly, R. W., & Weil, A. (2002). *Harper's
Illustrated Biochemisry* (28th Edition ed.). England: MC Graw Hill.

390 Noor, A., Gunasekaran, S., Manickam, A., & Vijayalakshmi, M. A. (2008). Antidiabetic
 391 activity of Aloe vera and histology of organs in streptozotocin –induced diabetic rats.
 392 *Curr. Sci.* 94:, 1070– 1076. .

393 Ochei, J., & Kollhatkar, A. (2007). *Medical laboratory science theory and practice*. 7 West
 394 patel. New Delhi: Tata McGraw-Hill.

395 Ogonnia, S. O., Anyika, E. N., Mbaka, G. O., Utah, P., Ugwu, D., Nwakakwa, N., et al.
 396 (2012). Antihyperglycaemic and antihyperlipidaemic effects of aqueous ethanol
 397 extract of *Tapinanthus globiferus* leaves and *Treculia africana* root bark and their
 398 mixture on alloxan diabetic rats. *Agriculture And Biology Journal Of North America*,
 399 3(6), 237-246.

400 Ogonnia, S. O., Enwuru, N. V., Onyemenem, E. U., & Oyedele, G. A. (2008).
 401 Phytochemical evaluation and antibacterial profile of *Treculia africana* Decne bark
 402 extract on gastrointestinal bacterial pathogens. *African Journal of Biotechnology*,
 403 7(10), 1385-1389.

404 Ogonnia, S., Odimegwu, J., & Enwuru, V. (2008). Evaluation of Hypoglycemic and
 405 Hypolipidemic Effects of Aqueous Ethanolic Extract of *Treculia africana* Decne and
 406 *Bryophyllum Pinnatum* Lam and Their Mixture on Streptozotocin(STZ)-induced
 407 Diabetic Rats. *Afr. J. Biotechnol.*, 15(7), 2535-2539.

408 Ohno, T., Horio, F., Tanaka, S., Terada, M., & Namikawa, T. a. (2000). Fatty liver and
 409 hyperlipidemia in IDDM (insulin dependent diabetes mellitus) of Streptozotocin
 410 treated shrews. *Journal of Life Science*, 125-131.

411 Ojokoh, A., Fayemi, O., & Ocloo, F. (2014). Proximate Composition, Antinutritional contents
 412 and Physicochemical properties of breadfruit (*Treculia africana*) and Cowpea (*Vigna*
 413 *Ungusticulata*) flour blends fermented with *Lactobacillus plantarum*. *African Journal*
 414 *of Microbiology Research*, 1352-1359.

415 Olapade, A., & Umeonuorah, U. (2014). Chemical and Sensory Evaluation of African
 416 Breadfruit (*Treculia africana*). *Nigerian Food Journal*, 80-88.

417 Omaye, K., Onoagbe, O. I., Georgina, E., Esosa, S. U., Ajeigbe, O. K., & Amegor, O. F.
 418 (2011). Effects of aqueous root extract of *Treculia africana* on blood glucose, lipid

419 profile and body weight changes of streptozotocin-induced diabetic and normal rats.
 420 *International Journal of Plant Physiology and Biochemistry*, 3(10), 169-175.

421 Oparinde, D., Atiba, A. S., Ajose, O., Eludoyin, A., & Adesiyan, A. (2014). Effect of
 422 *Moringa oleifera* leaf extract on serum lipids and glycaemic control in alloxan
 423 induced diabetic albino rats. *International Journal of Biomedical And Advance*
 424 *Research* 05 (10), 520-522.

425 Ortiz-Andrade, R., Rodriguez-lopez, V., & Estrack-Soto, S. (2005). Antidiabetic Effects on
 426 Alloxanized and Normaglycemic rat and Some Pharmaceutical Evaluation of
 427 *Tounetorniam Hartmegical*. . *Journal of Ethnopharmacology*: 100(1-3), 74-42.

428 Orwa, C., Mutua, A., Kindt, R., R., J., & Anthony, S. (2009). Agroforestry Database. A Tree
 429 *Reference and Selection Guide*, 1-5.

430 Osabor, V. N., Ogar, D. A., Okafor, P. C., & Egbung, G. E. (2009). Profile of the African
 431 Bread Fruit (*Treculia africana*). *Pakistan Journal of Nutrition*, 10005-1008.

432 Osabor, V., Ogar, D. A., Okafor, P. C., & Egbung, G. (2009). Profile of African bread fruit
 433 (*Treculia africana*). *Pakistan Journal of Nutrition*, 8(7), 1005-1108.

434 Oyedemi, S., E.A, A., Aijegro, O. A., & Akinpelu, D. (2011). Antidaibetic and
 435 Heamatolpgical Effect of Aqueous Extract of Stem Bark of *Alzelia africana* (smith)
 436 on Streptozotcin-Induced Diabetic Wistar Rats. *Asian Pacific Journal Tropical*
 437 *Biomedicine* 1 (5), 353-358.

438 Oyelola, O. O., Moody, J. O., Odeniyi, M. A., & Fakeye, T. O. (2007). Hypoglycemic Effect
 439 Of *Treculia Africana* Decne Root Bark In Normal and Alloxan-Induced Diabetic Rats.
 440 *Afr. J. Traditional, Complementary and Alternative Medicines*, 387 - 391.

441 Patil, R., Ahirwar, B., & Ahirwar, D. (2011). Current status of Indian medicinal plants with
 442 antidiabetic potential: A review. *Asian Pacific Journal of Tropical Biomedicine*. 1(2):
 443 , 5291–5298.

444 Petit, P., Baissac, Y., Ponsin, G., Ribes, G., Sauvaire, Y., Hillaire-Buys, D., et al. (1995).
 445 Steroid saponins from fenugreek seeds: Extraction, purification and pharmacological
 446 investigation on feeding behavior and plasma cholesterol. *Steroid* 60:, 674-680.

- 447 Piyachaturawat, P., Poprasit, J., & Glinsukon, T. (1990). The Gastric mucosal secretions and
448 lesions by different doses of Streptozotocin in rats. *Toxicol. Let.*, 55, (21-29).
- 449 Piyachaturawat, P., Poprasit, J., Glinsukon, T., & Warichanon. (1988). *Cell Biol. Intern. Rep.*,
450 12(1), 53-63.
- 451 Rafiu, A., & Luka, C. (2015). ANTI-DIABETIC ACTIVITY OF AQUEOUS SEED
452 EXTRACT OF (*Moringa oleifera*) IN NORMAL AND ALLOXAN INDUCED
453 DIABETIC RAT. *Journal of Biological Sciences and Bioconservation, Volume 7,*
454 *NUMBER 1*, 24-37.
- 455 Rashmi, Y., & Rahul, k. (2011). A study of phytochemical constituents and pharmacological
456 actions of *trigonella foenumgraecum*: A review. *International Journal of Pharmacy*
457 *and Technology 3 (2)*, 1022-1028.
- 458 Rasmi, Y., & Rahul, K. (2011). A study of phytochemical constituents and pharmacological
459 actions of *Trigonella foenum graecum*: A review. *International journal of pharmacy*
460 *and technology*, 1022-1028.
- 461 Richert, K., Martinez., R. R., & Martinez, T. T. (1999). Pharmacist knowledge of common
462 herbal preparations. *Proc. West. Pharmacol. Soc. 42*., 42, 1-2.
- 463 Rivera, L., Moron, R., Sanchez, M., Zarzuelo, A., & Galisteo, M. (2008). Quercetin
464 ameliorates metabolic syndrome and improves the inflammatory status in obese
465 Zucker rats Obesity. *Silver Spring*, 2081-2087.
- 466 Ruckmani, K., Kavimani, S., Anandan, R., & Jaykar, B. (1998). Effect of *Moringa oleifera*
467 Lam on paracetamol-induced hepatotoxicity. *Indian J Pharm Sci 60*: , 33–35.
- 468 Ruffo, C., Birnie, A., & Tengas, B. (2001). *Edible Wild Plants of Tanzania*. Nairobi:
469 Regional Land Management.
- 470 Saidu, Y., Nwachukwu, F., Bilbis, L., Faruk, U., & Abbas, A. (2010). Hypoglycaemic and
471 Hypolipidemic Effects of Root Extracts of *Albizzia chevalieri* in Alloxan-induced
472 Diabetic Rats. *Nigerian Journal of Basic and Applied Science*, 18(1), 72-78.
- 473 Sheikhla, A. (2013). *Trigonella foenum-graecum* L. (Fenugreek) as a Medicinal Herb in
474 Animals Growth and Health. *Science International Volume 1 Issue 6*., 1994-1998.

475 Sherifat, A. A., Olusegun, E., & Oluwole, O. (2007). Constituents of Breadfruit Leaves,
476 Stem, and Root barks. *Journal of Essential Oil Bearing Plants*, 189-193.

477 Stark, A., & Madar, Z. (1993). The effect of an ethanol extract derived from fenugreek on
478 bile acid absorption and cholesterol level in rats. *British Journal of Nutrition*:69,
479 277-287.

480 Swami, S., Sztalryd, C., & Kraemer, F. B. (1996). Effects of streptozotocin-induced diabetes
481 on low density lipoprotein receptor expression in rat adipose tissue. *Journal of Lipid*
482 *Research*, 37, 229-239.

483 Tietz, N. (1990). *Serum Triglyceride Determination*. In: *Clinical Guide to Laboratory Tests*
484 (Second Edition ed.). Philadelphia, Philadelphia, USA: W.B., Saunders Co.

485 Trinder, P. (1969). Determination of Blood Glucose in Blood Using Glucose Oxidase with an
486 Alternative Oxygen Acceptor. *Annals of Clin. Biochem*, 6, 24-25.

487 Uko, E., Erhabor, O., Isaac, I., Abdulrahman, Y., Adias, T., Sani, Y., et al. (2013). Some
488 Haematological Parameters in Patients With Type – I Diabetes in Sokoto, North
489 Western Nigeria. *J. Blood Lymph*, 3(110), 2165-7831.

490 Vos, T., Flaxman, A., Naghavi, M., Lozano, R., Michaud, C., Ezzati, M., et al. (2012). Years
491 lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990–
492 2010: a systematic analysis for the Global Burden of Disease Study. *Lancet* 380
493 (9859), 2163–96.

494 Xue, X., Li, W., Zhang, L., Liu, Y., Wang, Z., & Zhang, R. (2007). Effect of Trigonella
495 foenum-graecum (fenugreek) extract on blood glucose, blood lipid and
496 hemorheological properties in streptozotocin-induced diabetic rats. *Asia Pacific*
497 *Journal Clinical Nutrition*:16, 422-426.

498 Young, N. L., Saudek, C. D., & Crawford, S. A. (1982). Total hydroxymethylglutaryl CoA
499 reductase activity in the small intestine and liver of insulin-deficient rats. *Journal of*
500 *Lipid Research*, 23, 266-275.

501 Zafar, M., Naeem-ul-hassan Aaqvi, A., & Ahmed, A. &. (2009). Altered Liver Morphology
502 and Enzymes in Streptozotocin Induced Diabetic Rats. *International Journal of*
503 *Morphology* 27(3) , 719-725.

504 Zafar, M., Naqvi, *. N.-u.-H., & Kaimkhani, M. A. (2009). Altered Liver Morphology and
505 Enzymes in Streptozotocin Induced Diabetic Rats. *International Journal of*
506 *Morphology* 27(3), 719-725.

507 Zhang, A., Vetromenn, J., Van Gaal, L., & De Leeuw, I. (1995). Effects of pravastatin on
508 lipid levels in vitro oxidizability of non-HDL lipoproteins and microalbuminuria in
509 IDDM patients. *Diabetes Res. Clin. Pract.*, 29(3), 189-194.

510 Zhang, M., Swarts, S. G., Yin, L., Liu, C., Tian, Y., Cao, Y., et al. (2011). Antioxidant
511 properties of quercetin. *Adv. Exp. Med. Biol.* 915:, 283–289.

512

513