

Mandibular Second Molar with Three Mesial Canals:A Case Report

Abstact

Most mandibular second molar have two canals in mesial root and one in distal.This report represents a case of mandibular second molar with three mesial and two distal root canal.To achieve the ideal goal of endodontic treatment,clinicians should be aware of the complicated configuration of the root canal system and use suitable methods to identify the accessory canals that exist possibly.

Key words:mandibular seond molar;root canal treatment;the third mesial canal;dental operating microscope(DOM);cone-beam computed tomography(CBCT)

introduction

To achieve a successful root canal treatment,accomplishing detection,cleaning,shaping and complete obturation to the entire canal system is of most importance.Thus,the anatomy of the root canals such as the root numbers,numbers of canals in each root and their types and shapes plays a key role in the success of endodontic treatment.

Variations in the canals of the mandibular second molar have been reported by a number of investigators.It is described that the root and canal morphology is different according to the race and genetic[1]. The most common morphology was the presence of two canals in the mesial root and one in the distal root of the teeth with separate roots[2 3]. Otherwise,mandibular second molar commonly have fused roots and there was a high prevalence of C-shaped canals in Chinese population(39%-45.5%)[2 4 5 6].In addition,there were other types such as 2 roots and 2 canals,2 roots and 4 canals,3 roots and 3 canals,3 roots and 4 canals,etc[7 8 9]. While the presence of a third canal in the mesial root of the mandibular second molar with 2 roots and 5 canals has been reported rarely,to have an incidence of 0%-16.7%[1 10 11].Compared with distal root canals,the mesial root in mandibular molars is more complex than that was been commonly considered.[12]

This report describes a case of a mandibular second molar with two roots and five canals (three canals in the mesial root and two in the distal root)and discusses the significance of treating such cases.

Case report

A 68-year-old female patient presented to the Department of Endodontics of the JiLin University with a history of recurring painful swelling for 2 years and with a fistula on the buccal side in the lower right molar area.Her medical history had no relevant contribution.The mandibular right second molar was tender on vertical percussion and have a gomphiasis at I degree.A radiograph revealed deep carious lesion in distal proximal area approximating the pulp space area of teeth#47[Figure 1].A diagnosis of an chronic periapical periodontitis was made for the mandibular right second molar.

Local anesthesia(2% lidocaine with 1:40,000 adrenaline)was performed.The tooth was isolated by a


rubber dam,all carious tissue was removed and then an adequate access cavity was prepared.DG-16 Endodontic explorer(Hu-Freidy,USA)located the extra-canal orifice under dental operating microscope(DOM).The CBCT scan demonstrated there are there canals in mesial root and two canals in distal root.

After canal identification and negotiation,the working length was determined by an electronic apex locator (Propex II ,Dentsply Mailefer,israel).All canals were cleaned and shaped with ProTaper NiTi rotary instruments(Dentsply-Maillefer,Ballaigues,Switzerland)under copious irrigation with 1%sodium hypochlorite and saline.Canals were dried with paper point and temporized with calcium hydroxide(Pulpdent,USA)


At the second appointment after a week,all the canals were recapitulated,irrigated,dried and an angled radiograph was taken with master cones in all canals[Figures 5 and 6].The canals were obturated with continuous wave condensed gutta percha (0.04taper,Gapadent,China)and roeko sealer,and the access was closed with temporary restoration.A post-obturation angled radiograph was taken.The patient experienced no postoperative sequelae and was scheduled for appropriate coronal restoration.The efficacy and prognosis will be observed by follow up.



Figure.1 pre-operative radiograph



57




58

59 figure.2 pre-operative CBCT

60

61

62




63

64 figure.3 access opening


65

66



67
68

Figure.4 master cone in angled radiograph



69
70
71
72

Figure.5 CBCT after oburation




Figure.6 root canal orifice



figure.7 obturation

Discussion

One of the most common reason for the failure of endodontic treatment was missing canals,the bacteria might be located in such undetected or unprepared portion of the root canal,leading to the chronic symptoms and unhealed periapical lesions[13]. Just as this case,the complex nature of the root and root canal morphology of the mandibular second molars,especially the mesial root,has been underestimated easily. There are studies report 7.52% of the orifices were deeper than 2 mm so they could not be revealed via troughing[14.]Separation,curvature and fusion are more likely to occur to this location,so as a clinician,we should not only memorize the common anatomical configurations,but also identify the unusual ones.For the mesial root system of mandibular molars,besides mesiolingual(ML)

canal,mesiabuccal(MB) canal,sometimes its own an accessory mesial canal.The distribution pattern of the third canals was similar in mandibular molars.45%located closer to ML canals,30%located in the middle between the ML and MB canals(mesial middle canals),and 25%located closer to the MB canals.[12 15]

Every attempt should be made to find all canals to ensure successful endodontic treatment.Hence,some auxiliary tools in procedures may be necessary.It was demonstrated that the root canal orifices was more accurately detected under a dental operating microscope (DOM) than that with naked eye and the surgical loupes,[16 17]increased the detection rate to 93%,even if canal anatomy was complicate.[18]The two-dimensional radiography is the most conventional technique in endodontics therapy,however,it's difficult to achieve angled radiography,the overlapping and distortion of roots and canals may influence the judgment of clinicians.[18]The distinct advantages of cone beam computed tomography(CBCT) is that provides reliable information about the accessory roots and canals in three dimensions clearly without destructiveness,[7 8 18 19 23]and allow clinicians manipulate the images at different sections. The CBCT could detect the periodontal hard tissue damage which is inapparent in conventional radiography.[21]Compared with traditional CT,CBCT appeared a lower cost and radiation dose,was helpful and suitable in the field of endodontics.

Some diverged canal at a deep level and tiny ones only could be identified after removing precisely the dentin coverage with ultrasonic instrument and shaping.[18 23]Trying to detect and identify the aberrant canals that possibly exist should be persisted throughout the root canal treatment.

Conclusion

Although it's a extremely rare case,a five-canaled configuration may be exist in the mandibular second molar.The clinicians should aware of these accessory canals in root canal treatment and use suitable diagnostic methods to identify and detect to ensure the success at utmost.

References

- [1]Zare Jahromi M,Jafari Golestan F,Mashhadi Esmaeil,et al.Root and Canal Morphology of Mandibular Second Molar in an Iranian Population by Clearing Method.J DENT,2013,14:78-81.
- [2]Emmanuel João Nogueira Leal Silva,Yuri Nejaim,Amaro V.Silva,Francisco Haiter-Neto,Nestor Cohenca.Evaluation of Root Canal Configuration of Mandibular Molars in a Brazilian Population by Using Cone-beam Computed Tomography:An In Vivo Study.J Endod,2013,39:849-852.
- [3]Mohamed Idris,Nasil Sakkir,Nandakishore KJ,Annapurna Kini.Endodontic Retreatment of a Mandibular Second Molar with four Separate Roots:A Case Report.Journal of Clinical and Diagnostic Research,2014,8:280-282.
- [4]Q.Zheng,L.Zhang,X.Zhou,et al.C-shaped root canal system in mandibular second molars in a Chinese population evaluated by cone-beam computed tomography.International Endodontic Journal,2011,44:857-862.
- [5]Jafarzadeh,H(Jafarzadeh,Hamid),Wu,YN(Wu,You-Nong).The C-shaped root canal configuration:a review.Journal of Endodontics,2007,33:517-523.
- [6]Jin GC,Lee SJ,Roh BD.Anatomical study of C-shaped canals in mandibular second molars by analysis of computed tomography.Journal of Endodontics,2006,32:10-13.
- [7]Emmanuel João Nogueira Leal Silva,Yuri Nejaim,Amaro V. Silva,Francisco Haiter-Neto;Nestor Cohenca.Evaluation of Root Canal Configuration of Mandibular Molars in a Brazilian Population by Using Cone-beam Computed Tomography: An In Vivo Study.Journal of Endodontics,2013,39: 849-852.

- [8]Plotino G,Tocci L,Grande NM,Testarelli L,et al.Symmetry of Root and Root Canal Morphology of Maxillary and Mandibular Molars in a White Population: A Cone-beam Computed Tomography Study In Vivo.J Endod,2013,39: 1545-1548.
- [9]S.S.Patil.et al.The study of the morphologic variations in the root canal of human permanent mandibular second molar.Journal of Pierre Fauchard Academy,2009,23:97-99.
- [10]Sara A.Barsness,Walter R.Bowles,Alex Fok,et al. An anatomical investigation of the mandibular second molar using micro-computed tomography[J].Surgical and Radiologic Anatomy,2015,37:267-272.
- [11]Ozkan Miloglu,Hakan Arslan,Cagatay Barutcgil,Kenan Cantekin.Evaluating root and canal configuration of mandibular first molars with cone beam computed tomography in a Turkish population[J].Journal of Dental Sciences,2013,8:80-86.
- [12]Ali Nosrat,Raney J.Deschenes,Patricia A.Tordik,et al.Fouad.Middle Mesial Canals in Mandibular Molars:Incidence and Related Factors[J]. J Endod 2015,41:28-32.
- [13]Minju Song,Hyeon-Cheol Kim,Woocheol Lee,et al.Analysis of the Cause of Failure in Nonsurgical Endodontic Treatment by Microscopic Inspection during Endodontic Microsurgery. J Endod,2011,37:1516-1519.
- [14] Keleş A, Keskin C. Detectability of Middle Mesial Root Canal Orifices by Troughing Technique in Mandibular Molars: A Micro-computed Tomographic Study.[J]. Journal of Endodontics, 2017,43:1329-1331.
- [15]Karapinar Kazandag M,Basrani BR,Friedman S.The operating microscope enhances detection and negotiation of accessory mesial canals in mandibular molars. J Endod,2010,36:1289-1294.
- [16]Takatomo Yoshioka,Chihiro Kobayashi and Hideaki Suda.Detection Rate of Root Canal Orifices with a Microscope. J Endod,2002,28:452-453.
- [17]de Carvalho MC,Zuolo ML.Orifice locating with a microscope. J Endod,2000,26:532-534
- [18]Kurthukoti AJ,Sharma P,Swamy DF,et al.Computed Tomographic Morphometry of the Internal Anatomy of Mandibular Second Primary Molars.Int J Clin Pediatr Dent,2015,8:202-207.
- [19]Ran Yang,Chan Yang,Yuan Liu,Yong Hu,Jing Zou.Evaluate root and canal morphology of primary mandibular second molars in Chinese individuals by using cone-beam computed tomography.Journal of the Formosan Medical Association,2013,112:390-395.
- [20]Pita PARDO, Milena del Carmen,NISHIYAMA, Celso Kenji,et al.CBCT and microscopic analysis of the incidence of second mesiobuccal canal of maxillary molars.Dental Press Endodontics,2012,2: 61-66
- [21]Seema Dhananjay Pathak,Pradnya Vilas Bansode,Shraddha Pradeep Gite,et al. Cone Beam Computed Tomography:A New Boon and a Ray of Hope to the Endodontist—A Series of Cases-Review Article.International Journal of Prosthodontics and Restorative Dentistry,2015,5:101-109.
- [22]Flávia G.R.Cardoso,Nádia S.Ferreira,Frederico C.Martinho,et al. Correlation between Volume of Apical Periodontitis Determined by Cone-beam Computed Tomography Analysis and Endotoxin Levels Found in Primary Root Canal Infection.J Endod,2015,41:1015-1019.
- [23]Vertucci FJ.Root canal anatomy of the human permanent teeth.Oral Surgery, Oral Medicine, Oral Pathology,1984, 58: 589-599.