

BILIARY ATRESIA PRESENTING AS POST CIRCUMCISION EXSANGUINATION

ABSTRACT

An autopsy diagnosis of biliary atresia was made in a month old male infant who died following prolonged bleeding that lasted for about 8 hours after circumcision by a non-trained local circumcision practitioner.

Keywords:-circumcision, biliary atresia

CASE PRESENTATION

A 4 weeks old male delivered by a traditional birth attendant, was circumcised by a non-trained local circumcision practitioner (the only person performing such in a local village). Four hours later, the child presented with complaints of continued bleeding from the site of circumcision. The nurse applied vitamin K and penicillin ointment topically and sent the baby home. Approximately 3-4 hours later the baby died.

At the autopsy venue, relevant information was collected (first contact of the parents with a medical doctor since the child's delivery) were as follows: the child was delivered at 9 months (probably at term), the circumcision was delayed for 3 weeks because around 7 days of age, "the child was not strong enough" (in the words of the mother), was the 4th child in a family of 2 boys and 2 girls, the elder brother did not bleed much after circumcision and the stool was said to be pale coloured (non pigmented) in recent times.

AUTOPSY FINDINGS

Gross anatomy findings: An apparently healthy looking male infant was seen. The head circumference was 34cm, mid-arm circumference was 14cm and the length was 1.9ft. The baby was moderately jaundiced and the glands penis and site of circumcision were clean (no haematoma). All the organs were present, well formed and in their normal anatomical positions. The organs were very pale and there was no excessive fluid collection in any of the serosal spaces. The lungs were hyperventilated (with prominent rib markings), while the liver (weighed 140g) had a smooth shiny greenish surface with a homogenous greenish cut surface (figure 1a-c). There was no intra-cranial bleeding or bleeding elsewhere

Microscopic findings: The brain, heart, kidney, intestines, and lungs showed features of hypoxic/ischaemic injury. Sections of the liver showed bile duct proliferation, mixed inflammatory cell infiltration of the portal tract, grade 2 bridging fibrosis and bile plugging (features of biliary atresia (BA)). The autopsy diagnosis was concluded as massive haemorrhage (hypovolaemic shock) post-circumcision due to BA and defective hepatocyte function.

Informed written consent was obtained from the parents on with respect that the identity must remain anonymous.

LITERATURE REVIEW

Male Circumcision is one of the oldest and one of the most commonly performed surgical procedures in practice today. It is commonly practiced for religious, cultural and medical reasons.¹⁻² Complications of circumcision are broadly classified into early and late complications. Early complications include; bleeding (most common), pain, inadequate skin

45 removal, and surgical site infection. Others include chordee, iatrogenic hypospadias,
46 glandular necrosis, and glandular amputation. Post-circumcision bleeding in patients with
47 coagulation disorders can be significant and sometimes even fatal.^{2,3} Late complications of
48 circumcision include; epidermal inclusion cysts, suture sinus tracts, chordee, inadequate skin
49 removal resulting in the redundant foreskin, penile adhesions, phimosis, buried penis,
50 urethrocutaneous fistulae, meatitis, and meatal stenosis.^{2,3} Bleeding remains the commonest
51 complication encountered during and after circumcision.³ In most cases of bleeding, which is
52 usually minor, all that is required to achieve haemostasis is gentle pressure on the
53 area.² Excessive bleeding may be due to anomalous blood vessels or presence of bleeding
54 disorders.⁴ In situations where gentle pressure fails, the circumferential bandage may be used
55 to aid haemostasis.² Pharmacological agents like 1:100, 000 adrenaline can also be used.
56 Diathermy or suturing may be used for obvious bleeding vessels.² There are very few
57 documented cases of exsanguination following circumcision. **Death following circumcision is**
58 **a very rare occurrence.**⁵

59 **BA** is the single most frequent cause of death from liver disease in early childhood.⁶ BA is the
60 complete or partial obstruction of the lumen of the extrahepatic biliary tree within the first 3
61 months of life, that is characterized by progressive inflammation and fibrosis of intrahepatic
62 or extrahepatic bile ducts.⁷ BA mainly involves extrahepatic ducts, though intrahepatic ducts
63 can also be involved and are usually seen in about 1 in 10,000 births.⁸ There are two major
64 forms of biliary atresia (fetal and perinatal), based on the presumed timing of luminal
65 obliteration.⁶

66 The etiology of BA is not clear, though **aflatoxins**, toxic, infectious and autoimmune factors
67 have been implicated.^{6,9} Clinically, BA **can be seen in any sex**. The infants are usually
68 delivered at full term and jaundice develops during 2nd to 3rd week, while the stool colour is
69 usually acholic.¹ The index case was delivered at term, was jaundiced (onset not known) and

70 passed pale coloured stool (acholic). The fetal form is a rare form and usually associated with
71 other anomalies like interrupted inferior vena cava, polysplenia, congenital heart disease and
72 malrotation of abdominal viscera. It is seen in about 20% of cases. The suspected cause is
73 the aberrant intrauterine development of the extrahepatic biliary tree. The perinatal form of
74 BA is commoner, in which a presumed normally developed biliary tree is destroyed
75 following birth.¹⁰ The main cause of BA is unknown, though autoimmunity, viral infections
76 (reovirus, rotavirus, and cytomegalovirus) and toxic factors are known to play roles.^{8,10} The
77 index case is probably a perinatal form BA, because the features listed above for fetal form
78 were not seen.

79 Grossly, the liver surface is smooth and homogenously green. The case in the study was same
80 as shown in figure 1a. Micro morphologic features of BA which are evident in about two-
81 thirds of cases are marked bile ductular proliferation, portal tract oedema and fibrosis,
82 inflammation of the hepatic or common bile ducts, parenchymal cholestasis, and progressive
83 destruction of the intrahepatic biliary tree. If BA is unrecognized or uncorrected, cirrhosis
84 develops within 3 to 6 months of birth.^{6,11} These features were characteristically seen in the
85 index case with some bridging fibrosis. (Figures 2a & b, 3)

86 Clinically these infants appear almost normal at birth with normal birth weight and postnatal
87 weight gain. Initially, they pass normal coloured stools in the early days of life before it
88 progresses to acholic stools as the disease evolves.^{6,9} The index case though a male, was
89 claimed to be normal at birth by parents, though birth weight was not measured and passed
90 normal coloured stool (yellowish brown) initially. Investigations that aid diagnosis include:
91 elevated hyperbilirubinemia, moderately elevated aminotransferase and alkaline phosphatase
92 levels.^{6,9} Other investigations that aid diagnosis are ultrasonography, percutaneous liver
93 biopsy and technetium 99m hepatobiliaryiminodiacetic acid (HIDA), though more definitive

94 procedure for the diagnosis of BA currently is laparoscopic guided cholecysto
95 cholangiography.¹² No investigation was done in the index case.

96 Treatment of choice is hepatoportoenterostomy (kasaiprocedure) ie connecting the bowel lumen
97 with the porta hepatitis, or else patients die from liver cirrhosis before 2 years of age.⁶ The
98 success of this procedure is dependent on the level and variability in the anatomy of BA.⁶ For
99 those that may not benefit from Kasai procedure, the other alternative is liver transplant.^{6,9}


100 It was not seen in literature were circumcision lead to death in a BA patient, because in such
101 circumstances, the patients must have been stabilized and proper precautions taken. The
102 index case had no antenatal care, was not delivered in a hospital and never presented to a
103 hospital or qualified health worker.

104 Conclusion

105 In conclusion the post circumcision exsanguination seen in this case was probably due to
106 reduced / impaired hepatic functions with compromised production of clotting factors.
107 Though liver function test, clotting factor estimation and clotting time were not done. All
108 health workers should be trained and legislations should pave the way. Government should
109 promulgate a law that circumcision should be done only by medically qualified individuals
110 with necessary clinical skills and training. If traditional / untrained circumcision practitioners
111 must perform circumcision due to lack of personnel, they should be informed of symptoms
112 and signs (especially easily appreciated ones) that when seen, should discourage them from
113 circumcising a male child. Example that no jaundiced child should be circumcised except in a
114 medical facility. Bleeding tests are mandatory before circumcision in neonates, while
115 prophylactic vitamin K injection might be an alternative in areas where testing is not feasible.



118 Figure 1: **A)**The homogenously green coloured liver with a smooth and shiny surface. **B)** The
 119 inferior surface showing hypoplastic gallbladder. **C)** Hyper inflated lungs more marked on
 120 the left lung.



121
 122 Figure 2: A and B shows early bridging necrosis between portal tracts and parenchymal
 123 cholestasis
 124




Figure 3: Showing marked bile ductular proliferation, portal tract edema and fibrosis, and parenchymal cholestasis

REFERENCES

1. Nelson C P, Dunn R, Wan J, Wei J T. The increasing incidence of newborn circumcision: data from the nationwide inpatient sample. *Journal of Urology* 2005;173(3):978–981.
2. Weiss H A, Larke N, Halperin D, Schenker I. Complications of circumcision in male neonates, infants and children: a systematic review. *BMC Urology* 2010 10:2.
3. Aaron J K, Lane S P, Jeffrey S P. Complications of Circumcision. *The Scientific World Journal* 2011; 11: 2458–2468.
4. Sasmaz I, Antmen B, Leblebisatan G, ŞahinKaragün B, Kiliç Y, Tuncer R. Circumcision and complications in patients with haemophilia in southern part of Turkey: Çukurova experience. *Haemophilia* 2012;18:426-30.

5. Krill A J, Palmer L S, Palmer J S. Complications of Circumcision. *The Scientific World Journal* 2011; 11: 2458–2468.
6. Crawford J M, Liu C. Liver and biliary tract. In: Kumar V, Abbas A K, Fausto N, Aster J C (eds). *Robbins and Cotran Pathologic basis of diseases*, 8th Ed. Elsevier India, 2010. Pp 877-938.
7. Petersen C: Pathogenesis and treatment opportunities for biliary atresia. *Clin Liver Dis* 2006; 10:73.vi
8. Kirschner B S, Black D D. The gastrointestinal tract. In: Behrman R E, Kliegman R M (eds). *Nelson Essentials of Pediatrics*, 4th Ed. W B Saunders India 2002, Pp508-509.
9. Kotb MA. Aflatoxins in Infants with Extrahepatic Biliary Atresia. *Med. J. Cairo Univ* 2015; 83 (1): 207-210.
10. Mack CL: The pathogenesis of biliary atresia: evidence for a virus-induced autoimmune disease. *Semin Liver Dis* 2007; 27:233.
11. Desmet V J, Rosai J. Liver Non neoplastic diseases, tumors and tumorlike conditions. In: Rosai J (ed). *Rosai and Akerman's surgical pathology*, 10th Ed. Vol 1, Elsevier China, 2011. Pp 857-980.
12. Shreef K, Alhelal A. Evaluation of the use of laparoscopic-guided cholecystocholangiography and liver biopsy in definitive diagnosis of neonatal cholestatic jaundice. *Afr J Paediatr Surg* 2016;13(4):181-184.

Acknowledgement: The authors are grateful to the coroner and its officers.

Abbreviation list

Biliary atresia	BA
Kasai portoenterostomy	KPE
Hepatobiliaryiminodiacetic acid	HIDA

164

165