

A RARE CASE OF GALL STONE ILEUS WITH OBSTRUCTION OF SMALL BOWEL

Veeresh.H.M¹, Rajesh patil²

¹Senior Resident, General Surgery, Goa Medical College India

²Associate professor department of general Surgery, Goa Medical College, India.

Authors' contributions

This work was carried out in collaboration between both authors. Author VHM designed the study and wrote the first draft of the manuscript. Author HVM managed the literature searches. Author RP was the corresponding author and wrote the final draft. Both authors read and approved the final manuscript.

ABSTRACT:

Gallstone ileus is condition where small bowel is mechanically obstructed by gallstones. It's a true mechanical obstruction. It is uncommon entity which accounts for one to three percent of all small bowel obstruction². Elderly people are usually affected. X-ray, USG, CT SCAN are usual modalities of diagnosis. Mainstay of treatment is to relieve obstruction by surgery.

KEYWORDS: Small bowel obstruction, Gallstones, Gallstone ileus, , enterolithotomy

INTRODUCTION:

Gallstone ileus is true mechanical obstruction so word ileus is misnomer. This is one of the infrequent complication of gall stones which results as one or more gallstones enter INTO small bowel by cholecysto-enteric fistula. Open/laparoscopic surgery is the mainstay of treatment.

CASE REPORT:

A 48-Yr -old lady came with history of pain abdomen for three to four days, vomiting since two days and distention of the abdomen since one day. On examination 94 was pulse per minute, Blood pressure was 120/70 mm of mercury, mild abdominal distention was present. On auscultation bowel sounds were hyper, rectum was empty. Blood investigations showed raised white blood cell counts and renal function and liver function tests were normal. X –ray showed dilated small bowel loops. Ultrasound abdomen showed dilated small bowel loops and a

calculus measuring 1.5cm in gall bladder (which is pneumobilia on **Computed tomography**) and calculus in mid ileum measuring 3*3 cm (figure 1). **Contrast CT scan** abdomen was done which showed pneumobilia in gall bladder (initially thought as calculi in gall bladder on Ultrasound), and calculi in the mid ileum around 3cm*3cm*2.5cm in dimension with proximal dilated small bowel loops(figure 2). There was a cholecysto-duodenal fistula with stranding of omentum around the gallbladder(figure 3).

Figure 1. USG showing gall stone in small bowel




Figure 2. CT scan showing gallstone in small bowel




Figure 3. CT scan showing cholecysto-duodenal fistula.



Laparotomy with enterolithotomy and extraction of gall stone was carried out successfully(Figure 4 and 5). Nothing was done for Fistula between gall bladder and duodenum. Patient was extubated and shifted to recovery then ward. Recovery was uneventful. Patient didn't had any history of gallstones. At present patient is on follow up. Once patient gives **consent for surgery**, definitive surgery, i.e cholecystectomy with repair of the fistula will be done.

Figure 4. Intra operative findings dilated small bowel with gallstone in distal ileum





Figure 5: Gallstone of 3.5cm extracted from ileum



DISCUSSION:

The gallstone ileus term was coined by Dr. Erasmus Bartolini in 1654 [1]. It is a rare complication of cholelithiasis, accounting for one to three of all cases of bowel obstruction². Bouveret described a syndrome of gastric outlet obstruction caused by gallstone impaction in the duodenum after its migration through a cholecystoduodenal or choledochoduodenal fistula in 1896⁸. It is caused by one or more gallstone **getting** impacted within the lumen [3]. Adhesions forming between the inflamed gallbladder and an adjacent part of the gastrointestinal tract is the main pathogenesis. Large stones within the gallbladder subsequently cause pressure necrosis and forms a cholecysto–enteric fistula, which creates path for gallstones entry into the gut [7]. Elderly and females are usually affected. Any part of the bowel can be affected, most commonly ileum which is 60.5%, next is jejunum which is 16.1%, stomach is 14.2%, colon is 4.1% and duodenum is 3.5%. Terminal ileum and the ileocecal valve commonly affected because of the anatomically small lumen. [4,5]

Diagnosis of gall stone ileus should be considered in patients who are elderly and with history of gallstones and no history of previous surgery and tuberculosis. On plain abdominal radiography, classical signs of Rigler's triad i.e. pneumobilia, dilated intestinal loops, aberrant gallstone aid diagnosis [6].

The mainstay of treatment of gallstone ileus is a relief of mechanical obstruction of the bowel. Most of the patients with gallstone ileus requires surgical intervention. If stone is within proximal duodenum, then it can be removed by endoscope⁹. Extracorporeal shockwave lithotripsy can also be done but this method has its own limitations. Majority of patients requires surgery. Surgical options include enterotomy. Stones are removed by small incision through the bowel. Most of the surgeons favor enterolithotomy as first procedure. Cholecystectomy and repair of fistula is done in later settings.

CONCLUSION:

Even **though gallstone ileus** is rare cause of obstruction, when dealing with elderly patients without any history of surgery, gall stone ileus should be considered. **Early** surgery remains the main stay of treatment.

Consent and Ethical Approval:

Ethical committee clearance and consent for publication from patient is taken and there is no conflict of interest between authors.

References:

1. Syme RG. Management of Gallstone Ileus. Canadian Journal of Surgery. 1989; 32: 61- 64
2. Dai XZ, Li GQ, Zhang F, Wang XH, Zang CY. Gallstone ileus: case report and literature review. World J Gastroenterol 2013; 19 (33): 5586-89
3. Van Hillo M, van der JA, et al. Gallstone obstruction of the intestine: An analysis of ten patients and a review of the literature. Surgery. 1987; 101: 273-276
4. Reisner RM, Cohen JR. Gallstone ileus: a review of 1001 reported cases. Am Surg 1994; 60 (6): 441-6
5. GuptaM, Goyal S, Singal R, Goyal R, Goyal SL, Mittal A. Gallstone ileus and jejunal perforation along with gangrenous bowel in a young patient: a case report. N Am J Med Sci 2010; 2 (9): 442-3
6. Rigler LJ, Borman CN, Noble JF. Gallstone obstruction: pathogenesis and roentgen manifestations. JAMA 1941; 117: 1753-9
7. Windsor A, Heriot A. The small intestine. In: Burnand KG, Young AE, Lucas J, editors. The New Aird's Companion in Surgical Studies, 3rd edn. Amsterdam: Elsevier; 2005. pp. 623–44
8. Bouveret L. Stenose du pylore adherent a la vesicule calculeuse. Rev Med (Paris) 1896;16:1–16
9. Chen-Wang C, Shou-Chuan S, Shee-Chau L, Cheng-Hsin C: Gallstone ileus: a disease easily ignored in the elderly. Int J Gast enterology. 2008, 2 (1): 18-21.[Google Schola](#)