

CARDIO-PULMONARY RISK MARKERS IN AUTOMOBILE SPRAYERS AND SAW MILLERS

Abstract:

Other than oxygen, the air that enters the respiratory system contains several substances. These substances are derived from heterogeneous mixture of gases, liquids and particulate matter. Short term exposure to these pollutants cause deleterious effects on the cardio-respiratory systems. This study sought to investigate the effect(s) that air pollutants have on markers of cardiovascular and respiratory ailments in occupationally exposed individuals. An exposure factor study was conducted on 60 ethically recruited individuals from Enugu metropolis of Enugu state, Nigeria. They were then assigned into three groups (A-C) of 20 subjects each. With group A acting as control, groups B and C comprised of automobile spray painters and saw millers respectively. In any case, participants' respiratory and cardiovascular parameters [Forced Expiratory Volume (FEV), Forced Vital Capacity (FVC), Peak Expiratory Flow Rate (PEFR), Electrocardiogram (ECG) and haematological parameters (Fibrinogen concentrations and C-Reactive Proteins)] were assayed about 30 minutes before investigation; based on their weight height and Blood Pressure. Compared with control, Analysis of Variance (ANOVA) on collected data showed diastolic pressure (DBP) to have significantly increased with significant differences ($p < .05$) in FEV, whilst presenting insignificant values in PEFR and FVC of group C subjects. Also, a moderately strong and significant positive relationship (from Pearson Product Moment Correlation) was seen between ECG parameters (QRS and P-R) and duration of work for control (group A). Routine spirometry, haematological and ECG tests are recommended for individuals who are exposed to environmental pollutants as a result of occupational predisposition.

Keywords: Environmental pollution, Cardio-respiratory system, Lung volumes and Capacities

INTRODUCTION

Literarily known as the “apparatus of breathing”, the respiratory system supplies a critical component of life; the oxygen, and disposes a major waste product, carbon dioxide^{1&2}. The air that enters the respiratory system contains many substances other than oxygen, including natural constituents (e.g., nitrogen), as well as human and environmental contributions (e.g., fossil fuel combustion by-products). Various defence mechanisms of the respiratory system eliminate the unnatural components of air from the body and help repair any damage they may likely have. But exposure to large amounts of toxic substances or chronic exposure to lower levels can overwhelm the ability of the respiratory system to

protect and repair itself, sometimes resulting in impaired lung function. Such exposures to copious amounts of toxins can be traceable to a wide range of sources including the environment, human and industrial wastes/disposals, as well as exposures resulting from job-related or occupational activities³.

An occupational disease is any chronic ailment that occurs as a result of work or occupational activity. It is typically identified when its prevalence is more in a given body of workers than in the general population, or in other worker populations⁴. Occupational cardio-respiratory disease is the name for a collection of conditions of the cardiovascular and/or respiratory systems, which can have occupational risk factors for the particular disease. This includes acute, sub-acute and chronic diseases, which may be malignant, non-malignant or in infectious origin. Common occupational diseases of such may include occupational asthma, black lung disease(coal worker's pneumoconiosis), chronic obstructive pulmonary disease (COPD), mesothelioma, and silicosis. Infectious lung diseases can also be acquired in an occupational context. Exposure to substances like dusts, fumes, and gases such as isocyanate paint mist, silica, etc. can cause fibrotic lung diseases and impairments in cardio-respiratory functions, whereas exposure to carcinogens like asbestos and beryllium can cause lung cancer.

Significant decreases in Forced Expiratory Volume (FEV), Forced Vital Capacity (FVC) and Peak Expiratory Flow rate (PEF) show with varieties of symptoms including cough, dyspnoea, and phlegm that have been associated with exposures to gases and fumes⁵⁻⁷. This study was developed to ascertain the effect that occupational exposures have on the respiratory system of automobile sprayers and saw millers.

Aim of Study

This study was aimed at determining amongst workers in Enugu metropolis, the influence of airborne pollutants on markers of cardiovascular risk and lung function; using automobile sprayers and saw millers. Specifically, the study sought to examine:

1. The influence of occupational exposure to particulate air pollutants (saw dust and paint mist) on vital signs (systolic and diastolic blood pressures as well as heart rates).
2. The influence of occupational exposure to particulate air pollutants (saw dust and paint mist) on lung functions (Forced Expiratory Volume, Vital Capacity, and Peak Expiratory Flow rate)
3. The influence of occupational exposure to particulate air pollutants (saw dust and paint mist) on electrical activity (QRS duration, P-R interval and QTc value) of cardiac muscles.

80 **Materials and Methods**

81 **Scope of Study**

82 This study was conducted in Enugu metropolis of Enugu state, Nigeria. Enugu State is a 215
83 sq mi (556km²) area of land in Nigeria that lies approximately between Longitude 7⁰ 31' East
84 and Latitude 6⁰ 26' North (Duckworth, 1961).

85

86 **Study Design**

87 Study adopted the ex-pose facto research design. This was considered appropriate because
88 selected subjects were studied after changes in the dependent variables (Lung function,
89 electrocardiogram, fibrinogen, and C-reactive protein) have occurred due to occupational
90 exposure to air pollutants (independent variables), and thus, the investigator does not
91 manipulate the independent variable (polite et al., 2005). Sixty (60) participants were selected
92 for the study. They were randomly assigned into three groups (A-C) of 20 subjects each.
93 While group A served as control, group B comprised of automobile spray painters, leaving
94 group C exclusively for saw millers. An abridged version of the National Institute for
95 Occupational Safety and Health Asthma Identification questionnaire was administered before
96 eligible for inclusion; if proven to be non-asthmatic.

97

98 **Ethical Issues**

99 Ethical approval was obtained from the Research and Ethics committee of the University
100 of Nigeria, Enugu campus. Participants' confidentiality were maintained by using code
101 numbers instead of names, and ensuring that records were destroyed at the end of the study.
102 Subjects' informed consents were also obtained prior to study; after the procedure and
103 purpose of research had been clearly explained to them^{8&9}.

104

105 **Selection Criteria**

106 **Inclusion Criteria**

- 107 - only workers with minimum of 5 years uninterrupted work experience^{10&11}
- 108 - Only workers who have not taken any form of sedative for 72 hours (at least) before
109 investigation¹²
- 110 - Workers without any form of anti-hypertensive medication for the past 72 hours (at least)
111 before investigation

- Only paint workers who were directly physically, and routinely involved in paint mixing and vehicle spraying

Exclusion Criteria

- Individuals with any history of smoking were excluded. This was because, nicotine inhalation overtime has been found to alter cardio-pulmonary functionalities¹³
- Habitual tobacco snuffers were also excluded as it has been found to cause acute and chronic pulmonary diseases¹⁴.
- Subjects with known history of asthma or obstructive airway diseases
- Individuals whose ages were above 65years

Procedure

Questionnaires were administered to each participant to be sure if they are asthmatic or not. Next, they were required to rest for 30 minutes prior to investigation. For each subject, Lung function [Forced Expiratory Volume (FEV), Forced Vital Capacity (FVC), and Peak Expiratory Flow rate (PEF)], systolic and diastolic blood pressure, heart rates, as well as electrocardiogram readings (QRS, P-R interval and QTc values) were measured. To obtain spirometry readings, subjects were instructed to breathe in and out in upright, relaxed position. Forced expiration and inspiration readings were then read at maximum breathe outs and breathe in respectively

Analytical Approach

Obtained data were analysed using the one way analysis of variance (ANOVA). Post-hoc analysis was carried out with the Bonferroni test. Pearson Product Moment Correlation Coefficient was used to determine linear relationship between duration of exposure to air pollutant (work experience) and various parameters of lung function and electrocardiogram. All tests were one-tailed, with alpha (α) set at .05. Results are presented as means and standard deviations.

Results

Table I: Showing physical characteristics and occupational history of participants with ANOVA (n=60)

Variables	Category	X±SD	Range	F- Value	P-Value
Age (Years)	Control (A)	42.30+9.06	23-53	5.386	0.002*
	Paint (B)	40.75+9.85	17-65		
	Saw dust (C)	52.20+9.77	34-71		
Weight (kg)	Control (A)	79.60+11.38	49-105	5.064	0.003*
	Paint (B)	71.80+11.50	59-100		
	Saw dust (C)	78.65+13.48	51-102		
Height (m)	Control (A)	1.72+0.06	1-56	2.250	0.089
	Paint (B)	1.73+0.79	1.60-1.91		
	Saw dust (C)	1.69+0.08	1.58-1.90		
Height (m)	Control (A)	11.85+8.96	4.00-31.00	6.419	0.001*
	Paint (B)	19.10+9.80	4.00-41.00		
	Saw dust (C)	22.25+11.73	5.00-48.00		

*Significant at $p < .05$, $\bar{X}$ =mean, SD= standard deviation

Table II: Showing Cardiovascular Parameters of Participants

Variables	Control (Grp A) X±SD	Paint (Grp B) X±SD	Saw dust (Grp C) X±SD	f - Value	p - Value
HR	76.85±11.55	73.45±9.50	75.45±16.81	0.557	0.645
SBP	123.50±21.90	121.10±22.85	146.50±23.90	2.903	0.040*
DBP	94.85±12.58	63.30±14.86	96.20±12.91	19.073	0.001*

Keys: HR=Heart Rate, SBP=Systolic Blood Pressure, DBP=Diastolic Blood Pressure *Significant at $p < .05$

Table III: Showing Lung Function Parameters of Participants

Variables	Control (Grp A) X±SD	Paint (Grp B) X±SD	Saw dust (Grp C) X±SD	f - Value	p - Value
FVC	3.34±3.45	3.06±0.72	3.01±2.47	0.995	0.400
FEV	2.36±1.13	2.21±0.87	1.86±0.98	6.128	0.001*
PEF	3.74±1.71	4.40±1.71	3.12±1.31	5.523	0.002*

Keys: FVC=Forced Vital Capacity, FEV=Forced Expiratory Volume, PEF=Peak Expiratory Flow rate, $\bar{X}$ =mean, SD=Standard Deviation, *Significant at $p < .05$

Table IV: Showing Electrocardiogram profile of participants

Variables	Control (Grp A) X±SD	Paint (Grp B) X±SD	Saw dust (Grp C) X±SD	f - Value	p – Value
P-R Interval	154.20±22.65	160.90±27.86	168.40±21.43	1.097	0.356
QRS duration	87.40±13.37	86.70±12.44	84.60±15.32	0.859	0.466
QTc	418.05±24.44	405.35±4.13	431.50±5.75	4.521	0.054

Keys: X=mean, SD=Standard Deviation, *Significant at $p<.05$


Table V: Showing Haematological parameters of sampled participants

Variables	Control (Grp A) X±SD	Paint (Grp B) X±SD	Saw dust (Grp C) X±SD	f - Value	p – Value
FC	2.54±0.70	2.54±0.99	2.37±0.92	2.669	0.054
CRP level	3.06±0.82	2.90±1.19	2.79±1.19	0.864	0.464

Keys: X=mean, SD=Standard Deviation, FC=Fibrinogen Concentration, CRP=C-reactive protein.


*Significant at $p<.05$

Chart I: Showing relationship between participants' duration of work and Lung function parameters based on Pearson Product Moment Correlation Coefficient



Keys: FVC=Forced Vital Capacity, FEV=Forced Expiratory Volume, PEF=Peak Expiratory Flow rate, X=mean

Chart II: Showing relationship between participants' duration of work and Electrocardiogram parameters based on Pearson Product Moment Correlation Coefficient



Keys: *Correlation is Significant at the .05 level

Discussion

This study indicates that cardio-pulmonary risk markers vary in individuals according to their occupations and duration of exposure of to environmental pollutants. Our results show the following.

Systolic Blood Pressure and exposure to Paint Mist and Saw dust

Insignificant changes were seen in diastolic blood pressure of 50.5% of the population. The effects of paint mist and saw dust on systolic pressure was however high. From the etiological perspective, the insignificant effect indicates a weak influence of occupational exposure to sawdust on diastolic blood pressure. Therefore, the changes of diastolic blood pressure due to occupational exposure to sawdust in this population-based apparently healthy sample may not be clinically relevant. However, the high effect of paint mist indicates a strong influence of diastolic blood pressure.

Diastolic Blood Pressure and exposure to Paint Mist and Saw dust

Results from this study shows that changes in diastolic blood pressure (DBP) occurred in 50.5% of the population, revealing that the effect of size of saw dust ($d=0.105$) on DBP was small; even though it became high on systolic BP of paint mist and sawdust exposed individuals. Etiologically, the small effect indicates a weak influence of occupational exposure to sawdust on DBP. Therefore, the changes of DBP due to occupational exposure to sawdust in this population-based apparently healthy sample may be clinically irrelevant. However, the high effect size of paint mist indicates a strong influence of DBP. Thus, DBP

changes due to occupational exposure to paint mist and sawdust may be of great clinical relevance.

FVC, FEV, PEV and exposure to Paint Mist and Saw dust

FVC is a measure of lung volume and is usually reduced in diseases that cause the lungs to be smaller. Etiologically, the small effect is indicative of a weak influence of occupational exposure to paint mist and sawdust on measures of lung volume (FVC).

This study revealed a significant reduction in FEV and PEV of individuals occupationally exposed to sawdust. However, the reduction in FVC, FEV and PEF were insignificant as compared with control. Contrarily, there was virtually no change in FVC and FEV among those exposed to paint mist, while their PEF was higher but insignificant, compared to control. This explains the fact that various pollutants initiate different responses on lungs functions as also seen in the works of Chan-Yeung and Lee (1986). Chan-Yeung and Lee found no significant changes in FVC of their subjects who were exposed to sawdust, culminating with the fact that such inflammatory alterations did not occur in their subjects as the inert material which was used. Sawdust is probably removed by lung clearance before producing any significant modification of the bronchial mucosa.

For paint mist workers, study did not find any significant relationship between the duration of occupational exposure to paint mist and lung function parameters. However, result indicated that the magnitude of association between FEV and duration of work remained statistically significant; however, with a significant inverse relationship between duration of work and FEV. This finding was consistent with that of Baghdad, 2013¹⁴.

QT Interval and QRS complex in exposure to Paint Mist and Saw dust

Table IV Shows the Electrocardiogram profile of participants in the exercise. The study revealed that QTc interval for workers exposed to sawdust had the most prolonged value, while values obtained for control and other workers were nearly the same, with the least values observed for workers exposed to paint mist. However, all values obtained were less than 440ms, and were thus within the normal range. Also, the QRS complex had greatest values in workers exposed to cement dust, followed by those exposed to paint mist. Whereas, subjects exposed to sawdust had the least value. Invariably, the results suggest that with reference to control, QRS complex narrowed in workers exposed to paint mist, with the least recorded among those exposed to sawdust.

Previous studies on the daily variation of particulate nature of air pollution and heart rate variability in elderly subjects showed an increase in heartrate¹⁵ associated with particulate air pollution. Animal data support the concept that the autonomic nervous system may be a target for the adverse effects of air pollution. All of these findings suggest that inhaled particles may affect the balance between sympathetic and parasympathetic control of the heart, and promote a stress response that potentially leads to arrhythmias, which is one of the cardiac death triangle. This view appears corroborated by the findings of this study, suggesting that occupational exposure to airborne pollutants affect the other two components of the “Cardiac Death Triangle”

Again, a weak positive correlation was established between the QRS complex, P-R interval, QTc duration. And duration of exposure to paint mist among the relevant workers repectively. This implies that as the duration of exposure to these pollutants increased, there is a likelihood of concomitant increase in the QRS complex, P-R interval, and QT duration respectively. However, the magnitude of the relationship was insignificant.

Benefits of Study

Study will be of immense clinical value in various ways, including the provision of data that will clearly delineate the cardiovascular and pulmonary variables most affected by exposure to specific industrial pollutants, so that intervention is more precise and reflective of the defined changes in each instance. Study will also provide insight on the exact components on the lung function affected by specific pollutants across the groups. In essence, it will further provide a basis for appropriate policy and regulatory intervention, necessary to protect workers who are at risk of cardiovascular and pulmonary dysfunction due to exposure to airborne particulate pollutants.

Conclusion

Diastolic blood pressure has been shown to reduce in workers exposed to paint mist, while those exposed to sawdust recorded a significant increase in both systolic and diastolic pressures. Compared to control, Functional capacity (FCV) has also been shown to significantly reduce in workers exposed to paint mist and sawdust. Therefore, the changes in FVC due to occupational exposure to paint mist and saw dust in this population may be of huge clinical relevance.

Recommendations

Based on outcome of this study, we make the following recommendations:

- 1 Protective devices like aprons, hand gloves, long shoes dust preventive nose and mouth filters/masks be provided to exposed workers
- 2 Workers should regularly undertake periodic and routine checks for spirometry and ECG parameters in order to identify their susceptibility to hazardous particulate pollutants.

To government agencies, we make the following recommendations:

- 1 That more attention should be given to the environmental needs of exposed workers
- 2 Appropriate steps to avoid environmental disasters and adverse should must be legislated

References

1. Balogun, J.A., Obajuluwa, B.A., Olaogun, M.O., Abereoj, O.K., Oyeyem, A. Y., Adeodu, O.O., &Balogun, M.O. (1990). Anthropometric determinants of resting blood pressures and heart rates of Nigerian school children. *Annals of Tropical paediatric*; 10: 425-431.
2. Berry, P.A., Jones, S.W., Cicuttini, F.M., Wluka, A.E., &Maciewicz, R.A. (2011). Temporal relationship between serumadipokines, biomarkers of bone and cartilage turnover, and cartilage volume loss in a population with clinical knee osteoarthritis. *Arthritis Rheum*;63(3):700–7.
3. Alan, R. Schwartz., Susheel, P. Patil., Alison, M. Laffan., Vsevolod,Polotsky.,Hartmut, Schneider., and Philip, L. Smith.(2008). Obesity and Obstructive Sleep Apnea: Pathogenic Mechanisms and Therapeutic. *Approaches Proceedings of the American Thoracic Society*. Feb 15; 5(2): 185–192.
4. Escribano, B.M., Tunez, I., Requena, F., Rubio, M.D., Miguel, R.D., Montilla, P., Tovar, P. &Aguera, E.I. (2010). Effects of an aerobic training program on oxidative stress biomarkers in bulls. *VeterinariniMedicina*. 55(9): 422-428.
5. Grotle, M., Hagen, K.B., Natvig, B., Dahl, F.A., &Kvien, T.K. (2008). Obesity and osteoarthritis in knee, hip and/or hand: an epidemiological study in the general population with 10 years follow-up. *BMC Musculoskeletal Disorder*; 9:132–7.
6. Grundy, S.M. (1998).*Multifactorial causation of obesity: implications for prevention*. *Am J ClinNutr*. 67 (suppl 3): 563S–572S.

- 303 7. Grundy, S.M., Hansen, B., Smith, S.C. Jr., Cleeman, J.I., & Kahn RA (2004),
304 American Heart Association; National Heart, Lung, and Blood Institute; American
305 Diabetes Association. Clinical management of metabolic syndrome: report of the
306 American Heart Association/National Heart, Lung, and Blood Institute/American
307 Diabetes Association conference on scientific issues related to management.
308 *Circulation*.109:551–556.
- 309 8. Poirier, P., Giles, T.D., Bray, G.A., Hong, Y., Stern, J.S., Pi-Sunyer, F.X.,
310 Eckel,R.H(2006). American Heart Association; Obesity Committee of the Council on
311 Nutrition, Physical Activity, and Metabolism. Obesity and cardiovascular disease:
312 pathophysiology, evaluation, and effect of weight loss: an update of the 1997
313 American Heart Association Scientific Statement on Obesity and Heart Disease from
314 the Obesity Committee of the Council on Nutrition, Physical Activity, and
315 Metabolism. *Circulation*. 113: 898–918.
- 316 9. Popkin, B. (2003). Dynamics of the nutrition transition and its implications for the
317 developing world. *Forum Nutr*, 56:262-264.
- 318 10. Canadian Physical Activity, Fitness & Life Style Approach. (CPAFLA). (2003). CSEP-
319 Health & Fitness Program's Health-Related Appraisal and Counselling Strategy
320 ,Canadian Society for Exercise Physiology and Health Canada, Ottawa, Canada, 3rd
321 edition.
- 322 11. Caterson, I.D., Hubbard, V., Bray, G.A., Grunstein, R., Hansen, B.C., Hong, Y.,
323 Labarthe, D., Seidell, J.C., & Smith, S.C. Jr. (2004). *American Heart Association.*
324 *Prevention Conference VII: obesity, worldwide epidemic related to heart disease*
325 *and stroke: group III: worldwide comorbidities of obesity*. *Circulation*.;110: e476–
326 e483.
- 327 12. Morimoto, L.M., White, E., Chen, Z., Chlebowski, R.T., Hays, J., Kuller, L., Lopez,
328 A.M., Manson, J., Margolis, K.L., Muti, P.C., Stefanick, M.L., McTiernan, A. (2002).
329 Obesity, body size, and risk of postmenopausal breast cancer: the Women's Health
330 Initiative (United States). *Cancer Causes Control*; 13: 741–751.
- 331 13. Church, T. (2011): Exercise in obesity, metabolic syndrome, and diabetes. *Progress in*
332 *Cardiovascular Diseases* ,vol.53,no.6,pp. 412–418.
- 333 14. Manning Kelly. (2011): "Effects of Exercise Training on Fat Oxidation in Untrained
334 Overweight and obese Females." Dissertation Georgia State University,
- 335 15. Gold, A.R., Schwartz, A.R., Wise, R.A., & Smith, P.L. (1993). Pulmonary function
336 and respiratory chemo-sensitivity in moderately obese patients with sleep apnea.
337 *Chest*; 103:1325–1329.