

Original Research Article

Comparative performance of Pyrethrum (*Chrysanthemum cinerariifolium* Treviranus) extract and Cypermethrin on some field insect pests of groundnut (*Arachis hypogaea* L.) in southeastern Nigeria.

ABSTRACT

Aim: To evaluate the insecticidal efficacy of pyrethrum, *Chrysanthemum cinerariifolium*, relative to a synthetic insecticide (Cypermethrin 10 E. C) in the control of some field pests of groundnut.

Study Design: 2 x 4 factorial arrangement fitted into a Randomized Complete Block Design replicated three times.

Place and Duration of Study: Teaching and Research Farm, School of Agriculture and Agricultural Technology, Federal University of Technology, Owerri, southeastern Nigeria, between March to November, 2012.

Methodology: Groundnut seed variety, ICGV-IS 96894 (ICRISAT) was subjected to germination test to ensure viability. Seeds were planted at a spacing of 30 cm x 15 cm (220,000 plants/ha) on 24 (1 m²) beds with 1 m between furrows. Pyrethrum was extracted through a simple replicable procedure and tested at four rates (0.00, 0.25, 0.50, 0.75 g/100 ml of water). Cypermethrin 10 EC was tested at 0.00, 0.50, 1.00 and 1.50 ml/100 ml of water. Insecticides application, pest sampling and leaf damage assessments were carried out at 4, 6 and 8 WAP. Yield measurement parameters (seed weight, shell weights and pod density) were assessed.

Results: Major arthropod pests identified were; *Macrotermes bellicosus* Smeathman, *Peridontopyge* spp., *Helotrichia serrata* Fabricius, and *Oedaleus nigeriensis* Uvarov. Cypermethrin and pyrethrum applications reduced pest incidence (3.25, 2.50, 2.10 insects) and (4.09, 3.62, 3.42 insects), respectively, when compared with unsprayed plots (6.35, 6.16, 6.20 insects) at 4, 6 and 8 WAP. Insecticide type had no significant effect on the population of majority of sampled pests. Sprayed plots had less damaged leaves - 2.80 (cypermethrin) and 2.83 (pyrethrum) as against 4.52 leaves in unsprayed plots at 8 WAP. Sprayed plots also had increased fresh pod weight (0.61, 0.52 kg in cypermethrin and pyrethrum sprayed plots, respectively) as against 0.14 kg in unsprayed plots. Seed weights (0.26, 0.22 kg in cypermethrin and pyrethrum sprayed plots, respectively) were significantly distinct from the control (0.06 kg). There were no significant differences ($P = .05$) in the dry pod or shell weights based on insecticide types. The performance of the insecticides was dose related as higher rates gave better performances.

Keywords: *Arachis hypogaea*, Pyrethrum extract, Cypermethrin, *Macrotermes bellicosus*, *Peridontopyge* spp, *Helotrichia serrata*, *Oedaleus nigeriensis*.

1. INTRODUCTION

Groundnut (*Arachis hypogaea* L.), also called peanut, is a **specie** in the legume or bean family, Fabaceae. It is said to have originated from South America, in the region of Bolivia,

Argentina and Brazil and is one of the most popular commercial groups in West Africa, especially in Northern Nigeria (latitude 10°N). China is the largest producer of groundnut in the world followed by India, USA, Nigeria and Indonesia. In sub-Saharan Africa, groundnut is the 5th most widely grown crop, closely trailing behind maize, sorghum, millet and cassava. Nigeria is the largest producer in Africa, producing 30% of the Continent's total, followed by Senegal and Sudan (each with about 8%), and Ghana and Chad with about 5% each [1].

Groundnut, the third major oilseed of the world, next to soybean and cotton [2], provides a vital source of cash income and nutritious high protein and fatty food. The kernels are eaten raw, roasted, sweetened or processed into peanut butter which is rich in protein and vitamins A and B. They are also consumed as confectionary product. Groundnut oil is edible and is also used in the production of soap, cosmetics, lubricants, olein, stearin and their salts [3]. Groundnuts could prevent child malnutrition and are useful in the treatment of hemophilia, stomatitis and for the prevention of diarrhea. The meal is beneficial for growing children, pregnant women, nursing mothers [4], and is a good source of niacin, which contributes to brain health and blood flow [5].

Some insect pests such as white grubs (*Phyllophaga spp*), termites (*Macrotermes bellicosus* Smeathman), gram pod borers (*Helicoverpa armigera* Hubner), groundnut leaf miners (*Aproaerema modicella* Deventer), Aphids (*Aphis craccivora* Koch.), etc attack the plant and this may result to low yield, reduction in pod size, poor yield quality, and loss in market value.

The damages caused by these field pests have been mainly controlled with organochlorines (DDT, Thiodan etc); organophosphates (Monocrotophos, Dimethoate, etc); carbamates (Unden, Carbofuran, etc) and, more recently, pyrethroids (Permethrin, Cypermethrin, etc). Despite the efficacy of these synthetic insecticides, several adverse effects have been reportedly resulting from their misuse. These include; human poisonings, destruction of natural enemies, insecticide resistance, crop pollination problem due to honey bee losses, domestic animal poisonings, contaminated livestock products, fish and wildlife losses [6]; contamination of underground water and rivers, high persistence of the compounds, resurgence and genetic resistance of pests, adverse affect on non-target beneficial pests [7], etc. Cypermethrin 10 EC, used as a standard in this study, is a synthetic insecticide of choice in Nigeria. It is used in large scale commercial agricultural application as well as in consumer products for domestic purposes.

The ever- increasing problems associated with synthetic insecticides have synergized keen interest in the use of plant products as bioinsecticides. Botanicals are relatively safe, non-persistent, eco-friendlier and readily available [8]. Pyrethrum, a bioinsecticide, processed from the dried flower heads of *Chrysanthemum cinerariifolium* is said to have low mammalian toxicity, broad spectrum activity, is environmentally friendly and fast acting [9, 10]. Despite the worldwide acclaim for this plant, its cultivation and use in Nigeria is very limited.

Due to a dearth of information on the field pests of *A. hypogaea* in the Owerri west ecological zone of southeastern Nigeria, there is a compelling need to ascertain these field pests at different growth stages of the crop. In consideration of the debilitating effects of synthetic chemicals, it is also pertinent to screen biopesticides, especially pyrethrum, which would not be harmful to the farmers or end users of the treated produce.

2. MATERIAL AND METHODS

2.1 Site Location

The experiment was carried out at the Teaching and Research Farm of the Federal University of Technology, Owerri, Nigeria. The area is between latitude 5°25' N and longitude 7°2' E in the Tropical Rainforest Zone of southeastern Nigeria.

2.2 Collection and Preparation of Test Materials

Seeds of *Arachis hypogaea*, SAMNUT 23, were sourced from the Institute of Agricultural Research, Ahmadu Bello University, Zaria, Nigeria. This variety is also known as ICGV-IS 96894 under International Crops Research Centre for Semi-Arid Tropics (ICRISAT) nomenclature [11].

Pyrethrum, from the flower head of the plant, *Chrysanthemum cinerarifolium* Treviranus, was collected from Mawingo in Central Kenya. It was shade dried to prevent loss of active principle by sunlight, crushed into powdery form with pestle and mortar and weighed out at 0.25, 0.50 and 0.75 g. The weighed samples were mixed with 100 ml (respectively) of water in a container and allowed to settle for three hours. The solution was filtered through a fine muslin cloth and the active principle recovered and applied immediately.

Cypermethrin 10 E. C. was purchased from an agrochemical store in Owerri, Imo State, Nigeria. The insecticide was measured out with a syringe at 0.5, 1.0 and 1.5 ml, and mixed with 100 ml of water. The procedure was repeated on each spraying day.

2.3 Germination test

Germination test was carried out by random selection of 30 wholesome seeds from the groundnut seeds kept in a bag. Ten seeds each were placed inside 3 Petri-dishes that contained absorbent Whatman's No. 44 filter papers moistened with water. Daily checks were carried out for 10 days and germination rate recorded from the 4th day.

2.4 Land preparation

The site was cleared, pegged and beds made. Each bed had a dimension of 1m x 1m with furrows of 1m between. Altogether, the field contained a total of 24 beds.

2.5 Manure Application

Organic manure (cured poultry droppings) was basally applied on each of the beds and was incorporated into the soil at a rate of 2 kg per bed before planting.

2.6 Sowing and Weeding

The seeds were planted at a spacing of 30 cm x 15 cm. This gave a total of 22.0 plants per bed and 220,000 plants per hectare.

Weeding was done at 3 WAP and later at intervals with the use of hoes and hand pulling (rogueing).

2.7 Field Application of Plant Extract and Cypermethrin 10 EC

The different levels of plant extracts and Cypermethrin 10 EC, in water, were foliar- sprayed using a 250 ml hand sprayer, weekly for 3 weeks.

2.8 Pest Sampling

124 Insect pests were collected with a sweep net, cellophane bags and sample bottles. Others
125 were hand-picked using hand gloves and plastic forceps. Samplings were carried out at 4, 6
126 and 8 WAP. These were done a day before and 2 days after each spray regime and twice
127 during the pod and seed development stages. Collected insects were stored with chloroform
128 and later identified in the laboratory.

129

130 **2.9 Leaf Damage Assessment**

131

132 This was done through visual recording of the number of leaves damaged by insects. It was
133 carried out by counting the number of leaves damaged by insects on 6 selected and tagged
134 stands. Damage assessments were recorded at 4, 6 and 8 weeks after planting (WAP).

135

136 **2.10 Yield Measurement**

137

138 These were achieved through the measurements of the following parameters:

139

140 **2.10.1 Seed weight**

141

142 The seed weight was measured after the pods from each bed were harvested and the seeds
143 from dehiscent pods weighed, using Camry Emperor weighing balance (model J1111427541)
144 in kilogram (kg).

145

146 **2.10.2 Shell weight**

147

148 The shell weight measurement was carried out after drying.

149

150 **2.10.3 Pod density**

151

152 The pod density was assessed twice. First, immediately after harvest, the pods from each
153 bed were washed to remove soil and then weighed in the field. The second measurement
154 was taken after drying.

155

156 **2.11 Data Analysis**

157

158 The data collected were subjected to Analysis of Variance (ANOVA) for Randomized
159 Complete Block Design (RCBD) procedures as described by [12] and means separated
160 using Least Significant Difference at $P = .05$ level of significance.

161

162 **3. RESULTS AND DISCUSSION**

163

164 The major arthropod pests identified were termites, *Macrotermes bellicosus* Smeathman,
165 1781 (Blattodea: Termitidae -Macrotermitinae); millipedes, *Peridontopyge* spp. (Diplopoda:
166 Odontopygidae); white grub larvae, *Helotrichia serrata* Fabricius, 1781 (Coleoptera:
167 Scarabaeidae) and Nigerian grasshoppers, *Oedaleus nigeriensis* Uvarov, 1926 (Orthoptera:
168 Acrididae) (Table 1). Other arthropod pest incidences were occasional and insignificant to
169 report. The identified pests were in consonance with earlier works which implicated termites
170 (*Odontotermes* spp.), white grubs (*Holotrichia consanguinea*, *H. serrata*) and millipedes
171 (*Peridontopyge* spp.) as the major, widespread, economic pests of groundnut in Sub-
172 Saharan Africa [13, 14, 15]. Other coleopteran pests such as wireworms (*Agriotes lineatus*)
173 and false wireworms (*Gerocephalum* spp.) have been reported to be of occasional
174 importance [16].

175

176 The effect of pyrethrum, cypermethrin and their application rates on the field pests and leaf
177 damage of groundnut plants at different growth stages are also as contained in Table 1.
178 Cypermethrin and pyrethrum applications reduced total pest incidence (3.25, 2.50, 2.10
179 insects) and (4.09, 3.62, 3.42 insects), respectively, when compared with unsprayed plots
180 (6.35, 6.16, 6.20 insects) at 4, 6 and 8 WAP. Except for *O. nigeriensis* (1.75 and 0.50) and
181 *M. bellicosus* (0.50 and 1.25) at 4 and 8 WAP, respectively, insecticide type had no
182 significant effect on the population of majority of sampled pests. Sprayed plots had less
183 damaged leaves - 2.37, 2.73, 2.80 (average, 2.63)(cypermethrin) and 2.58, 2.90, 2.83
184 (average, 2.77)(pyrethrum) as against 3.17, 3.60, 4.52 (average, 3.76)(control) in unsprayed
185 plots at 4, 6 and 8 WAP, respectively. Insecticide type also had no significant effect on leaf
186 damage control.

187
188 The second rate gave the second best protection of groundnut against *M. bellicosus* (0.17,
189 0.50 and 0.50 insects) at 4, 6 and 8 WAP, respectively. The third (highest) rate recorded the
190 lowest number (0.00, 0.17 and 0.50) of sampled *M. bellicosus* across the growth stages,
191 respectively. Expectedly, the control plots were the most infested by the termites at all the
192 stages. The various application rates followed the same efficacy trend against other sampled
193 field pests.

194
195 Leaf damage result showed that the various rates of application significantly ($P = .05$)
196 reduced the number of damaged leaves sampled across the growth stages. Leaves from
197 groundnut plants sprayed with the third (highest) rate were the least damaged (1.82, 2.10
198 and 2.23 leaves) at 4, 6 and 8 WAP, respectively. Comparatively, the control plots recorded
199 the highest (3.17, 3.60 and 4.52) number of damaged leaves, respectively. It could be
200 reasonably inferred that cypermethrin and pyrethrum had basically the same statistical effect
201 on pest incidence and leaf damage of the plant.

202
203 The interactive effect of application rate with insecticide type is shown in Table 2. There was
204 only a significant interaction on the number of sampled *O. nigeriensis* all through the plant's
205 growth stages. Applying cypermethrin at the third rate (1.50 ml/ 100 ml) recorded the lowest
206 (0.33, 0.00 and 0.67) number of sampled target pests at 4, 6, and 8 WAP, respectively.

207
208 Include, as a matter of necessity, interactions between your two factors in, especially,
209 .. Tables 1 and 3, and discuss same where significant
210 .

211
212
213
214
215
216
217
218

Table 1: Main effects of application rate and insecticide type on target field pests of groundnut and leaf damage at different growth stages.

Treatment	4 WAP						6 WAP						8 WAP					
	Mb	Ps	Hs	On	Total	LD	Mb	Ps	Hs	On	Total	LD	Mb	Ps	Hs	On	Total	LD
Application Rates																		
Control	1.17	1.17	1.17	2.84	6.35	3.17	1.50	0.83	1.33	2.50	6.16	3.60	1.83	1.00	1.00	2.37	6.20	4.52
First Rate	0.33	0.50	0.50	2.17	3.50	2.57	0.83	0.33	0.33	1.60	3.09	2.88	0.67	0.17	0.33	1.34	2.51	3.02
Second Rate	0.17	0.50	0.50	1.84	3.01	2.33	0.50	0.33	0.50	0.84	2.17	2.67	0.50	0.00	0.30	0.67	1.47	2.77
Third Rate	0.00	0.33	0.33	1.17	1.83	1.82	0.17	0.00	0.00	0.50	0.67	2.10	0.50	0.00	0.17	0.17	0.84	2.23
LSD _{0.05}	0.816	NS	NS	0.204		0.396	0.758	NS	0.532	0.304		0.359	0.876	NS	NS	0.731		0.764
Insecticide Type																		
Cypermethrin	0.42	0.58	0.50	1.75	3.25	2.37	0.58	0.33	0.42	1.17	2.50	2.73	0.50	0.25	0.42	0.93	2.10	2.80
Pyrethrum	0.42	0.67	0.75	2.25	4.09	2.58	0.98	0.42	0.67	1.55	3.62	2.90	1.25	0.33	0.50	1.34	3.42	2.83
LSD _{0.05}	NS	NS	NS	0.123		NS	NS	NS	NS	NS		NS	0.619	NS	NS	NS		NS

Key:

NS: Non Significant
 Mb: *Macrotermes bellicosus* (Termites)
 Ps: *Peridontopyge spp* (Millipedes)
 Hs: *Helotrichia serrata* (White grub larvae)
 On: *Oedaleus nigeriensis* (Nigerian grasshoppers).
 LD: Leaf damage.

Rates:

Pyrethrum Extract: **Rate 1** = 0.25 g/100 ml of water; **Rate 2** = 0.50 g/100 ml of water; **Rate 3** = 0.75 g/100 ml of water
 Cypermethrin 10 EC: **Rate 1** = 0.50 ml/100 ml of water; **Rate 2** = 1.00 ml/100 ml of water; **Rate 3** = 1.50 ml/100 ml of water

Table 2: Interactive effects of application rate with insecticide type on target field pests of groundnut and leaf damage at different growth stages.

Application Rates	Insecticide Type	4 WAP							6 W AP							8 WAP						
		Mb	Ps	Hs	On	Total	LD		Mb	Ps	Hs	On	Total	LD		Mb	Ps	Hs	On	Total	LD	
Control	CYP.	1.33	1.33	1.00	2.67	6.33	3.10		1.33	0.67	1.33	2.33	5.66	3.5		1.00	1.00	1.00	2.40	5.40	4.30	
	PYM.	1.00	1.00	1.33	3.00	6.33	3.23		1.67	1.00	1.67	2.67	7.01	3.67		2.67	1.00	1.00	2.34	7.01	4.73	
First Rate	CYP.	0.00	0.33	0.33	2.00	2.66	2.47		0.67	0.33	0.50	1.33	2.83	2.83		0.33	0.00	0.33	1.00	1.66	2.93	
	PYM.	0.33	0.67	0.67	2.33	4.00	2.67		1.00	0.33	0.67	1.87	3.87	2.97		1.00	0.33	0.33	1.67	2.33	3.10	
Second Rate	CYM.	0.00	0.33	0.33	1.67	2.33	2.27		0.33	0.33	0.33	0.67	1.66	2.67		0.33	0.00	0.33	0.33	0.99	2.53	
	PYM.	0.33	0.67	0.67	2.00	3.67	2.40		0.67	0.33	0.33	1.00	2.33	2.73		0.67	0.00	0.33	1.00	2.00	3.00	
Third Rate	CYM.	0.00	0.33	0.33	0.67	1.33	1.63		0.00	0.00	0.00	0.33	0.33	2.10		0.33	0.00	0.00	0.00	0.33	1.77	
	PYM.	0.00	0.33	0.33	1.67	2.33	2.00		0.33	0.00	0.00	0.67	1.00	2.23		0.67	0.00	0.33	0.33	1.33	2.70	
LSD _{0.05}		NS	NS	NS	0.311		NS		NS	NS	NS	0.424		NS		NS	NS	NS	0.798		NS	

Key:

CYP.: Cypermethrin

PYM.: Pyrethrum

NS: Non Significant

Mb: *Macrotermes bellicosus* (Termites)

Ps: *Peridontopyge spp* (Millipedes)

Hs: *Helotrichia serrata* (White grub larvae)

On: *Oedaleus nigeriensis* (Nigerian grasshoppers).

LD: Leaf damage.

Rates:

Pyrethrum Extract: **Rate 1** = 0.25 g/100 ml of water; **Rate 2** = 0.50 g/100 ml of water; **Rate 3** = 0.75 g/100 ml of water

Cypermethrin 10 EC: **Rate 1** = 0.50 ml/100 ml of water; **Rate 2** = 1.00 ml/100 ml of water; **Rate 3** = 1.50 ml/100 ml of water

Table 3 records the effect of application rates and insecticide types on fresh pod, dry pod, seed and shell weights. The sprayed plots had increased fresh pod weight (0.61, 0.52 kg in cypermethrin and pyrethrum sprayed plots, respectively) as against 0.14 kg in unsprayed plots. Seed weights (0.26, 0.22 kg in cypermethrin and pyrethrum sprayed plots, respectively), were significantly distinct from the control (0.06 kg). There were no significant differences ($P = .05$) recorded in the dry pod or shell weights based on insecticide types.

There were significant ($P = .05$) differences with the various application rates on fresh pod and seed weights, respectively. The third rate gave the highest (0.57 and 0.30 kg) fresh pod and seed weights, respectively. The second rate was next with 0.56 and 0.27 kg fresh pod and seed weights, respectively. The performance of the insecticides was dose related as higher rates gave better performances.

Table 3: Main effects of application rate and insecticide type on fresh, dry, seed and shell weights (Kg).

Treatment	Fresh Pod Weight	Dry Pod Weight	Seed Weight	Shell Weight
Application Rates				
Control	0.14	0.31	0.06	0.11
First Rate	0.53	0.43	0.23	0.12
Second Rate	0.56	0.44	0.27	0.14
Third Rate	0.57	0.46	0.30	0.15
LSD_{0.05}	0.876	NS	0.057	NS
Insecticide Type				
Cypermethrin	0.61	0.42	0.26	0.14
Pyrethrum	0.52	0.40	0.22	0.12
LSD_{0.05}	0.079	NS	0.026	NS

Key: NS: Non Significant

Interaction effects of application rate with insecticide type on fresh pod, dry pod, seed and shell weights were not statistically significant (Table 4). However, the highest rate of cypermethrin (1.50 ml/ 100 ml) accounted for the highest fresh pod weight (0.58 kg), dry pod weight (0.47 kg), seed weight (0.32 kg) and shell weight (0.17 kg). These were, however, not statistically different from the effect of pyrethrum at the highest rate (0.75 g/ 100 ml).

Though cypermethrin performed marginally better than pyrethrum in some parameters, both treatments, however, showed no statistically significant differences ($P = .05$) in most parameters studied.

Table 4: Interactive effects of application rate with insecticide type on fresh, dry, seed and shell weights (Kg).

Application Rates	Insecticide Type	Fresh Pod Weight	Dry Pod Weight	Seed Weight	Shell Weight
Control	Cypermethrin	0.48	0.32	0.17	0.12
	Pyrethrum	0.50	0.33	0.16	0.11
1 st Rate	Cypermethrin	0.53	0.43	0.27	0.12
	Pyrethrum	0.53	0.42	0.20	0.12
2 nd Rate	Cypermethrin	0.57	0.45	0.28	0.15
	Pyrethrum	0.55	0.43	0.25	0.12
3 rd Rate	Cypermethrin	0.58	0.47	0.32	0.17
	Pyrethrum	0.57	0.45	0.28	0.13
LSD_{0.05}		NS	NS	NS	NS

Key: NS: Non Significant

The effectiveness of cypermethrin in the control of insect pests of cotton, fruits, vegetable crops and as an indoor insecticide has been noted [17]. The efficacy of the insecticide could be as a result of its action as a stomach and contact poison and its debilitating action on the nervous system [18]. It has also been reported that due to its low vapour pressure (1.3×10^{-9} mmHg at 20°C), the insecticide is absorbed as aerosols through stagnant surfaces like soil and foliage which are further exposed to atmospheric oxidation and solar radiation [19]. Though cypermethrin, which contain synthetic pyrethroids, is often dubbed as “safe as chrysanthemum flowers”, it should be noted that they are chemically engineered to be more toxic with longer breakdown times, often formulated with synergists for increased potency, thereby compromising the human body's ability to detoxify the pesticide [20]. Symptoms of cypermethrin poisoning in humans include numbness, burning, loss of bladder control, vomiting, incoordination, seizures, coma and death. The insecticide could cause reproductive toxicity, including abnormal sperm and disruption of sex hormones [21] and is very toxic to fish and other aquatic organisms [22].

The performance of pyrethrum extracts in the experiment is commendable when related to the crude extraction method employed. Its action could be attributed to the presence of pyrethrin as an active insecticidal component [23]. Pyrethrin is reported to attack the nervous systems of all insects, and when not present in lethal doses, could act as a repellent and ‘exciter’ – increasing their activity and ability to ‘flush’ out insects from their hiding places thereby increasing their exposure with the insecticide [10]. Earlier reports [24] have implicated six biologically active chemicals in pyrethrin: Pyrethrin I ($C_{21}H_{28}O_3$), Pyrethrin II ($C_{22}H_{28}O_5$), Cinerin I ($C_{20}H_{28}O_3$), Cinerin II ($C_{21}H_{28}O_5$), Jasmolin I ($C_{21}H_{30}O_3$) and Jasmolin II ($C_{22}H_{30}O_5$). Pyrethrin I, cinerin I, and jasmolin I are esters of chrysanthemic acid whereas pyrethrin II, cinerin II, and jasmolin II are esters of pyrethric acid. Chrysanthemic and pyrethric acids combine with one of three alcohols (pyrethrolone, cinerolone or jasmololone) to form the respective six active ingredients [25]. These acids are strongly lipophilic and rapidly penetrate many insects and paralyze their nervous system [26] and exert quick knockdown effects on a wide range of insect pests, causing paralysis within a few minutes

and acting as contact poison that affects their central nervous system by blocking their nervous function [27]. Despite its efficacy, the natural pyrethrin insecticides have the desirable environmental properties of being both non-toxic to mammals and non-persistent [28].

4. CONCLUSION

The efficacy of pyrethrum is statistically comparable to cypermethrin in its ability to control the field insect pests of groundnut and could serve as alternative to synthetic pesticides in the management of these pests in southeastern Nigeria.

The use of the bioinsecticide could assist in arresting the prevailing dumping of thousands of tons of poisonous pesticides on agricultural soils. It is recommended that the potential of pyrethrum extract as a protectant for field crops and its cultivation in Nigeria be further explored.

REFERENCES

1. HarvestChoiceinScience. Groundnut. 2010. Retrieved from: <http://harvestchoice.org/commodities/groundnut>
2. FAO. Food and Agricultural Organisation of the United Nations. FAO_Food outlook. 1990. Retrieved from: www.hort.purdue.edu/peanut.html
3. Himal J. Economic importance of groundnut oilseed crop. Oilseed Crop Groundnut India Meghmani Organics Limited, Ahmedabad. 2009; 14:12. Retrieved from: <http://agropedia.iitk.ac.in/?q=content/economic-importance-groundnut>
4. Kenny A, Finn K. Groundnuts (*Arachis hypogea*), women and development: Impact on nutrition and women's roles in West Africa. 2004. Retrieved from: <http://forest.mtu.edu/pcforestry/resources/studentprojects/groundnut.html>
5. Anonymous. Peanut as the world healthiest food. 2007. Retrieved from: <http://www.en.wikipedia.org/wiki/peanut>.
6. Pimentel D, Andow D, Dyson-Hudson K, Gallahan D, Jacobson S, Irish MK, Moss A, Schreiner1, Shepard M, Thompson T, Vinzant B. Environmental and Social Costs of Pesticides: A preliminary assessment. Oikos. 1980; 34: 126 -140.
7. Grzywacz D, Leavett R. Biopesticides and their role in modern pest management in West Africa. Natural Research Institute/University of Greenwich Collaboration. 2012. Retrieved from: <http://www.nri.org/news/archive/2012/20120413-biopesticides.htm>
8. Isman MB. Botanical insecticides, deterrents, and repellents in modern agriculture and an increasingly regulated world. Ann. Rev. Entomol. 2006; 51: 45–66.
9. Dayan FE, Cantrell CL, Duke SO. Natural products in crop protection. Bioorganic & Medicinal Chem. 2009; 17: 4022 – 4034.
10. Pavela R. Effectiveness of some botanical insecticides against *Spodoptera Littoralis* Boisduvala (Lepidoptera: Noctuidae), *Myzus persicae* Sulzer (Hemiptera: Aphididae) and *Tetranychus urticae* Koch (Acari: Tetranychidae). Plant Protect. Sci. 2009; 45: 161 – 167.
11. Ndjunga J, Ntare BR, Waliyar F, Echekwu CA, Kodio O, Kapran I, Diallo AT, Amadou A, Bissala HY, Da Sylva A. Early adoption of modern groundnut varieties in West Africa. Working Paper Series No. 24. Sahelian Centre, BP 12404 Niamey, Niger. International Crops Research Institute for the Semi-Arid Tropics. Patancheru, Andhra Pradesh, India. 2008; 62 pp. Retrieved from: <http://oar.icrisat.org/2360/>
12. Wahua TAT. Applied statistics for scientific studies Africa-link press, Ibadan, Nigeria. 1999; 108- 248.

- 424 13. Johnson RA, Gumel MH. Termite damage and crop loss studies in Nigeria – the
425 incidence of termite scarified groundnut pods and market samples. *Trop. Pest*
426 *Managem.* 1981; 27: 343 -350.
- 427 14. Johnson RA, Lamb RW, Wood TG. Termite damage and crop loss studies – a
428 survey of damage to groundnut. *Trop. Pest Managem.* 1981; 27: 325- 342.
- 429 15. Lynch RE, Ouedrago AP, Dicko I. Insect damage to groundnut in semi-arid tropical
430 Africa.1986; 175 -183. In *Agrometeorology of Groundnut*. Proceedings of an
431 International Symposium, 21-26 August, ICRISAT Sahelian Centre, Niamey, Niger.
432 Patancheru, A. P. 502 324, ICRISAT, India.
- 433 16. Umeh VC, Youm O and Waliyar F. Soil pests of Groundnut in Sub-saharan Africa
434 *.Intnl. J. Trop. Insect Sci.* 2001; 21: 23 - 32.
- 435 17. Takahashi N, Mikami M, Matsuda T, Miyamoto J. Photodegradation of the pyrethroid
436 insecticide Cypermethrin in water and on soil surface. *J. Pesticide Science.* 1985;
437 10: 629-642.
- 438 18. Jin H, Webster GRB. Persistence, penetration and surface availability of
439 Cypermethrin and its major degradation products in elm bark. *J. Agric. Food*
440 *Chemistry.* 1998; 46: 2851- 2857.
- 441 19. Bacci E, Calamari D, Gaggiand C, Vighi M. An approach for the prediction of
442 environment distribution and fate of Cypermethrin. *Chemosphere.* 1987; 16: 1373 –
443 1380.
- 444 20. Beyondpesticides. Pyrethroids/ Pyrethrins. Beyond Pesticides Rating: Toxic. 2014.
445 Retrieved from:
446 [http://www.beyondpesticides.org/info services/pesticidefactsheets/toxic/pyrethroid.ph](http://www.beyondpesticides.org/info services/pesticidefactsheets/toxic/pyrethroid.php)
447 [p](http://www.beyondpesticides.org/info services/pesticidefactsheets/toxic/pyrethroid.php), on October 8, 2014.
- 448 21. Elbetieha A, Daas SI, Khamas W, Darmani H. Evaluation of the toxic potential of
449 cypermethrin pesticide on some reproductive and fertility parameters in the male
450 rats. *Arch. Environ. Contam. Toxicol.* 2001; 41: 522-528.
- 451 22. ETN. Extension Toxicology Network. Cypermethrin: *Pesticide Information Profiles.*
452 1996. Retrieved from: <http://ace.orst.edu/cgi-bin/mfs/01/pips/cypermeth.htm>
- 453 23. Morris SE, Davies NW, Brown PH, Groom T. Effect of drying conditions on
454 pyrethrins content. *Industrial Crop Production.* 2006; 23(1): 9 -14.
- 455 24. Moorman R, Nguyen K. Identification and qualification of the six active compounds
456 in a pyrethrin standard. *J. Assoc. Off. Anal. Chem.* 1997; 65: 921- 926.
- 457 25. Head SW. Composition of pyrethrum extract and analysis of pyrethrins. In:
458 *Pyrethrum: The Natural Insecticide.* Cadisa JE (ed.). Academic Press. New York,
459 NY. 1973; 25-49.
- 460 26. Reigart JR, Roberts JR. *Recognition and Management of Pesticide Poisonings,*
461 U.S. Environmental Protection Agency. 1999. 401 M Street SW (7506C),
462 Washington, DC 20460. Sourced from:
463 <http://www.epa.gov/pesticides/safety/healthcare/handbook/handbook.pdf>
- 464 27. Bhat B, Menary R. Pyrethrum production in Australia: its past and present potential.
465 *J. Australian Institute of Agric. Sci.* 1984; 494: 189 -192.
- 466 28. Bullivant MJ, Pattenden G. Photodecomposition of natural pyrethrins and related
467 compounds. *Pesticide Science.* 1976; 7(3): 231 – 35.
- 468