

Increase of spinach growth through the use of larger plug cell volume and an exogenous BAP spray

J. Di Matteo¹, J. Rattin¹, A. Di Benedetto^{1-2*}

¹Faculty of Agricultural Sciences, National University of Mar del Plata, Route 226, km. 73.5 (B7620ZAA), Balcarce, Province of Buenos Aires, Argentina.

²Faculty of Agronomy, University of Buenos Aires, Avenida San Martín 4453 (C1417DSE), Buenos Aires, Argentina.

E-mail address: dibenede@agro.uba.ar (A. Di Benedetto)

Authors' contributions

This work was carried out in collaboration between all authors. All authors read and approved the final manuscript.

ABSTRACT

Aims: In spinach, transplanting has replaced direct seeding but small containers may cause root restriction effects that would be decreased post-transplant yield. Although a single hormonal regulator (BAP) spray is effective in overcoming root restriction, the mechanisms involved are unknown. The aim of this work was to analyze spinach growth changes by the use of different plug cell volumes and BAP as foliar spray.

Study design: A randomized complete block factorial design with three blocks was used.

Place and Duration of Study: The experiments were conducted at the INTA Balcarce Experimental Station campus, Argentina (37°45' S, 58°18' W and altitude 130 m) from 5th March to 24th May 2008 and repeated once from 15th March to 19th May 2009.

Methodology: We analyzed growth changes by the use of different plug cell volumes and BAP as foliar spray on dry weight accumulation and partitioning in *Spinacea oleracea* plants in two field experiments.

Results: the use of large plug cell volumes and BAP sprays at the pre-transplant stage increased plant growth, through an increase in the relative growth rate (RGR) or the rate of leaf area expansion (RLAE). We found a close direct relationship between RGR and net assimilation rate (NAR) but an inverse relationship between RGR and leaf area ratio (LAR). In addition, we found that NAR significantly increased as leaf thickness, intercellular spaces and stomatal density increased. A change in dry weight partitioning toward shoots was found as well.

Conclusion: From a grower's point of view, the use of large plug cell volume would increase spinach growth. On the other hand, a promising approach to increasing crop productivity is the use of plant growth and development regulators such as BAP.

Keywords: Leaf traits, nursery, photoassimilate partitioning, productivity, root restriction.

1. INTRODUCTION

Technological advances in transplanting have contributed to the growth of the industry by reducing costs and increasing the reliability of production. However, small containers may cause root restriction effects, i.e. a physical stress imposed on a root system which leads to a significant decrease in post-transplant growth [1, 2, 3]. Surprisingly, the root restriction associated with container volume seems to have received little consideration in the scientific literature [4].

We have previously shown that a single exogenous application of cytokinin 6-benzylaminopurine (BAP) to plants grown in small pots at the pre-transplant stage may override the shoot growth limitation due to root restriction [2, 5]. We have also shown that the pre-transplant application of BAP increases both lettuce and celery yield [1, 3] through higher shoot fresh weight accumulation and the rate of leaf area expansion (RLAE) as a result of higher relative growth rate (RGR) and the rate of leaf appearance (RLA).

In addition, it has been shown that an endogenous decrease in cytokinins levels decreased the development of the aerial part of plants [6, 7, 8, 9]). Root and shoot growth are closely coordinated, presumably by a cytokinin signaling pathway which is largely hormonal in nature with a major site of control located in the root system [10, 11]. Thus, increasing root growth may lead to a corresponding increase in the synthesis of cytokinins [12].

Spinach (*Spinacea oleracea* L.), a leaf vegetable with a high nutritive value, is of considerable economic importance. Since transplanting has almost replaced direct seeding, the objective of this work was to analyze spinach growth changes by the use of different plug cell volumes and the hormonal regulator BAP through the physiological mechanisms involved.

2. MATERIALS AND METHODS

2.1. Plant Material, Treatments and Experiments

The experiments were conducted at the INTA Balcarce Experimental Station campus, Argentina (37° 45' S, 58° 18' W and altitude 130 m) from 5th March to 24th May 2008 and repeated once from 15th March to 19th May 2009.

Spinach (cv. 'Ballet') seeds (a flowering-resistant genotype) were sown in 288- (6.18 cm³ cell⁻¹) or 200- (13.90 cm³ cell⁻¹) cell trays and were transplanted to a Typic Argiudol soil. Soil containing clay-loam-sand at 17.5, 47.5 and 35% (v/v) respectively with pH = 6.09; EC = 0.16 dS/m; N = 0.26% and P = 0.91 mmol L⁻¹. Seedlings were sprayed with BAP (6-benzylaminopurine) (SIGMA EC 214-927-5) (Sigma-Aldrich Co., St. Louis, MO, USA) solutions (0, 5, 50, 100 and 200 mg L⁻¹) either when first true leaf pairs appeared (pre-transplant application) or one week after transplant (post-transplant application) [1]. BAP was previously diluted in alcohol 80%.

Plants were grown under greenhouse facilities and were irrigated as needed (to compensate 80% relative evapo-transpiration) with high quality tap water; a weekly fertilization (Stage 2: 50 mg L⁻¹ N; Stage 3-4: 100 mg L⁻¹ N; post-transplant: 150 mg L⁻¹ N) [13] was used.

Half hourly averages air temperature was measured using a HOBO H08-001-02 data logger (Onset Computer Corporation, MA, U.S.A) protected from direct radiation by aluminum foil shades. Mean temperatures during the experiments ranged from 8.9 to 18.2°C.

2.2. Growth Evaluations

For destructive measurements, plants (10 per treatment and block) were harvested at the transplant stage and both in the middle and at the end of the experiments. The numbers of new leaves that had appeared were also recorded on the same dates and the rate of leaf appearance (RLA) was

calculated as the slope of the number of new (visible) leaves vs. time (in weeks). The fresh weight (FW) of roots and shoots was determined separately for each plant. Separate plant parts were dried at 80°C for 48 hours and weighed to obtain the dry weight of shoots (including leaf blades and stems) and roots. Leaves were oven-dried at 80°C for 48 hours. Leaf area was determined with a LI-COR 3000A automatic leaf area meter (LI-COR BioSciences, Lincoln, NE, USA).

The relative growth rate (RGR) was calculated as the slope of the regression of the natural logarithm (ln) of the whole plant on a dry weight basis vs. time (in days), while the rate of leaf area expansion (RLAE) was calculated as the slope of the regression of the ln of total leaf area vs. time (in days). Mean net assimilation rate (NAR) and leaf area ratio (LAR) were calculated according with Potter and Jones [14] as follows:

$$\text{NAR} = (k_w W_0 e^{k_w t}) / (A_0 e^{k_a t})$$

$$\text{LAR} = (k_a A_0 e^{k_a t}) / (k_w W_0 e^{k_w t})$$

where: k_w : RGR (days^{-1}); W_0 : extrapolated value of total dry weight at time zero (g); A_0 : extrapolated value of leaf area at time zero (cm^2); k_a : RLAE (days^{-1}); t : time (in days) at the midpoint of the experimental period and e : base of natural logarithms.

The specific leaf area on a FW basis (SLA_{FW}) was calculated as the ratio between area of the new individual leaf and leaf FW. The leaf weight rate (LWR) values measured biomass allocated to leaves vs. biomass allocated to other plant parts. The allometric coefficients between roots and shoots were calculated as the slope of the straight-line regression of the ln of root dry weight vs. the ln of shoot dry weight.

Tissue from the middle region of the lamina was fixed in formalin-acetic-alcohol. The size and densities of epidermal stomata size were recorded on this tissues discolored with ethyl alcohol (96%) at boiling temperature. Leaf thickness was determined from leaf lamina tissues embedded in paraffin, sectioned at 20 μm on a rotary microtome. Leaf samples were stained with safranin-crystal violet-fast green. Data are the mean of three leaves per treatment and block from ten leaf cross-sections per leaf. An image analysis system (Image Pro Express v 6.0, Media Cybernetics, USA) facilitated quantitative anatomical measurements.

2.3. Statistical Analysis


We used a randomized complete factorial design with three blocks of each treatment: Cell size, BAP concentration (BAP_c) and BAP application time (BAP_{at}). Since we found no significant differences between the two experiments, we considered them together. Data were subjected to two way analysis of variance and means were separated by Tukey's test ($P < 0.05$); STATISTICA 8 (StatSoft) software was used. Straight-line regression slopes were tested for parallelism (test for equal slope) [15].

3.1. RESULTS

3.1. Leaf area and biomass accumulation

The shoot fresh weight at the end of the experiments was higher for control plants grown in 200-cell trays than for those grown in 288-cell ones and for those sprayed at the pre-transplant stage with 5 to 100 mg L^{-1} BAP. The higher the BAP dose, the higher the shoot fresh weight. On the other hand, a high (200 mg L^{-1}) pre-transplant or post-transplant dose decreased aerial fresh weight (plants grown in 200-cell trays) or showed little changes (Figure 1).

Spinach plants from 200-cell trays showed higher total leaf area than those from 288-cell trays at the end of the experiment. In general, total leaf area in control plants was increased by a single BAP spray, but was highest when BAP was sprayed at the pre-transplant stage. The higher leaf area was the result of an increase in both the individual leaf area and RLA (Table 1).



ANOVA	
Significantly	
PCV	***
BAP _c	***
BAP _{at}	***
PCV x BAP _c	***
PCV x BAP _{at}	***
PCV x BAP _c x BAP _{at}	***

Significance *** 0.001

Fig. 1. The effects of two plug cell volume (288- and 200-cells tray⁻¹) (PCV) and a pre-transplant or pos-transplant BAP sprays (5, 50,100 or 200 mg L⁻¹) on spinach shoot fresh weight at the end of the experiment (n = 6). Both, standard errors and the analysis of variance (ANOVA) are indicated.

RLAE was higher in control plants from 200-cell than from 288-cell trays and was significantly increased with a BAP spray at almost all doses tested at both pre- and post-transplant stages. RGR showed similar results except when BAP was applied at the post-transplant stage in plants grown in 200-cell trays. NAR was higher in plants grown in 200-cell trays than in those grown in 288-cell ones. A single BAP spray (at all doses) increased NAR plants grown in 288-cell trays both at the pre- and at the post-transplant stage; BAP failed to increase NAR at the post-transplant stage in 200-cell grown plants. Results for LAR were opposite. Specific leaf area (SLA) was higher in control plants grown in 288-cell trays than in those grown in 200-cell ones, but in both cases was significantly decreased by a BAP spray. Leaf weight ratio (LWR) increased when plants were sprayed with BAP but even was higher in plants grown in 200-cell plug trays than in those grown in 288-cell ones (Table 2).

Table 1. Changes in the total leaf area, the individual leaf area and the rate of leaf appearance (RLA) at the end of the experiment in spinach plants grown at two plug cell volume (PCV) and sprayed with different concentrations of BAP solutions at the pre- or post-transplant stage (data are the mean of two experiment) ($n = 6$) $\pm$ SE. Different lower-case letters indicate statistically significant differences ($P = .05$) between BAP treatments for each cell volume while different capital letters indicate statistically significant differences ($P = .05$) for each BAP treatment between different cell volumes. The probability of the RLA slope being zero was $P = .001$. The significance of interactions (ANOVA) has been indicated as well.

PCV	Total leaf area (cm ² plant ⁻¹)	Individual leaf area (cm ² leaf ⁻¹)	RLA (leaves week ⁻¹)
288 cells tray⁻¹			
Control	499.37 ^{cB}	34.07 ^{cB}	1.122 ^{cB}
Pre-5	598.74 ^{bB}	42.71 ^{bB}	1.262 ^{aB}
Pre-50	681.25 ^{aB}	45.18 ^{bA}	1.233 ^{aB}
Pre-100	670.18 ^{aB}	53.72 ^{aA}	1.292 ^{aB}
Pre-200	656.61 ^{aA}	41.50 ^{bB}	1.142 ^{bB}
Post-5	595.80 ^{bA}	41.48 ^{bA}	1.108 ^{bB}
Post-50	598.93 ^{bA}	42.43 ^{bA}	1.157 ^{bB}
Post-100	610.34 ^{bA}	38.90 ^{bA}	1.131 ^{bB}
Post-200	638.42 ^{bA}	43.02 ^{bA}	1.147 ^{bB}
200 cells tray⁻¹			
Control	562.04 ^{CA}	39.21 ^{CA}	1.219 ^{CA}
Pre-5	743.23 ^{aA}	50.94 ^{aA}	1.358 ^{aA}
Pre-50	768.69 ^{aA}	46.04 ^{aA}	1.384 ^{aA}
Pre-100	773.18 ^{aA}	51.30 ^{aA}	1.350 ^{aA}
Pre-200	649.18 ^{bA}	49.12 ^{aA}	1.202 ^{bA}
Post-5	604.75 ^{bA}	45.76 ^{bA}	1.272 ^{bA}
Post-50	642.65 ^{bA}	48.80 ^{bA}	1.224 ^{CA}
Post-100	639.43 ^{bA}	45.58 ^{bA}	1.212 ^{CA}
Post-200	654.23 ^{bA}	42.58 ^{bA}	1.225 ^{CA}

ANOVA			
Significantly			
	Total leaf area	Individual leaf area	RLA
PCV	***	***	***
BAP _c	***	***	***
BAP _{at}	*	*	***
PCV x BAP _c	***	***	***
PCV x BAP _{at}	**	**	***
PCV x BAP _c x BAP _{at}	**	**	***

Significance *** 0.001 ** 0.01 * 0.05

Table 2. Changes in the rate of leaf area expansion (RLAE), the relative growth rate (RGR), the

net assimilation rate (NAR), the leaf area ratio (LAR), the specific leaf area (SLA) and the leaf weight rate (LWR) in spinach plants grown at two plug cell volume (PCV) and sprayed with different concentrations of BAP solutions at the pre- or post-transplant stage (n = 6) $\pm$ SE. Different lower-case letters indicate statistically significant differences between BAP treatments for each cell volume while different capital letters indicate statistically significant differences for each BAP treatment between different cell volumes at $P = .05$ by Tukey's test (SLA and LWR) or according the test for equal slope (testing for parallelism) (RLAE, RGR, NAR and LAR); the probability of these slopes being zero were $P = .001$.

PCV	RLAE (cm ² cm ⁻² day ⁻¹)	RGR (g g ⁻¹ day ⁻¹)	NAR (g cm ⁻² day ⁻¹) x 10 ⁵	LAR (cm ² g ⁻¹)	SLA (cm ² g ⁻¹)	LWR (g g ⁻¹)
288 cells tray⁻¹						
Control	0.028 ^{bB}	0.076 ^{bB}	4.65 ^{bB}	1856.20 ^{aA}	3.53 ^{aA}	1.389 ^{dB}
Pre-5	0.036 ^{aA}	0.084 ^{aB}	5.97 ^{aB}	1540.71 ^{bA}	3.11 ^{bA}	1.532 ^{cB}
Pre-50	0.036 ^{aA}	0.082 ^{aB}	5.96 ^{aB}	1535.37 ^{bA}	2.61 ^{bA}	2.822 ^{bA}
Pre-100	0.037 ^{aA}	0.083 ^{aB}	5.71 ^{aB}	1511.63 ^{bA}	2.86 ^{bA}	2.531 ^{bB}
Pre-200	0.038 ^{aA}	0.075 ^{bB}	5.78 ^{aB}	1556.20 ^{bA}	2.76 ^{bA}	2.838 ^{aB}
Post-5	0.037 ^{aA}	0.074 ^{bB}	5.49 ^{aB}	1594.26 ^{bA}	2.73 ^{bA}	1.450 ^{cA}
Post-50	0.037 ^{aA}	0.071 ^{bB}	5.37 ^{aB}	1577.40 ^{bA}	2.61 ^{bA}	1.542 ^{cA}
Post-100	0.036 ^{aA}	0.071 ^{bB}	5.27 ^{aB}	1548.86 ^{bA}	2.77 ^{bA}	1.790 ^{cA}
Post-200	0.035 ^{aA}	0.070 ^{bB}	5.26 ^{aB}	1511.29 ^{aB}	2.87 ^{bA}	1.587 ^{cA}
200 cells tray⁻¹						
Control	0.032 ^{bA}	0.082 ^{bA}	7.15 ^{bA}	1248.38 ^{aB}	3.13 ^{aB}	1.590 ^{dA}
Pre-5	0.038 ^{aA}	0.092 ^{aA}	8.16 ^{aA}	906.87 ^{bB}	2.66 ^{bB}	2.005 ^{cA}
Pre-50	0.040 ^{aA}	0.091 ^{aA}	7.90 ^{aA}	1007.95 ^{bB}	2.64 ^{bA}	2.507 ^{bA}
Pre-100	0.041 ^{aA}	0.092 ^{aA}	8.49 ^{aA}	977.50 ^{bB}	2.84 ^{bA}	3.021 ^{aA}
Pre-200	0.035 ^{aA}	0.084 ^{bA}	6.94 ^{bA}	1224.15 ^{aB}	2.80 ^{bA}	3.008 ^{aA}
Post-5	0.036 ^{aA}	0.080 ^{bA}	6.86 ^{bA}	1224.15 ^{aB}	2.88 ^{bA}	1.754 ^{cA}
Post-50	0.036 ^{aA}	0.080 ^{bA}	6.85 ^{bA}	1182.04 ^{aB}	2.71 ^{bA}	1.776 ^{cA}
Post-100	0.034 ^{bA}	0.079 ^{bA}	7.20 ^{bA}	1124.39 ^{aB}	2.42 ^{bB}	1.729 ^{cA}
Post-200	0.033 ^{bA}	0.080 ^{bA}	6.67 ^{bA}	1170.28 ^{aB}	2.57 ^{bB}	1.753 ^{cA}

ANOVA						
Significantly						
	RLAE	RGR	NAR	LAR	SLA	LWR
PCV	***	***	***	***	***	***
BAP _c	***	***	***	***	***	**
BAP _{at}	ns	**	**	**	***	**
PCV x BAP _c	***	***	***	***	***	**
PCV x BAP _{at}	**	**	**	**	***	**
PCV x BAP _c x BAP _{at}	**	**	**	**	***	**

Significance *** 0.001 ** 0.01 'ns' No significant

3.2. Dry weight partitioning

The allometries from Table 3 show that controls plants grown in 288- and 200-cell trays partitioned a similar photo-assimilate proportion to roots and that a single BAP application significantly changed photo-assimilate partitioning towards shoots.

Table 3. Changes in the allometric relationships between shoots and roots for spinach plants grown at two plug cell volume (PCV) and sprayed with different BAP solutions at the pre- or post-transplant stage using a straight-line regression analysis between the natural logarithm of the roots dry weight and the natural logarithm to the shoot dry weight ($\ln \text{Root dry weight} = \alpha + \beta \times \ln \text{Shoot dry weight}$). Mean values ($n=6$) $\pm$ SE. The coefficients of determination (r^2) are indicated. Different lower-case letters indicate statistically significant differences between BAP treatments for each cell volume while different capital letters indicate statistically significant differences for each BAP treatment between different cell volumes according the test for equal slope (testing for parallelism). The probability of the slope being zero was $P = .001$.

PCV	α	β	r^2
288 cells tray⁻¹			
Control	-1.77	0.69 ^{aA}	0.675
Pre-5	-1.97	0.58 ^{bA}	0.788
Pre-50	-2.25	0.55 ^{bA}	0.807
Pre-100	-2.17	0.60 ^{bA}	0.748
Pre-200	-2.33	0.59 ^{bA}	0.661
Post-5	-1.77	0.60 ^{bA}	0.782
Post-50	-2.04	0.53 ^{bA}	0.722
Post-100	-1.92	0.60 ^{bA}	0.836
Post-200	-1.76	0.61 ^{bA}	0.822
200 cells tray⁻¹			
Control	-1.65	0.65 ^{aA}	0.771
Pre-5	-1.76	0.50 ^{bB}	0.730
Pre-50	-1.93	0.60 ^{bA}	0.663
Pre-100	-2.01	0.55 ^{bB}	0.750
Pre-200	-1.91	0.55 ^{bA}	0.821
Post-5	-1.50	0.60 ^{bA}	0.854
Post-50	-1.70	0.56 ^{bA}	0.841
Post-100	-1.78	0.58 ^{bA}	0.841
Post-200	-1.81	0.52 ^{bB}	0.836

ANOVA	
Significantly	
PCV	***
BAP _c	***
BAP _{at}	**
PCV x BAP _c	**
PCV x BAP _{at}	**
PCV x BAP _c x BAP _{at}	**

Significance *** 0.001 ** 0.01

218

219 **3.3. Anatomical traits**

220 Leaf thickness was higher in plants grown in 200-cells trays than in those grown in 288-cell ones.
 221 Leaf thickness was significantly increased by a BAP spray, but the response was different according
 222 to the BAP dose applied and application time; the same response was found when size and density
 223 of stomata were recorded (Table 4).

224

225 **Table 4. Changes in leaf thickness, stomata density and stomata size in the last expanded**
 226 **leaf at the end of the experiments in spinach plants grown at two plug cell volume (PCV) and**
 227 **sprayed with different concentrations of BAP solutions at the pre- or post-transplant stage.**
 228 **Mean values (n=6) $\pm$ SE. Different lower-case letters indicate statistically significant**
 229 **differences ($P = .05$) between BAP treatments for each cell volume while different capital**
 230 **letters indicate statistically significant differences ($P = .05$) by Tukey's test for each BAP**
 231 **treatment between different cell volumes.**

232

	Leaf thickness ($\mu\text{m leaf}^{-1}$)	Stomata density (stomata mm^{-2})	Stomata size ($\mu\text{m stomata}^{-1}$)
288 cells tray⁻¹			
Control	418.18 ^{cB}	55.73 ^{cB}	904.80 ^{aB}
Pre-5	535.15 ^{aB}	87.58 ^{aB}	723.76 ^{bB}
Pre-50	542.42 ^{aB}	79.62 ^{aB}	837.33 ^{bA}
Pre-100	538.18 ^{aB}	79.85 ^{aB}	842.00 ^{bA}
Pre-200	445.45 ^{bB}	79.78 ^{aB}	659.40 ^{cB}
Post-5	433.33 ^{bB}	85.80 ^{aB}	586.13 ^{cA}
Post-50	448.48 ^{bB}	79.83 ^{aB}	628.00 ^{cA}
Post-100	425.76 ^{cB}	85.54 ^{aB}	732.67 ^{bA}
Post-200	415.15 ^{cB}	71.66 ^{bB}	779.20 ^{bA}
200 cells tray⁻¹			
Control	469.70 ^{dA}	63.69 ^{cA}	1014.10 ^{aA}
Pre-5	562.12 ^{aA}	87.58 ^{aA}	921.07 ^{bA}
Pre-50	566.97 ^{aA}	111.46 ^{aA}	805.93 ^{cA}
Pre-100	568.18 ^{aA}	108.50 ^{aA}	853.67 ^{cA}
Pre-200	513.64 ^{bA}	87.80 ^{bA}	879.20 ^{bA}
Post-5	483.33 ^{cA}	99.62 ^{aA}	565.20 ^{eA}
Post-50	457.88 ^{cA}	87.58 ^{bA}	733.70 ^{cA}
Post-100	475.76 ^{cA}	95.54 ^{aA}	659.40 ^{eA}
Post-200	439.39 ^{dA}	88.51 ^{bA}	742.30 ^{cA}

233

ANOVA			
Significantly			
	Leaf thickness	Stomata density	Stomata size
PCV	***	***	***
BAP _c	***	***	***
BAP _{at}	***	***	***
PCV x BAP _c	***	***	***
PCV x BAP _{at}	***	***	***
PCV x BAP _c x BAP _{at}	***	***	***

234

3.4. Growth rates and anatomical changes ratios

When plotting the data from all treatments, we found a close direct relationship ($r^2 = 0.822$) between RGR and NAR (Figure 2a) and an inverse relationship between RGR and LAR ($r^2 = 0.761$) (Figure 2b). Nevertheless, control plants grown in 200-cell trays showed higher NAR and lower LAR values than those grown in 288-cell trays.




Fig. 2. The net assimilation rate (NAR) (a) and the leaf area ratio (LAR) (b) related to the relative growth rate (RGR). The straight-line regressions were: $\text{NAR} = 192.11 \text{ RGR} - 10.07$ ($r^2 = 0.822$) and $\text{LAR} = -52,175.00 \text{ RGR} + 5,856.20$ ($r^2 = 0.761$). $\diamond$: 288-grown controls; $\triangle$: 200-grown controls.

On the other hand, NAR significantly increased according to an increase in leaf thickness ($r^2 = 0.694$) (Figure 3a), intercellular space proportion ($r^2 = 0.727$) (Figure 3b) and stomata density ($r^2 = 0.495$) (Figure 3c). NAR relationships were higher in control plants grown in 200-cell trays than in those grown in 288-cells ones.




Fig. 3. The net assimilation rate (NAR) related to leaf thickness (a), intercellular spaces (b) and stomata density (c). The straight-line regressions were: $\text{NAR} = 0.015 \text{ leaf thickness} - 0.95$ ($r^2 = 0.694$), $\text{NAR} = 0.110 \text{ intercellular spaces} + 0.40$ ($r^2 = 0.727$) and $\text{NAR} = 0.061 \text{ stomata density} + 1.22$ ($r^2 = 0.495$) respectively. $\diamond$: 288-grown controls; $\triangle$: 200-grown controls.

4. DISCUSSION

Leaf expansion and synthesis of photo-assimilates are essential for plant growth. Our results showed that a higher total leaf area in spinach (Table 1) as a result of higher RLAE (Table 2) was closely related to plug cell volume and both BAP spray concentration and BAP application time (pre- or post-transplant stage). Previous reports have shown similar increases in total leaf area in other leafy vegetables such as lettuce and celery [1, 3], but the fact that this result would be associated with a higher individual leaf area and RLA (Table 1) has not been previously published in vegetables. We have recently suggested that an increase in plug cell size or exogenous BAP would increase leaf size or leaf appearance as an expression of meristem growth and phylochron decrease respectively in ornamental foliage plants [16, 5, 17]. Cytokinins control growth mainly through the regulation of cell division and differentiation in the stem apical meristem [18]. Similarly, the size of the apical

meristem is in turn related to the rate of leaf appearance, as shown in experiments with cytokinins deficient *Arabidopsis* mutants [19].

The productivity of most typical field grown leafy vegetables is closely associated with a high both fresh and dry weight per plant. The possibility to reach a higher plant yield was strongly restricted by the plug cell volume used, perhaps related to insufficient cytokinins supply [1, 3]. Figure 1 shows that the higher spinach fresh weight was found in control plants transplanted from 200-cell trays and those which were sprayed with a single 100 mg L⁻¹ BAP at the pre-transplant stage. These results are in agreement with the fact that availability of hormones synthesized in the root apex and reallocated to shoots would be reduced when the vertical cell root growth is impeded by the container base [20].

Since there were no significant differences in dry weight content for control or treated plants (data not shown), it is possible to describe the photo-assimilate acquisition and partition rates on a dry weight base. Thus, RGR increased in plants grown in 200-cell trays and when a 5 to 100 mg L⁻¹ BAP spray was applied at the pre-transplant stage (Table 2). RGR can be described as the product between NAR or 'physiological component', and LAR or 'morphological component'. In our experiment, the higher spinach RGR would be explained by an increase in NAR (Table II) and by the close positive relationship between NAR and RGR ($r^2 = 0.822$) (Figure 2a). An increased efficiency of carbon fixation would be associated with higher NAR values, since this variable is largely the net result of carbon gain and carbon loss [21].

Both direct additions of cytokinins [22] and over expression in transgenic plants [23] have been reported to increase leaf thickness, which may further enhance carbon assimilation per unit leaf area. On the other hand, fresh biomass per unit leaf area is well correlated to leaf thickness [24]. Resistances encountered in the mesophyll from the intercellular spaces to the chloroplasts may relevantly decrease CO₂ concentration in the chloroplast and may contribute to limiting photosynthesis [25]. Aalto and Juurola [26] showed that large gradients in CO₂ concentrations can be observed in the stomatal capillary tube and in the mesophyll cells and that the concentration changes less than 1% across the airspaces of the mesophyll. This suggests that the gas phase diffusion is fast enough to provide a plentiful CO₂ supply, even in the narrowest passages between the cells [27]. Airspace effects may be larger in some species. It is profitable to have a large amount of absorbing surface because CO₂ transport is fast in air and slow in solution. The concentration decreased faster in the mesophyll cells than in the airspaces, because the diffusion coefficient is four orders of magnitude smaller in liquids than in gases. Due to the fast gas phase diffusion and equal distance from cell walls, chloroplasts in different parts of the leaf obtain almost equal quantities of CO₂ for photosynthesis. SLA and internal leaf anatomy often affect net gas exchange because of their effects on internal CO₂ conductance to the site of carboxylation, internal shading and competition for CO₂ among carboxylation sites. On the other hand, Magyar-Tábori et al. [28] have indicated that cytokinins can influence the leaf structure. Although a negative relationship between LAR and RGR ($r^2 = 0.761$) (Figure 2b) was found in the present study, there were statistically significant changes in the 'morphological component' of growth (NAR) through different leaf attributes such as leaf thickness, stomata density and stomata size related to cell size or BAP concentration and time of application (pre- or post-transplant) (Table 4). We also found a change in specific leaf area (SLA) on a fresh weight basis (Table 2). A significant proportion of NAR values would be explained by the positive relationship between leaf thickness ($r^2 = 0.694$) (Figure 3a), intercellular spaces ($r^2 = 0.727$) (Figure 3b) and stomata density ($r^2 = 0.495$) (Figure 3c). The BAP-mediated increase in NAR might be the consequence of different possible effects of cytokinin on the efficiency of carbon fixation. Another possibility would be that BAP could promote leaf blade thickening, thus leading to an increase in the levels of photosynthetic machinery per unit leaf area [29]. Mediavilla et al. [30] and Gandolfo et al. [31] showed that changes in leaf dry mass per unit area are caused by variations in internal anatomy and leaf tissue density, and are not simply a consequence of changes in leaf thickness. Our results are in agreement with these previous findings (Table 4; Figure 3).

Photo-assimilates from leaves are used for root and stem growth. In addition, the size of the different plant sinks determines the photo-assimilate partitioning among the plant organs. Our results showed that a decrease in root restriction through the use of larger plug cell and an exogenous BAP application increased dry weight partitioning to shoots as indicated by both the allometric relationships between roots and shoots (Table 3) and the leaf weight rate (LWR), which measures the allocation of biomass to leaves vs. that to other plant parts [21] (Table 2). However, complementary information about sugar contents in different organs of the plant is needed.

Changes in the allometric slopes reflect variations in carbon partitioning [32], and carbohydrate partitioning between competing sites is fundamental to plant growth; especially because plants are capable of modifying their resource allocation to favor the growth of their growing parts [33]. Plant tissues and organs rich in cytokinins are known to attract assimilate translocation. Ron'zhina (2004) [34] showed that the sink-promoting effect of cytokinins in detached leaves was related to the source-to-sink transformation of the portion of the mature leaf and manifests itself by changing the direction of the phloem assimilate translocation. Incorporation of cytokinin-producing bacteria into the root zone of lettuce plants has been shown to double the speed of accumulation of shoot biomass at the normal level of water supply [35].

5. CONCLUSION

In summary, spinach yield is limited by root restriction during the plug tray propagation stages mainly associated with a decrease in dry weight accumulation. The use of larger plug cells and BAP sprays at the pre-transplant stage increased plant productivity (expressed as RGR or RLAE) through an increase in NAR as a result of higher leaf thickness, stomata size, stomata density and intercellular spaces and a change in dry weight partitioning toward shoots.

From a grower's point of view, the use of large plug cell volume would increase yield. On the other hand, a promising approach to increasing crop productivity is the use of plant growth and development regulators such as BAP. This compound, even at very low doses, is capable of changing the intensity and direction of plant physiological processes. Because BAP is not phytotoxic, it can be used in the nursery plant industry on a large scale.

6. ACKNOWLEDGEMENTS

This work was supported by the University of Buenos Aires Science Program under Grant G054 and W193 and the University of Mar del Plata Science Program under AGR 317/010 and AGR 379/12.

REFERENCES

1. Araki A, Rattin J, Di Benedetto A, Mirave P. Temperature and cytokinin relationships on lettuce (*Lactuca sativa* L.) and celery (*Apium graveolens* L.) nursery growth and yield. Int. J. Agric. Res. 2007; 2(8):725-730.
2. Di Benedetto A. Root restriction and post-transplant effects for bedding pot plants. In: Aquino, J.C., editor. Ornamental Plants: Types, Cultivation and Nutrition. Nova Science Publishers, Inc. NY, USA. pp. 47-79. 2011.
3. Coro M, Araki A, Rattin J, Miravé P, Di Benedetto A. Lettuce and celery responses to both BAP and PBZ related to the plug cell volume. Amer. J. Exp. Agric. 2014; 4(10): 1103-1119.
4. Poorter H, Bühler J, van Dusschoten D, Climent J, Postma JA. Pot size matters: a meta-analysis of the effects of rooting volume on plant growth. Funct. Plant Biol. 2012; 39(11): 839-850.
5. Di Benedetto A, Pagani A. Dry weight accumulation in the *Impatiens walleriana* pot plant in responses to different pre-transplant plug cell volume. Eur. J. Hortic. Sci. 2013; 78(2): 76-85.

6. Werner T, Motyka V, Laucou V, Smets R., Van Onckelen H. Cytokinin-deficient transgenic *Arabidopsis* plants show multiple developmental alterations indicating opposite functions of cytokinins in the regulation of shoot and root meristem activity. *Plant Cell*. 2003; 15(11): 2532-2550.
7. Shani E, Yanai O, Ori N. The role of hormones in shoot apical meristem function. *Curr. Op. Plant Biol.*, 2006; 9(5): 484-489.
8. Kurepin LV, Emery RJN, Pharis RP, Reid DM. Uncoupling light quality from light irradiance effects in *Helianthus annuus* shoots: putative roles for plant hormones in leaf and internode growth. *J. Exp. Bot.* 2007; 58(8): 2145-2157.
9. Kyoizuka J. Control of shoot and root meristem function by cytokinin. *Curr. Op. Plant Biol.* 2007; 10: 442-446.
10. Dodd IC. Root-to shoot signalling: Assessing the roles of 'up' in the up and down world of long-distance signalling *in planta*. *Plant Soil* 2005; 274(1-2): 251-270.
11. Hirose N, Takei K, Kuroha T, Kamada-Nobusada T, Hayashi H, Sakakibara H. Regulation of cytokinin biosynthesis, compartmentalization and translocation. *J. Exp. Bot.* 2008; 59(1): 75-83.
12. O'Hare TJ, Turnbull CGN. Root growth, cytokinin and shoot dormancy in lychee (*Litchi chinensis* Sonn.). *Sci. Hortic.-Amsterdam* 2004; 102(2): 257-266.
13. Styer RC, Koranski DS. Plug and Transplant Production. A grower's Guide. Ball Publishing, Batavia, Illinois, USA. 1997.
14. Potter J, Jones JW. Leaf area partitioning as an important factor in growth. *Plant Physiol.* 1977; 59 (1): 10-14.
15. Motulsky H, Christopoulos A. Fitting models to biological data using lineal and nonlinear regression. A practical guide to curve fitting. GraphPad Software Inc., San Diego, CA, USA. 351 pp. 2003.
16. Di Benedetto A, Galmarini C, Tognetti J. Changes in leaf size and in the rate of leaf production contribute to cytokinin-mediated growth promotion in *Epipremnum aureum* L. cuttings. *J. Hortic. Sci. Biotechnol.* 2013; 88(2): 179-186.
17. De Lojo J, Di Benedetto A. Biomass accumulation and leaf shape can be modulated by an exogenous BAP spray in the ornamental foliage plant *Monstera deliciosa* (Liebm.). *J. Hortic. Sci. Biotechnol.* 2014; 89(2): 136-140.
18. Jaillais Y, Chory J. Unraveling the paradoxes of plant hormone signaling integration. *Nat. Structur. Mol. Biol.* 2010; 17(6): 642-645.
19. Francis D, Halford NG. Nutrient sensing in plant meristems. *Plant Mol. Biol.* 2006; 60(6): 981-993.
20. Di Benedetto A, Molinari J, Rattin J. The effect of transplant in sweet maize (*Zea mays* L.). II. Container root restriction. *Int. J. Agric. Res.* 2006; 1(6): 555-563.
21. Shipley B. Net assimilation rate, specific leaf area and leaf mass ratio: which is most closely correlated with relative growth rate? A meta-analysis. *Funct. Ecol.* 2006; 20(4): 565-574.
22. Bosselaers JP. Cytokinin effects on leaf architecture in *Phaseolus vulgaris* L. *J Exp. Bot.* 1983; 34(145): 1007-1017.
23. Rupp HM, Frank M, Werner T, Strnad M, Schmülling T. Increased steady state mRNA levels of the STM and KNAT1 homeobox genes in cytokinin overproducing *Arabidopsis thaliana* indicate a role for cytokinins in the shoot apical meristem. *Plant J.* 1999; 18(5): 557-563.
24. Milla R, Reich PB, Niinemets Ü, Castro-Diaz P. Environmental and developmental controls on specific leaf area are little modified by leaf allometry. *Funct. Ecol.* 2008; 22(4): 565-576.
25. Flexas J, Barbour MM, Brendel O, Cabrera HM, Carriqui M, Diaz-Espejo A, Douthe C, Dreyer E, Ferrio JP, Gago J, Galle A, Galmes J, Kodama N, Medrano H, Niinemets Ü, Peguero-Pina JJ, Pou A, Ribas-Carbo M, Tomas M, Tosens T, Warren CR. Mesophyll diffusion conductance to CO₂: An unappreciated central player in photosynthesis. *Plant Sci.* 2012; 193-194: 70-84.
26. Aalto T, Juurola E. A three-dimensional model of CO₂ transport in airspaces and mesophyll cells of a silver birch leaf. *Plant, Cell Environ.* 2002; 25(11): 1399-1409.
27. Kondo T, Kajita R, Miyazaki A, Hokoyama M, Nakamura-Miura T, Mizuno S, Masuda Y, Irie K, Tanaka Y, Takada S, Kakimoto T, Sakagami Y. Stomatal density is controlled by a mesophyll-derived signaling molecule. *Plant Cell Physiol.* 2010; 51(1): 1-8.

- 442 28. Magyar-Tábori K, Dobránszki J, Teixeira da Silva JA, Bulley SM, Hudák I. The role of
443 cytokinins in shoot organogenesis in apple. *Plant Cell Tiss. Org.* 2010; 101(3): 251-267.
- 444 29. Ferjani A, Yano S, Horiguchi G, Tsukaya H. Control of leaf morphogenesis by long-and
445 short-distance signaling: differentiation of leaves into sun or shade types and compensated
446 cell enlargement. *Plant Cell Monogr.* 2008; 10: 47-62.
- 447 30. Mediavilla S, Escudero A, Heilmeyer H. Internal leaf anatomy and photosynthetic resource-
448 use efficiency: interspecific and intraspecific comparisons. *Tree Physiol.* 2001; 21(4): 251-
449 259.
- 450 31. Gandolfo E, De Lojo J, Gómez D, Pagani A, Molinari J, Di Benedetto A. Anatomical changes
451 involved in the response of *Impatiens wallerana* to different pre-transplant plug cell volumes
452 and BAP sprays. *Eur. J. Hortic. Sci.* 2014; 79(4): 226-232.
- 453 32. Kobe RK, Iyer M., Walters MB. Optimal partitioning theory revisited: Nonstructural
454 carbohydrates dominate root mass responses to nitrogen. *Ecology.* 2010; 91(1): 166-179.
- 455 33. Price CA, Weitz JS. Zero-sum allocational strategies determine the allometry of specific leaf
456 area. *Am. J. Bot.* 2010; 97(11): 1808-1815.
- 457 34. Ron'zhina ES. Structural and functional rearrangements of mesophyll as a probable basis for
458 the cytokinin-dependent assimilate translocation in detached leaves. *Russ. J. Plant Physl.*
459 2004; 51(3): 333-341.
- 460 35. Arkhipova TN, Prinsen E, Veselov SU, Martinenko EV, Melentiev AI, Kudoyarova GR.
461 Cytokinin producing bacteria enhance plant growth in drying soil. *Plant Soil.* 2007; 292(1-2):
462 305-315.