

1
2
3
4
5
6
7

Original Research Article

GENETIC EFFECTS FOR MAIZE P EFFICIENCY TRAITS IN ACID AND NON-ACID SOILS OF WESTERN KENYA

ABSTRACT

Soil acidity is a major constraint to maize (*Zea mays* L.) productivity in tropical soils due to toxic levels of aluminium (Al) and phosphorus (P) deficiency. The objectives of this study were to: (i) determine the genetic effects of certain traits associated with phosphorus efficiency in maize (ii) compare the genetic control of maize P efficiency traits in acid and non-acid soils. Six F₁ single crosses derived from acid soil tolerant and susceptible lines were used in this study. The parental inbred lines, the F₁'s, F₂'s, BC₁P₁, BC₁P₂, from each of the six crosses were evaluated in two low P acid and two low P non-acid soils in Kenya. Mean genetic effect (m), additive genetic effects (a), dominant genetic effects (d) and epistatic digenic effects (aa, ad, dd) were computed for Shoot dry matter (SDM), Root Length density (RLD), P content (PC), P utilization efficiency (PUE) and P efficiency (PE). For most of the traits, greater variation was accounted for by dominance followed by epistatic and additive genetic effects in both acid and non-acid soils. Means for all the traits studied were significantly higher at high P conditions (36kgP/ha) in non-acid soils compared to acid soils for all the generations. Both RLD and PE exhibited higher means under nonacid soils compared to acid soils under both P conditions. Mean heritabilities were generally higher in non-acid soils compared to acid soils. There was higher reduction in PE in acid soils (25-50%) compared to non-acid soils (15 to 30%). The magnitude of both additive and non-additive gene effects were always greater in non-acid compared to acid soils pointing to the effects of soil acidity on gene action. The inheritance of major PE traits did not differ in acid and non-acid soils even though soil acidity affected the magnitude of the gene effects.

Keywords: maize, acid soils, additive, dominant, epistatic effects, phosphorus efficiency.

8
9

1. INTRODUCTION

10
11

Soil acidity is a major constraint to maize (*Zea mays* L) productivity worldwide mainly because of Aluminium (Al) toxicity and phosphorus (P) deficiency [1]. Al toxicity limits plant growth through its effects on root growth and development while P starvation leads to stunted growth, thin and spindly stems with purpling of leaves, reduced grain yields etc [2,3,4]. Soil acidity covers extensive areas in tropical, subtropical and temperate zones, and occurs in 30-40% of the world's arable soils [5]. They are found mainly in South America (26.7%), North America (19.4%), Africa (19.1%) and Asia (15.1%) while the rest occur in Australia and New Zealand, Europe and Central America [6, 3]. Different strategies have been suggested to improve the productivity of these soils including lime application, P replenishment through organic and inorganic sources and the development of tolerant

20
21

cultivars [7,8,4]. The latter approach has been preferred as the most suitable, sustainable and cost effective. Enormous genetic variation for tolerance to soil acidity that has been reported in several studies. [9] reported additive, dominance, and epistatic effects for P efficiency in maize with additive effects being more important while [10,11] reported the importance of both additive and dominance effects in controlling maize P efficiency traits. Other reports by [12, 13] showed non-additive effects to be more important than additive effects for tolerance to low P soils. Further results from studies carried out by [14,15,16] have shown that for grain yield, additive effects accounted for the major part of the total genetic variance, although non-additive effects were also significant. [17] reported that dominance followed by additive effects were more important for grain yield, plant height and days to anthesis in both acid and non-acid soils. According to [18], both additive and dominance effects were more important than epistatic effects in the inheritance of grain yield in acid soils. [18] further reported that for grain yield the additive-dominance model accounted for 91.1% of the variation in non-acid soils and 70.0% of the variation in acid soils, and that epistatic effects were more important in acid than in non- acid soils. These studies give hope that selection for P efficiency under acid and non-acid soils is possible.

So far, estimation of genetic effects on several important traits in maize evaluated under non-acid soils has been well documented [20, 21, 9, 10, and 11]. However information on the genetic control of maize P efficiency traits in acid soils is still inadequate given that the area under maize production in acid soils is quite substantial. Besides, with increased use of inorganic P-based fertilizers to restore soil fertility in agricultural systems, it is expected that the world acid soil area will increase in the future [18]. This is because some of the available inorganic P sources contribute to further soil acidification. A clear selection criteria for P efficiency in acid and non-acid soils and a better understanding of their genetic control is crucial in selecting for target traits for yield improvement and QTLs mapping studies. This is expected to accelerate the development of P efficient maize cultivars. The objectives of this

study were to (i) determine the genetic effects of traits associated with phosphorus efficiency in maize (ii) compare the genetic control of maize P efficiency traits in acid and non-acid soils.

2. MATERIAL AND METHODS

2.1 Genetic Materials

A total of six single crosses (KML 036 X MUL 229, HSL3 X 5046-2 X S396-16-1, KML 036 X S396-16-1, HSL3 X 5046-2XMUL 229, HS 228 X S 396-16-1 and HS 228 X MUL 229) were used to estimate the genetic effects in two acid soils sites (Sega and Chepkoilel) and two non-acid soils sites (Migori and Koyonzo). The parents were selected based on tolerance to low P conditions and their combining abilities [8]. The crosses were developed in 2010. For each cross, the F1 was advanced by selfing to obtain F2 generation in 2011. Backcross 1 to each parent (BC1P1 and BC1P2), were also obtained by crossing the F1 for each cross with each of its 2 parents, with the F1 as the female parent in 2012. At least 10 ears were saved and balance bulked to represent each generation.

2.1.1 Experimental Conditions

A total of 23 maize genotypes comprising 6 backcrosses (BC1), 6 F1 single crosses (SCH), 5 parental lines and 6 F2s were evaluated for tolerance to low P in a replicated trial at Sega, Chepkoilel, Migori and Koyonzo sites (Table 1) during the long rains of 2013.

Table 1: Site biophysical characterization

Site	Latitude	Longitude	Altitude	pH	P	Al.sat	Temp	Annual	Soil type
	(°)	(°)	masl		mgkg-1	(%)	range (oC)	rainfal (mm)	
Chepkolel	0° 37' N	035° 15' E	2143	4.8	4.4	45.6	13-26	1100	Chromic ferralsols
Sega	0° 15' N	34° 20' E	1200	4.5	2.2	44	17-30	1000	Orthic Acrisols
Migori	1° 03' S	34° 24' E	1381	5.7	2.6	12	22-24	1200	humic ferralsols
Koyonzo	0° 25' N	34° 25' E	1310	5.4	3	15	20-22	1400	Luvisols

Note Al. sat- Aluminium saturation, P- Amount of soil available phosphorus (0-30 cm) (Kisinyo et al., 2013).

The experiment was laid out in an RCBD replicated three times. Treatment consisted of the 24 genotypes and 2 levels of P described as low P (6 KgP/ha) and high P (36 KgP/ha) supplied as TSP). Generations were allocated to different blocks and randomized

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

independently. A two row plot measuring three metres long, with inter and intra-row spacing 0.75 m x 0.30 m was used for each generation except the F2 where four row plots were used. Two seeds were sown per hill and later thinned to one. Genstat software [22] was used to generate the randomization and field layout. All the plots were side-dressed using calcium Ammonium Nitrate (CAN) at the rate of 75 Kg N/ha. Standard agronomic practices were followed to maintain the experimental plots.

Root length density (RLD), shoot dry matter (SDM), phosphorus utilization efficiency (PUE), Shoot P concentration (SPC) and P efficiency (PE) were measured at anthesis. Destructive sampling was done on 16 randomly selected plants for all generations except F2 where 30 samples were used. Root sampling was done using the root box technique as described by Vepraskas and Hoyt [23] and Manske [24] in order to determine RLD. The line-intercept method described by Tennant [25] was used to determine RLD. Shoot samples were oven dried at 80°C, ground and ashed at 550 °C for determination of P concentration in the whole shoot. The ground samples were then dissolved in 3.3% HCl and analyzed for P using the method of Barton [26]. Based on shoot dry matter yield, and P concentration in these plant components, the P content in the shoot (PC) and PUE were determined using the method of [27,28]. The P efficiency ratio was calculated as the ratio of shoot dry matter production under low P to that under adequate P supply [29].

2.1.1.1 Data Analysis

Generation means for each cross and P treatment were used to estimate the gene effects according to the [29] model following [31] notation to define the genetic parameters in the model. This model was as follows:

$Y_k = m + \alpha a + \beta d + \alpha^2 aa + \alpha\beta ad + \beta^2 dd$, where α and β are the coefficients for additive and dominance effects,

Y_k = the observed mean across locations of the kth generation

102 m = mean of all possible homozygous locus, considering all locus controlling the trait;

103 a = pooled additive effects

104 d = pooled dominance effects

105 aa = additive x additive gene interaction effects

106 ad = additive x dominance gene interaction effects

107 dd = dominance x dominance gene interaction effects

108 Estimates of additive, dominance and epistatic effects were computed for each cross by
109 weighted least square regression analysis [32,30] using the equation $b = (X' D^{-1} X)^{-1} (X' D^{-1} y)$, where b is the vector of genetic effects (m, a, d, aa, ad, and dd), X is the incidence
110 matrix of the genetic effects coefficients (α , β , α^2 , $\alpha\beta$, and β^2), y is the column vector of the
111 generation means and D-1 is a weighted diagonal matrix, where the diagonal elements were
112 the reciprocals of the variances of each generation mean computed for each generation (P's,
113 F1's, F2's, and BC's). Statistical Analysis System [33] was used to estimate the genetic
114 effects from the generation means of each cross at each P level and combined over
115 locations. F test of the sum of squares for the genetic effects was used to reduce the model
116 appropriately. In the selected model, genetic parameters having significant effects were
117 included and all the non-significant parameters excluded from the model. For each cross in
118 each trait, the ratios a/m, d/m and epistasis/m were calculated using absolute values. Only
119 data where the parameter estimates (a, d and epistasis) were significant were used in these
120 calculations. For each trait and at the two P levels across the locations, a general mean of
121 the ratios a/m and d/m was calculated using data from all crosses with significant effects.
122 Broad sense heritability (H^2) was estimated by variance components using linear mixed
123 models (REML) as follows:
124

125 $H^2 = \sigma_g^2 / \{\sigma_g^2 + (\sigma_{error}^2/r)\}$, Where H^2 is broad sense heritability, σ_g^2 is the generic variance,
126 σ_{error}^2 is the error variance, r is the number of replicates per genotype [34].

3. RESULTS AND DISCUSSION

3.1 Trait means and heritabilities in acid soils of western Kenya.

In acid soils, shoot dry matter yields (SDM) were significantly higher at high P compared to low P ones for all the generations tested. Higher P supply increased mean SDM from 0.17 to 0.25 kg per plant in the parentals, 0.26 to 0.46 kg/plant, in the F1s, 0.25 to 0.45 kg/plant for the backcrosses and 0.13 to 0.26 kg/plant for the F2s (Table 2). The F1s attained the highest SDM under both high (0.46 kg/plant) and low P (0.26 kg/plant) supplies while the parental lines yielded the least (0.25 kg/plant) under high P and the F2s gave the least under low P (0.13 kg/plant). Mean H^2 for SDM was generally higher under high P compared to low P conditions for all the generations except for the F2s where the reverse was true. The F1s exhibited the highest heritability at high (0.603) and low P conditions (0.57) (Table 2). The highest mean root length density (RLD) was obtained in the F1 at high P (10.23 cm/cm³) and lowest in the parents (6.66 cm/cm³) while the backcrosses exhibited the highest RLD (6.37 cm/cm³) under low P. The highest heritability for RLD was obtained in the backcrosses while lowest in the F2s. Low P supply resulted in a significant reduction by (23 to 50%) in shoot P concentration of genotypes in all the generation. Backcrosses exhibited the highest mean PUE (559.28 gSDM/gP) while parentals the least (520.9 gSDM/gP) although the difference was not large. The highest mean PE (54%) was measured in the F1s that also exhibited very high mean SDM under low P conditions

3.2. Trait means and heritabilities in non-acid soils of western Kenya.

In non-acid soils, SDM yields were significantly higher under high P compared to low ones for all the generations. Mean SDM was increased at least two folds by the application of higher P in all the generations. The F1s attained the highest SDM under both high (0.52 kg/plant) and low P (0.229 kg/plant) supplies while the F2s yielded the lowest under high P (0.24 kg/plant) and low P (0.12 kg/plant). Mean broad sense heritability for SDM did not show any clear pattern. The BC1 exhibited the highest heritability under both high (0.76) and low P conditions (0.72) (Table 3). Under high P supply, the highest mean RLD was obtained in the BC1 (20 cm/cm³) and the lowest in the F2 generation (11.87 cm/cm³) while at low P,

the backcrosses exhibited the highest RLD (14.1 cm/cm³) and the F₂ generation the lowest (8.16 cm/cm³). The highest mean H₂ was obtained in the F₁s and BC₁s under high and low P respectively while the lowest in the F₂s. Low P supply resulted in a significant reduction (15.5 to 31%) in shoot P concentration (SPC) of genotypes in all the generations. Backcrosses exhibited the highest mean PUE (645.1 gSDM/gP) while parentals the least (485.88 gSDM/gP). The highest mean PE (59%) was measured in the F₁s that also exhibited high mean SDM under low P conditions.

Table 2: Mean shoot dry matter (SDM), root length Densitydensity, (RLD) and P concentration (PC) of maize genotypes evaluated in 2 acid soil sites in western Kenya.

Formatted: Highlight

ENTRY	SDM (kg/plant)		RLD (cm/cm3)		PC (%)		PUTE	PE
	P36	P6	P36	P6	P36	P6	gSDM/gP	%
S1 (Parents)	0.248	0.185	5.657	3.244	0.163	0.126	570	48
K1	0.278	0.187	7.69	4.407	0.161	0.107	488	51
H2	0.253	0.172	7.455	3.349	0.17	0.13	382	56
H1	0.273	0.185	8.391	4.572	0.142	0.117	582	50
H3	0.176	0.13	5.422	3.611	0.157	0.121	566	48
M1	0.245	0.143	5.359	3.569	0.153	0.12	537	42
MEAN	0.25	0.17	6.66	3.79	0.16	0.12	520.98	49.22
H2	0.581	0.395	0.227	0.607	0.645	0.404		
SE	0.02	0.02	1.162	0.854	0.01	0.006		
LSD (0.05)	0.016	0.011	0.77	0.47	0.01	0.008	35.2	
K1XS1 (F1s)	0.485	0.253	8.457	6.455	0.149	0.125	583	48
H1XS1	0.452	0.287	9.372	6.147	0.144	0.116	463	59
H2XS1	0.46	0.233	8.591	5.011	0.155	0.117	588	45
H1XM1	0.385	0.237	10.974	5.435	0.156	0.11	491	65
H2XM1	0.477	0.252	9.331	5.844	0.133	0.127	567	48
K1XM1	0.518	0.297	14.681	8.898	0.139	0.103	474	60
MEAN	0.46	0.26	10.23	6.3	0.15	0.116	527.66	54.03
H2	0.603	0.576	0.638	0.583	0.1	0.441		
SE	0.025	0.019	1.009	0.821	0.006	0.005		
LSD (0.05)	0.038	0.021	0.79	0.48	0.01	0.008	37.4	
H1XS1*H1(E	0.458	0.242	13.255	6.759	0.158	0.127	589	52
H1XS1*S1	0.493	0.222	8.844	6.016	0.167	0.112	773	40
K1XS1*K1	0.46	0.278	8.989	4.944	0.156	0.13	641	58
K1XM1*K1	0.447	0.27	9.899	8.202	0.157	0.131	538	56
K1XM1*M1	0.388	0.186	11.226	8.339	0.162	0.12	602	45
H2XM1*H2	0.418	0.232	7.27	5.627	0.179	0.114	406	51
H1XM1*M1	0.487	0.247	9.516	5.828	0.156	0.133	553	52
H1XM1*H1	0.412	0.262	7.135	5.734	0.266	0.117	384	57
H2XS1*H2	0.467	0.277	11.198	5.856	0.167	0.116	548	56
MEAN	0.45a	0.25a	9.70b	6.37a	0.17	0.12	559.28	51.84
H2	0.749	0.471	0.67	0.721	0.1	0.384		
SE	0.02	0.016	0.546	0.156	0.019	0.006		
LSD (0.05)	0.029	0.017	0.7	0.63	0.009	0.0075	36.8	
K1XM1(F2 s	0.26	0.135	4.626	3.862	0.141	0.117	689	42
H1XM1	0.31	0.175	7.394	4.905	0.16	0.1	492	50
H1XS1	0.243	0.133	8.25	4.44	0.145	0.128	391	50
K1XS1	0.27	0.113	7.341	5.01	0.145	0.098	689	40
H2XM1	0.23	0.147	10.714	5.184	0.143	0.115	294	62
H2XS1	0.231	0.087	6.197	4.271	0.149	0.112	682	49
MEAN	0.26	0.13	7.42	4.61	0.15	0.11	539.38	49.02
H2	0.407	0.549	0.417	0.504	0.256	0.15		
SE	0.019	0.015	1.212	0.598	0.006	0.006		
LSD (0.05)	0.017	0.008	0.9	0.51	0.008	0.006	40.4	

Note: H1-HSL3 X 5046-2, H2-HS 228, M1-MUL 229, S1-S396-16-1

Table 3: Mean shoot dry matter (SDM), root length density, (RLD) and P concentration (PC) of maize genotypes evaluated in 2 non-acid soil sites in western Kenya.

ENTRY	SDM (kg/plant)		RTLD (cm/cm ³)		PC (%)		PUTE gSDM/gP	PE %
	P36	P6	P36	P6	P36	P6		
S1 (Parents)	0.353	0.107	11.430	8.074	0.177	0.133	409	43
K1	0.258	0.120	12.906	10.650	0.174	0.163	693	53
H2	0.362	0.157	13.376	6.730	0.159	0.145	608	66
H1	0.338	0.217	13.091	10.958	0.165	0.133	307	65
H3	0.277	0.140	11.126	10.644	0.155	0.160	557	58
M1	0.262	0.140	12.704	7.489	0.134	0.110	341	54
MEAN	0.31	0.15	12.44	9.09	0.161	0.141	485.88	56.53
H2	0.391	0.703	0.718	0.485	0.591	0.30		
SE	0.035	0.024	0.868	0.703	0.009	0.026		
Lsd (0.05)	0.022	0.0107	0.85	0.64	0.0115	0.010	35.14	
K1XS1 (F1s)	0.492	0.203	20.000	12.575	0.144	0.142	630	59
H1XS1	0.561	0.263	18.998	12.976	0.127	0.130	608	62
H2XS1	0.574	0.227	21.998	11.214	0.141	0.142	760	52
H1XM1	0.518	0.277	10.606	10.407	0.164	0.132	361	73
H2XM1	0.497	0.177	13.215	12.622	0.189	0.123	457	47
K1XM1	0.483	0.230	25.291	17.333	0.158	0.136	413	64
MEAN	0.521	0.229	18.351	12.855	0.154	0.134	538.298	59.432
H2	0.504	0.618	0.781	0.621	0.598	0.159		
SE	0.025	0.017	1.281	0.865	0.011	0.024		
Lsd (0.05)	0.025	0.015	1.42	0.85	0.0107	0.010	37.23	
H1XS1*H1(Bcs)	0.365	0.320	22.540	15.465	0.156	0.132	440	75
H1XS1*S1	0.378	0.240	17.778	9.570	0.141	0.113	659	70
K1XS1*K1	0.416	0.250	19.201	16.769	0.146	0.134	530	63
K1XM1*K1	0.462	0.237	27.044	18.140	0.150	0.116	694	58
K1XM1*M1	0.341	0.177	21.663	14.094	0.158	0.157	752	49
H2XM1*H2	0.418	0.187	19.928	14.680	0.166	0.140	716	49
H1XM1*M1	0.371	0.203	17.008	15.469	0.142	0.100	492	59
H1XM1*H1	0.362	0.167	16.801	13.777	0.162	0.102	874	47
H2XS1*H2	0.435	0.200	18.182	9.005	0.156	0.114	650	47
MEAN	0.394	0.220	20.016	14.108	0.153	0.123	645.107	57.406
H2	0.757	0.716	0.747	0.78	0.123	0.190		
SE	0.021	0.019	0.941	0.851	0.006	0.028		
Lsd (0.05)	0.034	0.025	1.33	0.75	0.011	0.007	46.07	
K1XM1(F2 s)	0.236	0.143	11.796	7.893	0.170	0.126	393	64
H1XM1	0.300	0.113	12.568	9.808	0.167	0.101	653	42
H1XS1	0.231	0.123	11.897	9.063	0.172	0.105	329	53
K1XS1	0.205	0.150	13.229	8.317	0.153	0.124	482	67
H2XM1	0.236	0.103	10.350	8.608	0.167	0.128	591	46
H2XS1	0.244	0.097	11.401	5.291	0.187	0.114	648	41
MEAN	0.242	0.122	11.873	8.163	0.169	0.117	515.892	52.290
H2	0.524	0.45	0.454	0.509	0.316	0.17		
SE	0.023	0.016	2.156	1.808	0.010	0.014		
Lsd (0.05)	0.0184	0.0093	0.91	0.63	0.013	0.0092	37.80	

Note: H1-HSL3 X 5046-2, H2-HS 228, M1-MUL 229, S1-S396-16-1

3.3.1 Mean comparison and heritabilities under acid and non-acid soils

Mean SDM was greater under nonacid soils compared to acid soils at high P supply (Table 4). However under low P, SDM values were generally higher in acid compared to non-acid soils for all the generations evaluated. Both RLD and PE exhibited higher means under nonacid soils compared to acid soils under both P conditions. Highest PE was exhibited by the F1s in both acid and non-acid soils although it was higher in non-acid (59%) compared to

acid soils (54 %). There was higher reduction in PE in acid soils (25-50%) compared to non-acid soils (15 to 30%). Measurement of PUE was greatly dependent on the generation for both soils. Mean PUE was higher for parentals and the F2 generations in acid (520.98 gSDM/gP and 539.38 gSDM/gP) compared to non-acid soils (485.88 gSDM/gP and 515 gSDM/gP), while it was higher for F1s and BC1 in non-acid soils (538.29 gSDM/gP and 645.1 gSDM/gP) compared to acid soils (527.66 gSDM/gP and 559.28 gSDM/gP). However, there were no significant differences for PC between the two soils, although higher magnitudes of PC were detected for non-acid soils (Table 4). Mean heritabilities were generally higher in non-acid soils compared to acid soils although this depended on the generation, the trait and the level of P applied (Table 4).

Table 4: Mean broad sense heritabilities for SDM and other P efficiency traits in acid and non-acid soils.

ACID SOILS (CHEPKOILEL AND SEGA)						
Trait generation	SDM (kg/plant)		RLD (cm/cm ³)		PC (%)	
	P36	P6	P36	P6	P36	P6
Parents	0.581	0.395	0.227	0.607	0.645	0.404
F1S	0.603	0.576	0.638	0.583	0.1	0.441
Backcrosses	0.749	0.471	0.67	0.721	0.1	0.384
F2S	0.407	0.549	0.417	0.504	0.256	0.15
NON ACID SOILS (KOYONZO AND MIGORI)						
Trait generation	SDM (kg/plant)		RLD (cm/cm ³)		PC (%)	
	P36	P6	P36	P6	P36	P6
Parents	0.391	0.703	0.718	0.485	0.591	0.3
F1S	0.504	0.618	0.781	0.621	0.598	0.159
Backcrosses	0.757	0.716	0.747	0.78	0.123	0.19
F2	0.524	0.45	0.454	0.509	0.316	0.17

3.4 Gene effects in acid and non-acid soils

3.4.1. Shoot dry matter

In high P acid soils, majority of the crosses (83%) exhibited significant dominance gene action compared to additivity (33%) although additivity was more pronounced under low P supply (67%). Consequently the magnitude of mean dominance was higher under both P conditions compared to mean additive genetic effects (0.88 and 0.55 vs. 0.075 and 0.085) for dominance and additive under high and low P respectively. Epistatic effects were only detected for a few crosses under both P conditions. The mean of ratio "a/m", "d/m", and "epistasis/m" at high P was 0.27, 3.58 and 1.69 respectively while they were 0.74, 6.1 and

5.6 at low P, respectively. This indicates that dominance and epistatic effects were more important in the expression of SDM than additive effects under both P conditions in acid soils. However, both dominance and additive effects had higher magnitude under low P conditions (d/m, a/m ratios 6.1 and 0.74) compared to high P conditions (d/m, a/m ratios of 3.5 and 0.027) (Table 5). In non-acid soils, majority of the crosses exhibited significant dominance gene action (83% and 67%) compared to additive (33 and 17%) gene action under high and low P respectively. Therefore mean dominance was higher compared to mean additive genetic effects (0.81 vs. 0.11 and 0.55 vs. 0.14) in high and low P conditions. Significant epistatic effects (dominance x dominance) were detected at both P levels. The magnitude of epistatic effects were greater than dominance under both P conditions, although they were more pronounced under low P (6.13) compared to high P(4.29) (Table 6). Therefore epistasis followed by dominance was more important in SDM inheritance compared to additive effects in non-acid soils.

Table 5: Estimates of genetic effects for shoot dry matter (Kg/plant) evaluated in two acid soil locations in in western Kenya.

Shoot dry matter at High P (acid soils of Chepkoilel and Segla)									
Cross	M	A	d	aa	ad	dd	a/m	d/m	Epist/m
K1XS1	0.27**	0.05	0.80**	-	-	-	-	2.96	-
H1XS1	0.24**	-0.04	1.04**	0.93*	-	-1.25*	-	4.28	1.33
H2XS1	0.23**	0.05	0.98**	0.85*	-	-1.04*	-	4.23	0.82
H1XM1	0.31**	-0.08*	0.60	-	-	-0.91*	0.24	-	2.93
H2XM1	0.23**	0.07*	0.76*	-	-	-	0.30	3.32	-
K1XM1	0.26**	0.06	0.81*	-	-	-	-	3.11	-
Mean	0.26	0.075	0.88	-	-	-	0.27	3.58	1.69
Shoot dry matter) at low P (acid soils of Chepkoilel and Segla)									
Cross	M	A	d	aa	ad	dd	a/m	d/m	Epis/tm
K1XS1	0.11*	0.10*	0.54*	0.47*	-	-	0.85	4.80	4.3
H1XS1	0.13*	0.02	0.50	-	-	-	-	-	-
H2XS1	0.09*	0.09*	0.64*	-	-	-0.69*	1.00	7.35	6.9
H1XM1	0.18*	0.02	0.37	-	-	-	-	-	-
H2XM1	0.15*	0.08*	0.27	-	-	-	0.52	-	-
K1XM1	0.14*	0.08*	0.48	-	-	-	0.62	-	-
Mean	0.13	0.085	0.59	-	-	-	0.74	6.1	5.6

Gene effects: m: mean; a: additive; d: dominance; aa, ad and dd are additive x additive, additive x dominance and dominance x dominance epistasis, respectively; Only significant estimates of the parameters were used to obtain the means. . Means were obtained using absolute values; Significance at 5% (*) and 1% (**) probability level

Table 6: Estimates of genetic effects for shoot dry matter evaluated in two low P non-acid soil locations in western Kenya.

Shoot dry matter (Kg/plant) at High P (non-acid soils of Migori and Koyonzo)									
Cross	m	A	d	aa	ad	dd	a/m	d/m	Epist/m
K1XS1	0.20**	-0.04	0.98**	--	--	-1.37*	-	4.78	6.85
H1XS1	0.23**	-0.01	0.64*	--	--	--	-	2.75	-
H2XS1	0.24**	-0.09*	1.01**	--	--	-1.26*	0.35	4.16	5.25
H1XM1	0.30**	-0.01	0.34	--	---	--	-	-	-
H2XM1	0.24**	0.04	0.70*	--	--	-0.91*	-	2.95	0.67
K1XM1	0.24**	0.12*	0.74**	--	--	-1.06*	0.51	3.14	4.41
Mean	0.242	0.11	0.81	-	-	-1.23	0.43	3.56	4.29
Shoot dry matter (Kg/plant) at Low P (non-acid soils of Migori and Koyonzo)									
Cross	m	A	d	aa	ad	dd	a/m	d/m	Epist/m
K1XS1	0.14**	0.05	0.15	-	-	-	-	-	-
H1XS1	0.12**	0.01	0.59*	-	-	-0.72*	-	4.92	6.0
H2XS1	0.11**	-0.02	0.71*	-	-	-0.86*	-	6.46	7.8
H1XM1	0.13**	-0.06	0.46*	-	-	-	0.46	3.44	-
H2XM1	0.13**	0.05	0.21	-	-	-	-	-	-
K1XM1	0.15**	0.14*	0.46*	-	-	-0.69*	0.93	3.18	4.6
Mean	0.13	0.14	0.55	-	-	-0.75	0.67	4.5	6.13

Gene effects: m: mean; a: additive; d: dominance; aa, ad and dd are additive x additive, additive x dominance and dominance x dominance epistasis, respectively; Only significant estimates of the parameters were used to obtain the means. Means were obtained using absolute values; Significance at 5% () and 1% (**) probability level*

3.4.2 Root Length Density

The number of crosses under high P acid soils with significant additive effects was larger (100%) than those with significant additive effects in low P (50%) (Table 7). No epistasis was detected for RLD in acid soils. The overall mean ratio for "a/m", "d/m", at low P was 0.39 and 2.88, respectively while they were 0.17 and 2.13 respectively at high P. This shows that dominance was more important than additive effects in the inheritance of RLD in both P conditions. Both dominance and additive effects were more pronounced at high P conditions compared to low P conditions.

Table 7: Estimates of genetic effects for Root Length density evaluated in two acid soil locations in western Kenya.

Root Length density (cm/cm ³) at High P (acid soils of Chepkoilel and Sega)									
Cross	M	a	d	aa	ad	dd	a/m	d/m	Epist/m
K1XS1	7.34**	2.99*	2.40	-	-	-	0.41	-	-
H1XS1	8.25**	4.41*	13.55*	-	-	-	0.53	1.64	-
H2XS1	6.20**	4.20*	13.64*	-	-	-	0.68	2.20	-
H1XM1	7.39**	2.38*	6.82	-	-	-	0.32	-	-
H2XM1	10.71**	1.27*	-13.39*	-	-	-	0.12	1.25	-
K1XM1	4.63**	-1.33*	29.90**	-	-	-	0.29	6.46	-
Mean	7.42	2.76	15.46				0.39	2.88	-
Root Length density (cm/cm ³) at Low P acid soils of Chepkoilel and Sega)									
Cross	m	a	d	aa	ad	dd	a/m	d/m	Epist/m
K1XS1	5.01**	0.62*	0.27	-	-	-	0.12	-	-
H1XS1	4.44**	0.74*	8.78*	-	-	-	0.17	1.98	-
H2XS1	4.27**	0.89*	5.84*	-	-	-	0.21	1.37	-
H1XM1	4.91**	-0.29	4.12*	-	-	-	-	0.84	-
H2XM1	5.18**	-0.25	4.46*	-	-	-	-	0.86	-
K1XM1	3.86**	-0.34	21.8*	-	-	-	-	-	-
Mean	4.61	0.75	9.00				0.17	2.13	-

Gene effects: m: mean; a: additive; d: dominance; aa, ad and dd are additive x additive, additive x dominance and dominance x dominance epistasis, respectively; Only significant estimates of the parameters were used to obtain the means. . Means were obtained using absolute values; Significance at 5% (*) and 1% (**) probability level

The number of crosses with significant additive effects was smaller (33.3%) in high P compared to low P (50%) non-acid soils (Table 8). At both P conditions dominance effects were more important followed by epistatic effects and additive effects, although both dominance and additive effects had higher magnitude under low P conditions (d/m, a/m ratios of 3.55 and 0.36) than under high P conditions (d/m, a/m ratios of 2.0 and 0.33). Epistatic effects (aa and dd) for RLD were detected at both P conditions in non-acid soil in four out of the six crosses although they were more pronounced at high P (epist/m ratio of 1.77) compared to low P (epist/m ratio of 1.08). These results agree with those of Wolf and Hallauer (1997) who concluded that epistasis in maize seems to be more important in either poorer or better environments. These findings also suggest that variation in P levels in non-acid soils did not affect the detection of epistatic effects for RLD.

Table 8: Estimates of genetic effects for Root-root Length-length density evaluated in two non-acid soil locations in western Kenya.

Formatted: Highlight

Root Length density (cm/cm ³) at High P (non-acid soils of Migori and Koyonzo)									
Cross	m	a	d	aa	ad	dd	a/m	d/m	Epist/m
K1XS1	13.2**	4.77*	32.1**	24.27*	-	-	0.36	2.43	1.83
H1XS1	11.9**	5.89*	20.6**	-	-	-	0.50	1.76	-
H2XS1	11.4**	3.00	18.44*	-	-	-30*	-	1.62	2.6
H1XM1	12.6**	2.31	8.97	-	-	-	-	-	-
H2XM1	10.3**	3.68	17.10*	16.93*	-	-	-	1.65	1.64
K1XM1	11.8**	4.05*	45.4**	32.89*	-	-21.17*	0.24	3.85	0.99
Mean	11.87	4.9	26.7	-	-	-	0.37	2.26	1.77
Root Length density (cm/cm ³) at Low P (non-acid soils of Migori and Koyonzo)									
Cross	m	a	d	aa	ad	dd	a/m	d/m	Epist/m
K1XS1	8.32**	1.77	36.48**	30.27*	-	-43.9*	-	4.39	1.63
H1XS1	9.06**	5.89*	20.28**	-	-	-	0.65	2.24	-
H2XS1	5.29**	1.00	19.66**	-	-	-	-	3.72	-
H1XM1	9.81**	-1.69	23.44**	19.26	-	-32.49*	-	2.39	1.32
H2XM1	8.61**	3.68	25.44**	16.93	-	-22.82*	-	2.96	0.68
K1XM1	7.89**	4.05*	44.16**	32.89*	-	-38.56*	0.51	5.59	0.718
Mean	8.16	4.97	28.24	-	-	-	0.58	3.55	1.08

*m: mean; a: additive; d: dominance; aa, ad and dd are additive x additive, additive x dominance and dominance x dominance epistasis, respectively; *, and **, Significance at 5% (*) and 1% (**) probability level*

3.5 DISCUSSIONS

Maize genotypes differed significantly both in shoot and root growth at low P supply and in response to P fertilization. Such substantial genetic variation in response to P deficiency and P supply was also observed in past studies for maize hybrids [2, 35, and 4] sorghum [36, 37], *Brassica oleracea* [27] and wheat [29]. The application of high P fertilizer increased SDM, RLD, PE and PUE in both acid and non-acid soils because of the increased soil available P, which is necessary for healthy plant growth and high grain production. Besides, Soil P availability is critical for the early growth and development of maize as it affects root morphological and physiological characteristics that are important for eventual P uptake since P is immobile and often unavailable in most soils [38, 39]. Similar results have been reported in maize for increased root length density, grain yield, PE and PUE due to increased P application. [38,40] and in wheat [41, 29].The increments in P efficiency traits measured were also due to the fact that P is involved in several key plant functions including energy transfers, photosynthesis, transformation of sugar and starches, nutrient movement

within the plants e.t.c [42] hence the increase in SDM and root growth and development. The finding for RLD is however contrary to those of [43] who recorded a decrease in root growth as a result of increased P supply in bean recombinant inbred lines. The contrasting results could be explained by the difference in plant types, adaptation and growth habits between maize and beans.

There was a general reduction in SDM and other P efficiency traits in acid soils compared to non-acid soils. This was in addition to the higher reduction in PE in acid soils (25-50%) compared to non-acid soils (15 to 30%). These results compare well with those of [17] who observed significant decrease in maize grain yields and plant height for parentals, F1, F2 and Backcross generations. [24] also reported decrease in RLD as a result of soil acidity in bread wheat genotypes. These observations can be attributed to the detrimental effects of soil acidity on maize performance, grain yield and other agronomic traits [44, 19, 2, 7, 8, 4].

Apart from the low available P, the crops in the acid soils also suffered from Al toxicity which decreased their cell division hence reduced SDM, RLD and PE.

Lower heritabilities in acid soils were probably due to high experimental error and low generic variations depicted under such conditions. These findings compare well with those of [45] who obtained lower heritabilities estimates under stress environments. However, in part of this study, higher heritabilities were reported in acid soils compared to non-acid soils. These findings compare well with those of [46, 47] who reported greater genetic variation under stress environments and suggested that heritability in such environments can sometimes be comparable to non-stress environments or even higher if the experimental error is of the same magnitude.

In both acid and non-acid soils, dominance and epistasis were more important than additive portion although epistatic effects were more pronounced in non-acid soils while dominance more pronounced in acid soils for SDM. The magnitude of additive, dominance and epistasis was always higher in low P compared to high P in both acid and non-acid soils. This implies that selecting for SDM is more suitable under low P soils in both acid and non-acid soils.

These findings compare well with those of [29, 43, and 27] who found SDM as suitable selection criteria for P efficiency under low P conditions for beans, brassica and wheat genotypes. For RLD, there was higher magnitude of additivity in high P acid soils compared to the low magnitude of additivity at low P acid soils (Table 7 and 8). These findings imply that selection for RLD in acid soils is more suitable under high P conditions because of high additive effect. Such selection strategy may lead to identifying good responders rather than efficient genotypes. However, it would still be suitable in acid soils considering that a large proportion of soil P is held very tightly to the surface of soil particles as organic phosphorus compounds and hence unavailable even at high P supplementation. In contrast, selection for RLD in non-acid soils is more suitable at low P levels because of high additivity and dominance gene effects

4. CONCLUSION

Both additive and non-additive effects were detected in both acid and non-acids soils although this was more dependent on the trait studied and the level of available P. Dominance effects played a more important role than epistatic effects and the latter were more important than additive effects in the inheritance of majority of P efficiency traits studied in maize in both acid and non-acid soils. In most cases, epistasis was specifically more important in non-acid soils while dominance in acid soils. Additive gene effects were fairly of similar magnitude across the acid and non-acid soils. The magnitude of both additive and non-additive gene effects were always greater in non-acid soils compared to acid soils pointing to the detrimental effects of soil acidity in the detection of gene actions in maize. Our results suggest that the inheritance of Root Length Density and Shoot Dry matter did not differ in acid and non-acid soils. The overall results of this study showed that soil acidity significantly reduced P efficiency traits in maize and affected the detection of the genetic effects for these traits

ACKNOWLEDGEMENTS

The authors acknowledge the financial support provided by the Generation challenge program (GCP). We also thank the farmers for providing land for conducting the Research.

REFERENCES

1. Kochian L.V, Migue, A. P, Jiping L and. Magalhaes J.V. Plant Adaptation to Acid Soils: The Molecular Basis for Crop Aluminum Resistance. Annual Review of Plant Biology. 2015. 66: 571-598
2. Parentoni, S.N., Souza JR., Alves, V.M.C.; Gama, E.E.G.; Coelho, A.M.; Oliveira, A.C. Guimaraes, P.E.O., Guimaraes, C.T., Vasconcelos, M.J.V., Pacheco, C.A.P., Magalhães, J.V., Meirelles, W.F.,Guimarães, L.J.M., Silva, A.R., Mendes, F.F., Schaffert, R.E..Inheritance and breeding strategies for phosphorus efficiency in tropical maize (*Zea mays* L.). *Maydica*. 2010. 55: 1-15.
- 3.Ouma E., Ligeyo, D., Matonyei, T., Were B., Agalo, J., Emily. T, Onkware, A., Gudu S., Kisinyo, P., Okalebo, J and Othieno C. Development of maize single cross hybrids for tolerance to low phosphorus. African Journal of plant science. 2012. 6(14):394-402.
4. Ligeyo D.O, E.Ouma, S.Gudu, P.O.Kisinyo, T. Matonyei, J.R. Okalebo and C.O. Othieno. Response of maize top cross hybrids to low phosphorous in acid soils of western Kenya. East. Africa. Forestry.J. 2014. 80(1) 25-30.
5. Von Uexkull, H.R. and Mutert, E. Global extent, development and economic impact of acid soils. Plant soil.1995.171: 1-15
6. Eswaran, H., Reich, P. and Beinroth, F. Global distribution of soils with acidity. "In Plant-Soil Interaction at Low PH" (A.C. Moniz, ed.).1997.159-164. Brazilian soil science society.
7. Kisinyo P, Gudu S, Othieno C, Okalebo J, Ochuodho J, Agalo J, Ng'etich W, Opala R, Maghanga, J.K, Kisinyo J.A., Opile W.R, Ogola B.O. Phosphorus Sorption and Lime Requirements of Maize Growing Acid Soils of Kenya. Sustainable Agriculture Research. 2013. 2(2):116-123.
8. Ouma E., Ligeyo, D., Matonyei, T., Were, B., Agalo, J., Emily. T, Onkware, A., Gudu S., Kisinyo, P., Okalebo J.and Othieno C. Enhancing maize grain yield in acid soils of western Kenya using aluminium tolerant germplasm. Journal of Agricultural Science and Technology A .2013.3.33-46.
9. Duncan, R.R. Genetic manipulation.1994.1-38. In: R.E. Wilkinson (Ed.), Plant-environment interactions. New York: Marcel Dekker.
10. Chaubey, C.N., D. Senadhira, G.B. Gregorio. Genetic analysis of tolerance for phosphorous deficiency in rice (*Oryza sativa* L.). Theor. Appl. Genet. 1994. 89: 313-317.
11. Furlani A.M.C., LIMA, M., NASS, L.L. Combining ability effects for P-efficiency characters in maize grown in low P nutrient solution. *Maydica* .1998. 43: 169-174
12. Parentoni, S.N., Souza JR., Alves, V.M.C.; Gama, E.E.G.; Coelho, A.M.; Oliveira, A.C. Guimaraes, P.E.O., Guimaraes, C.T., Vasconcelos, M.J.V., Magalhães, J.V and Schaffert, R.E.Breeding maize for Al tolerance, P use efficiency and acid soil adaptation for the cerrado areas of Brasil: the EMBRAPA's experience. pp. 129-131. In: Proc. 3rd Intl. Symposium on Phosphorus Dynamics in the Soil-Plant Continuum. Uberlândia, Minas Gerais, Brazil.2006. May 14- 19.
13. Chen, J., XU, L., CAI, Y. XU, J. Identification of QTLs for phosphorus utilization efficiency in maize (*Zea mays* L.) across P levels. *Euphytica* .2009.167: 245-252.
14. Magnavaca, R., Gardner, C.O and Clark, R.B. Inheritance of aluminum tolerance in maize. In: Gabelman, H.W and Loughman, B.C. (eds) Genetic Aspects of Plant Mineral Nutrition. Martinus Nijhoff Publishers.1987b.Dordrecht: 201-212.

Formatted: Highlight

Formatted: Normal, Justified

Formatted: Font: (Default) Helvetica

15. Pandey, S., Ceballos, H., Granados, G. and Knapp, E. Developing maize that tolerates aluminium toxic soils. In "Stress tolerance Breeding: Maize That resist insects, Drought, Low Nitrogen and Acid Soils" eds Edmeades, G.E. and Deutsch, J.A. 1994. CIMMYT, Mexico
16. Salazar, F., Pandey, S., Narro, L., Pérez, J.C., Ceballos, H., Parentoni, S., and Bahia Filho A.F.C. Diallel analysis of acid soil tolerant and intolerant tropical maize populations. *Crop Sci.* 1997. 37:1457-1462.
17. Velásquez J.C., Pérez, C., Lopes de Souza Jr., Luis A. N., Shivaji P. and Carlos D.L. Genetic effects for maize traits in acid and non-acid soils. *Genetics and Molecular Biology.* 2008. 31, (1) 89-97.
18. Pandey, S., Narro, L., Friesen, D.K. and Waddington, S.R. Breeding maize for tolerance to soil acidity. *Plant Breed Rev.* 2007. 28:59-100.
19. Ceballos, H., Pandey, S., Narro, L. and Pérez, J.C. Additive, dominant, and epistatic effects for maize grain yield in acid and non-acid soils. *Theor Appl Genet.* 1998. 96:662-668.
20. Hallauer, A.R., Russell, W.A and Lamkey, K.R. Corn breeding. In: Sprague, G.F and Dudley, J.W (eds) *Corn and Corn Improvement.* Am Soc Agron. 1988. Madison: 463-564.
21. Hallauer, A.R and Miranda Filho, J.B. *Quantitative Genetics in Maize Breeding.* Iowa State Univ. Press. 1988. Ames, 468 pp.
22. Payne, R.W., Murray, D.A., Harding, S.A., Baird, D.B. & Soutar, D.M. *GenStat for Windows* (18th Edition). VSN International. 2014. Hemel Hempstead
23. Vepraskas, M.J., and Hoyt, D. Comparison of the trench profile and core methods for evaluation of root distribution in tillage studies. *Agron. J.* 1988. 80:166-177.
24. Manske, G.G.B, Ortiz-Monasterio, J.I., van Ginkel, R.M., Rajaram, S. and Vlek P.L.G Phosphorus use efficiency in tall, semi- dwarf and dwarf near-isogenic lines of spring wheat. *Euphytica* .2002. 125, 113–119.
25. Tennant, D. A test of a modified line intersects method of estimating root length. *J. Ecol.* 1975. 63:995-1001.
26. Barton C J. Photometric analysis on phosphate rock. *Ind. Anal. Eng. Chem.* 20.1948.1068–1073.
27. Hammond J.P., Martin R. B., Philip J. W., Graham J. K., Helen C.B., Rory H., Mark C.M., Andrew M., Tracey O., William P. S., and Duncan J.G. Shoot yield drives phosphorus use efficiency in Brassica oleracea and correlates with root architecture traits. *Journal of Experimental Botany.* 2009. 7. (60):1953–1968
28. Moll, R.H., Kamprath, E.J., Jackson, W.A. Analysis and interpretation of factors which contribute to efficiency of nitrogen utilization. *Agronomy J.* 1982. 74: 562-564
29. Ozturk, L., Eker, S., Torum, B and Cakmak, I. Variation in phosphorus efficiency among 73 bread and durum wheat genotypes grown in a phosphorus-deficient calcareous soil. *J. Plant Soil.* 2005. 269:69-80.
30. Mather, K., and Jinks, L. *Biometrical Genetics.* Chapman and Hall. 1982. New York, 396 pp.
31. Gamble E.E. Gene effects in corn. Separation and relative importance of gene effects for yield. *Can J. Plant Sci.* 1962. 42:339-348.
32. Rowek.E, W.L. Alexander. Computations for estimating the genetic parameters in joint-scaling tests. *Crop Sci.* 1980. 20: 109-110
33. Statistical Analysis Institute .SAS User's Guide Statistics, Release 8.2 (TS2 MO), Windows, Edition. Cary, NC: 2000. SAS Institute Inc
34. Ribaut, J.M., Hoisington D.A., Deutsch J.A., Jiang, C and Gonzalez-de- L.D. Identification of quantitative trait loci under drought conditions in tropical maize. I. Flowering parameters and the anthesis-silking interval. *Theor Appl Genet.* 1996. 92: 905-914.

35. Yan D., Keru C., Wan T., Ai Z., Yiping T., Gu F., Zhenling C., Fusuo Z., Xinping C.
Is the Inherent Potential of Maize Roots Efficient for Soil Phosphorus Acquisition?
PLOS one. 2014. 9 (3) e90287. doi:10.1371/journal.pone.0090287
36. Hufnagel, B., Sylvia M.S., Lidianne A., Claudia T. G., Willmar L., Gabriel C. A.,
Barbara N., Brandon G. L., Jon E. S., Maria M.P., Beatriz A. B., Eva W., Frederick
W. R., Joao H. V., Randy T. C., Alexandre F., Rodrigo G., Antonio A.F., Robert E.
S., Leon V. K and Jurandir V. M. Duplicate and Conquer: Multiple Homologs of
PHOSPHORUS-STARVATION TOLERANCE1 enhance Phosphorus Acquisition
and Sorghum Performance on Low-Phosphorus Soils. Plant Physiology.2014.166.
659–677
37. Leiser W.L, Frederick W.R, Eva, W.Z., and Bettina I.G. H. Phosphorus uptake and
use efficiency of diverse west and central African Sorghum under field conditions in
Mali. Plant soil. 2014. 377: pp. 383-394. Doi 10.1007.
38. Hajabbasi, M.A and Schumacher, T.E. Phosphorus effects on root growth and
development in two maize genotypes. Plant and Soil.1994.158: 39-46.
39. Obura, P.A. Effects of soil properties on bioavailability of aluminium and phosphorus
in selected Kenyan and Brazilian soils. PhD thesis .2008. Purdue University, USA.
40. Deng, Y., Keru C., Wan T., Ai Z., Yiping T., Gu, F., Zhenling, C., Fusuo Z., Xinping C.
Is the Inherent Potential of Maize Roots Efficient for Soil Phosphorus Acquisition?
PLOS one .2014. :9 (3) pp 1-9.
41. Monasterio, O.I.J., Pena, R.J., Pfeiffer, W.H., and Hede, A.H. P use efficiency, grain
yield, and quality of triticale and durum wheat under irrigated conditions.
Proceedings of the 5th International Triticale Symposium, Annex. June 30- July 5.
2002. Radzikow, Poland
42. White, PJ and Hammond .JP. Phosphorus nutrition of terrestrial plants. In: White PJ,
Hammond JP, eds. The ecophysiology of plant phosphorus interactions. Dordrecht,
the Netherlands: Springer, .2008. 51–81.
43. Cichy, K.A., Sieglinde, S.S and Matthew, W.B. Plant growth habit, root architecture
traits and tolerance to low soil phosphorus in an Andean bean population.
Euphytica.2009.165 (2), pp 257-258.1007/s10681-008-9778-2
44. Borrero J.C., Pandey S., Ceballos, H., Magnavaca, R and Bahia Filho A.F.C.
(1995).Genetic variances for tolerance to soil acidity in a tropical maize population.
Maydica.1995.40:283-288.
45. Ceccarelli, S.. Specific adaptation and breeding for marginal conditions. Euphytica.
199477:205–219. doi:10.1007/BF02262633
46. Ceccarelli, S.. Adaptation to low/high input cultivation. Euphytica .1996.92:203–214.
doi:10.1007/BF00022846