

Impact of preceding legume and Nitrogen levels on Maize –Preliminary year study

ABSTRACT

Experiment on summer legumes (mungbean, cowpea and sesbania) - maize cropping system with different nitrogen levels were carried out in University of Agriculture, Peshawar during 2011-2012 to determine the immediate and residual effects on soil and plant system. The summer legumes such as mungbean, cowpea and sesbania were sown for grain, fodder and green manure purposes respectively along with fallow as control. After legumes four levels of nitrogen (0, 90, 120 and 150 kg ha⁻¹) were taken for maize. Post-harvest soils of legume experiment results were not significantly affected. However, soil pH was lower in mungbean plots than fallow plots. While on other soil properties viz., Total Soluble Salts, Electrical Conductivity and Calcium Carbonate content were recorded higher in mungbean plots than fallow plots. All the parameters under investigation were significantly affected by nitrogen levels except emergence m⁻² of maize. The effect of legumes was significant for maize grains per ear and ear weight. Taller plants (206 cm) were recorded on 150 kg N ha⁻¹ applied plots and followed by N application of 120 kg ha⁻¹ while control plots resulted in short stature plants (188 cm). Plots fertilized with high N (150 kg ha⁻¹) had significantly higher ear weight which was at par with N application at the rate of 120 kg ha⁻¹. Higher grains per ear were found in ear collected from plots treated with N at the rate of 120 kg ha⁻¹ which was at par with plots fertilized with N at the rate of 150 kg ha⁻¹ and control plots resulted in fewer grains ear⁻¹. The above results suggested that preceding legume plots perform better than fallow plots and it may be concluded that the use of summer legumes in maize cropping system will be a good strategy for sustainability.

Key words: Summer legumes, maize, Cropping systems and nitrogen levels,

I. INTRODUCTION

Cereal based cropping systems are predominantly practiced in Pakistan because of higher productivity and profitability, and for ensuring food security. Previously, balanced fertilization with N, P and K were thought to be essential for achieving high levels of productivity. However, with the passage of time, there is a gradual decline in factor of productivity and increasingly

34 higher doses of nutrients are required to be applied to achieve the potential yields of these crops.
35 This situation is associated with deteriorating fertility of the soils and consequent emergence of
36 multiple nutrient deficiencies including those of micronutrients (Timsina and Connor, 2001).
37 Organic matter is the life of soil, and the practices that support organic matter build-up also favor
38 sustainable productivity (Katyal *et al.*, 2001). Incorporation of plant residues is a useful means to
39 sustain organic matter content and thereby enhance the biological activity, improve soil physical
40 properties and increase nutrient availability (Arif *et al.*, 2014). Organic manuring by way of
41 direct application of conventional farmyard manure, mulching or green manuring, has declined
42 over the years due to various operational constraints (Herridge *et al.*, 1995). Cultivation of
43 legumes for seed, fodder or green manure helps in sustaining the productivity of cereal based
44 cropping systems and improves soil fertility through nutrient cycling (Herridge *et al.*, 1995).

45 In most parts of Pakistan, there is a hot summer period of 70–80 days which occurs from
46 last week of April to mid July, a period between after wheat harvesting and before maize
47 planting. This period can be utilized by short duration legume crops for enhancing grain legumes
48 production or fodder availability or green manuring. Green manuring with annual leguminous
49 crops like Sesbania is a widely known practice has been found beneficial not only for realizing
50 potential yields but also for N economy and improving soil fertility (Singh *et al.*, 1991).
51 Likewise, mungbean as well as cowpea have direct value in human diet and potential in adding
52 N to the cropping system, especially when their residues with nodules are incorporated into the
53 soil (Glasener *et al.*, 2002).

54 It is not only desirable but quite possible to reduce the reliance on synthetic N fertilizers
55 and move towards greater use of legumes to supply N for crop production. In fact, legume-based
56 systems are likely to be more sustainable than cereal-based or fertilizer based systems. There are
57 few studies quantifying the relative contribution of legumes in summer gap towards N economy,
58 productivity, profitability and soil fertility in cereal based cropping system (Arif *et al.*, 2014)

59 Hence, a comprehensive study was undertaken to determine the immediate and residual effects
60 of legumes on soil fertility and maize performance in cereal based cropping system.

61

62

II. MATERIALS AND METHODS

The experiments were conducted at University of Agriculture Peshawar during 2011-2012 in Summer and Kharif season. The experimental farm is located at 34.01° N latitude, 71.35° E longitude, at an altitude of 350 m above sea level in Peshawar valley and has continental type of climate. Soil is clay loam, low in organic matter (0.87 %), phosphorus (6.57 mg kg⁻¹), potassium (121 mg kg⁻¹) and alkaline (pH 8.2) and is calcareous in nature (Amanullah et al., 2009). Summer legumes-maize cropping pattern was used for the experiments. The summer legumes were adjusted in planted during the period between after wheat harvesting and the planting of maize for grain, fodder and green manure purposes. Mungbean for grain, cowpea for fodder, and Sesbania was purely for green manure production. A fallow was included in the experiment as control. After the harvest of wheat crop, summer legumes i.e. cowpea (*Vigna unguiculata*, cv. Ebony), mungbean (*Vigna radiata*, cv. NIAB-2006) and sesbania (*Sebania aculeata*, cv. Pashawari) were sown in the first week of May according to the recommended agronomic practices. The biomass of Sesbania was incorporated into the field with a disc harrow in early July.

After thorough land preparation, each plot from the previous legumes experiment was split into four sub plots to accommodate four levels of N to maize crop. Maize was sown with four levels of N fertilizer (0, 90, 120 and 150 kg ha⁻¹) in mid July. Nitrogen as urea was applied to maize in two equal splits, half at sowing and another half at 30 days of growth. Similarly, the recommended rate of P fertilizer at the rate of 90 kg ha⁻¹ was applied to maize at sowing. For the legume experiment, a randomized complete block design (RCBD) with four replications was used in a plot size of 5 m x 16 m. For maize experiment, the plot (5 m x 16 m) was split into four plots to accommodate four levels of N to maize. In this way, each subplot size was 5 m x 4 m.

Data were recorded on emergence per meter square by counting total number of plants that emerged in one meter row length at three randomly selected rows in each sub plot and then converted to emergence m^{-2} . Plant height of each crop was measured from base to the tip of the plant and five randomly selected plants in each sub-plot were used. Grains per ear of five randomly selected ears was counted and averaged.

For the analysis of soil physio-chemical characteristics, three soil samples were taken from each subplot at random and were mixed together for the determination of different physiochemical properties of soil (pH, EC, total soluble salts (TSS) and CaCO_3). Soil sampling was done after the harvest of legumes.

Statistical analysis

The data were analyzed according to analysis of variance technique appropriate for RCB design for legume experiment and RCB design with split plot arrangement for maize and wheat experiments using Genstat discovery software. The treatment means were compared at $P < 0.05$ level of probability using LSD test (Jan et al., 2009).

IV. RESULTS AND DISCUSSION

Soil parameters after legumes experiment

The values of the soil pH, total soluble salts (TSS), electrical conductivity (EC) and CaCO_3 content were not significantly affected by preceding legumes (Table 1).

Emergence m^{-2}

Statistical analysis of the data showed that nitrogen levels (N) and legumes (L) had non-significant effects on emergence m^{-2} . The N x L interaction was also not significant (Table 2).

Plant height (cm)

Different nitrogen levels significantly affected maize plant height while the effect on legume cropping system remained not significant (Table 2). The Nitrogen x Legumes interaction on maize height was not significant. Plant height increased as nitrogen level increased from 0 to 150 kg ha^{-1} . The tallest maize plants (206 cm) were recorded in plots where N was applied at the rate of 150 kg ha^{-1} (Table 2b) followed by that 120 kg ha^{-1} (202 cm), while control plots recorded the shortest (188cm) maize plants. This increase in plant height could be attributed to positive effect of N on vigorous vegetative growth. The results are in agreement with the finding of Khan *et al.*, (2009) who reported that maize growth increased with increasing nitrogen fertilizer.

Ear weight (g)

Ear weight of maize was significantly affected by nitrogen levels and previously grown legumes (Tables 2a and 2b). Cowpea and mungbean enhanced ear weight of maize than previous sesbania and fallow. The N x L interaction was not significant. Linear increase was observed in ear weight of maize as N level increased. Plots fertilized with high N (150 kg ha^{-1}) had significantly higher ear weight (75 g) which was at par with nitrogen application at

the rate of 120 kg ha⁻¹ as compared to control plots (54 g). Our findings are in line with the observation of Lopez *et al.*, (2007) who reported that nitrogen application has a positive relation with yield of maize. Similarly, ear weight was higher in plots previously sown with cowpea or mungbean. Increase in ear weight is due to effect of preceding legumes which might increase the availability of essential nutrient for plant growth. Similar results are also reported by Sharma *et al.*, (2009) that Sesbania green manuring and legumes in cropping system have a favorable effect on maize yield and yield attributes.

Grains ear⁻¹

Data on grains ear⁻¹ is presented in Table 2. Statistical analysis of the data indicated that nitrogen levels significantly affected grain ear⁻¹ of maize while the effect of legumes were not significant. Similarly, N x L interaction remained not significant. Grains ear⁻¹ increased as N increased from 0 to 120 kg ha⁻¹. Higher grains ear⁻¹ were found in ears collected from plots treated with N at the rate of 120 kg ha⁻¹ which was at par with plots fertilized with N at the rate of 150 kg ha⁻¹. Control plots resulted in fewer grains ear⁻¹. All N amended plots resulted in higher grain ear⁻¹ over control. Timely availability of nutrient mainly N in N amended plots increased in dry matter accumulation and better crop growth that could positively change the physiological functions of the maize crop (Ali *et al.*, 1999).

Table 1. Soil pH, TSS, EC and CaCO₃ as affected by preceding legumes.

Legumes	pH	TSS (%)	EC (dsm ⁻¹)	CaCO ₃ (ppm)
Cowpea	8.200	0.060	0.187	12.833
Mungbean	8.167	0.066	0.207	12.917
Sesbania	8.167	0.059	0.183	13.250
Fallow	8.233	0.053	0.167	12.000
LSD(0.05)	ns	ns	ns	ns

Means in the same column followed by different letters are significantly different from one another at 5% level of probability.

ns = non-significant

Table 2a. Emergence m⁻², plant height (cm), ear weight (g) and grains ear⁻¹ of maize as affected by preceding legumes.

	Emergence m ⁻²	Plant height (cm)	Ear weight (g)	Grains ear ⁻¹
Legumes				
Cowpea	11	205	77a	259 a
Mungbean	11	201	74a	263 a
Sesbania	10	191	63b	242 b
Fallow	11	193	60b	244 b
LSD (0.05)	ns	ns	9.23	13.54

Table 2b. Emergence m⁻², plant height (cm), ear weight (g) and grains ear⁻¹ of maize as affected by different nitrogen levels.

	Emergence m ⁻²	Plant height (cm)	Ear weight (g)	Grains ear ⁻¹
Nitrogen (kg ha⁻¹)				
0	10	188 d	54c	234 c
90	11	195 c	64b	252 b
120	11	202 b	77a	264 a
150	11	206 a	79a	257ab
LSD (0.05)	ns	3.54	4.12	14.68

Means of the same category followed by different letter(s) are significantly different from one another at 5% level of probability.

ns = non-significant

Conclusions and recommendation

It is concluded from the above results that summer legumes increase the availability of soil nutrient, reduce nutrient losses, reduce fertilizer cost and thus have the continuous use of this system will improve the yield of maize on sustainable basis in future.

- 167 Ali S, Naseem WB, -- Ali A. Establishment of productive and sustainable legume-cereal rotations in
168 Pothwar, Pakistan. 1999; Pak. J. Soil Sci. (17): 23-28.
- 169 Ali M. Effect of summer legumes on productivity and nitrogen economy of succeeding rice (*Oryza*
170 *sativa*) in sequential cropping. 1992; Indian J Agric. Sci. 62(7): 466-467.
- 171 Amanullah., R.A. Khattak and S.K. Khalil. 2009b. Effects of plant density and N on phenology and
172 yield of maize. J Plant Nutr 32: 245-259.
- 173 Arif, M. and M.A. Malik. 2014. Economic feasibility of proposed cropping patterns under different
174 moisture regimes of Pothwar plateau. Int. J. Agric. Biol. 11: 27-32.
- 175 Glasener KM, Waggoner MG, Mackown CT, --Richard JV. Contributions of shoot and root nitrogen-
176 15 labeled legume nitrogen sources to a sequence of three cereal crops. 2002; Soil Sci. Soc.
177 Am. J. 66 (2): 523-530.
- 178 Herridge DF, Marcellos H, Felton WL, Truener GL, ---Peoples MB. Chickpea increases soil-N
179 fertility in cereal systems through nitrate sparing and N₂ fixation. 1995; Soil Biol. Biochem.
180 27(3): 545-551.
- 181 Jan M T, Shah P, Hollington PA, Khan MJ, ---Sohail Q. Agriculture Research: Design and
182 Analysis, A Monograph. 2009; NWFP Agricultural University Peshawar, Pakistan.
- 183 Katyal JCN, Rao H, Reddy MN. Critical aspects of organic matter management in the tropics. The
184 Example of India. 2001; Nutr. Cycl. Agro-ecosyst. 61(4): 77-88.
- 185 Khan A, Jan MT, Marwat KB, --Arif M. Organic and inorganic nitrogen treatment effect on plant
186 and yield attributes of maize in a different tillage system. 2009; Pak. J. Bot. 41(6): 99-108.
- 187 López-Bellido, L., R.J. López-Bellido and R. Redondo. 2007. Nitrogen efficiency in wheat under
188 rainfed Mediterranean conditions as affected by split nitrogen application. Field Crops
189 Res. 94: 86-97.

190 Ojiem JO, Vanlauwe B, de Ridder N, --Giller KE. Niche based assessment of contributions of
 191 legumes to the nitrogen economy of Western Kenya smallholder farms. 2007; Plant.Soil.
 192 292(5): 119–135.

193 Safdar A, Schwenke GD, Peoples MB, Scott GF, --Herridge DF. Nitrogen, yield and economic
 194 benefits of summer legumes for wheat production in rainfed northern Pakistan. 2002;
 195 Pakistan J. Agron., 1: 15–19.

196 Sharma AR, ---Behera UK. Nitrogen contribution through Sesbania green manure and dual-purpose
 197 legumes in maize-wheat cropping system: agronomic and economic considerations. 2009;
 198 Plant and Soil J. 325(5): 289-304

199 Singh Y, Khind CS, --Singh B. Efficient management of leguminous green manures in wetland rice.
 200 1991; Adv. Agron. 45(1): 135-189.

201 Timsina, J., M.A. Quayyum, D.J. Connor, M. Saleque, F. Haq, G.M. Panaullah, M. Jahan and
 202 R.A. Begum. 2006. Effect of fertilizer and mungbean residue management on total
 203 productivity, soil fertility and N-use efficiency in intensified rice–wheat systems. Int. J.
 204 Agric. Res. 1(1): 41-52.