



The effect of Jasmonic Acid (JA) as seed treatment and soil drench on **Some** Morphological parameters of Moneymaker tomato (*Solanum lycopersicum* L.)

ABSTRACT

The aim of this research was determine the effect of Jasmonic Acid (JA) application on some morphological parameters of money maker tomatoe. This research was undertaken in the botanical garden of the department of Conservation, Lancaster university, United kingdom. Results showed that Tomato plants that received 3mM JA seed treatment showed a reduction in their mean height and are significantly different ($p < 0.001$) to control soil treated plants. Tomato plants of JA soil drench treatment also showed a significant reduction in their mean height ($p = 0.001$) compared to the control soil drenched plants. JA seed primed Moneymaker tomato plants showed significant increase in their mean root weight compared to the control at ($p = 0.026$). However, there were no significant differences in mean dry weight between JA soil drenched plants and their controls at ($p = 0.110$). Although, significant increase in mean root weight was found in soil-drenched plants compared to control seed treated plants ($p = 0.002$),. Increase in total mean dry weight of Moneymaker tomato plant (*Solanum lycopersicum* L.) was detected in those of JA seed treatment (figure 5, $p = 0.018$) compared to control seed treatment (Tukey's *post hoc* test). Similarly, there is significant increase in mean dry weight of JA soil drench plants than the control soil drenched plants (figure 5, $p = 0.001$) using Tukey's *post hoc* test. Finally it was also observed that no growth cost was associated with priming by exogenous JA application except for stunting of plant height.

Keywords: jasmonic Acid, seed treatment, morphological parameters, soil drench, money maker

Introduction

In inducible defence, plants are known to possess three key signaling hormones, which are Jasmonic acid (JA), Salicylic acid (SA) and Ethylene (ET). Each of these phytohormones has a dependent pathway that aids plants in battling particular class of pathogens (Abd El Rahman *et al.*, 2012).



In the wild, plants may be challenged with different forms of pathogenic **and non-pathogenic** invasions at the same time (Koornneef and Pieterse, 2008). This as a result, builds- up the innate defense processes within the plant (inducible defence) against these simultaneous attacks. The plant makes use of defence mechanisms that are either destructive to all aggressors, or at least does not impede resistance to other attackers.

The main methods of exogenous chemical application for inducing prime state in plant includes soil drenching, leaf spraying and seed treatment. However, their usefulness are limited as they tend to reduce plant growth and reproduction in the course of triggering strong plant defence (Zhang *et al.* 2003;). However, as described by Ahmad *et al* (2010), exogenous chemical application may guarantee efficient and more reliable priming reactions in plants with insignificant cost on growth using appropriate concentrations.

In addition, Worrall *et al* (2012), observed no growth cost as JA seed treated tomato plants with sustained priming response effective against *B. cinerea* and other necrotrophic pathogens, and was recommended for field trials.

Leaf and root drench were familiar methods of priming plants by means of synthetic chemicals. Though it was recently discovered that priming plant defence by seed treatment is though very effective, it is unknown if its resistance, particularly to necrotrophic pathogens is mediated by hydrogen peroxide.

However, it is the intention of this research to focus on the Effect of Jasmonic Acid (JA) seed treatment and soil drench on Some Morphological parameters of Moneymaker tomato (*Solanum lycopersicum* L.)

Materials and Methods

Seed treatment

Seeds of Moneymaker tomato cultivar (*Solanum lycopersicum* L.) of were supplied from moles seeds (www.molesseed.co.uk). The procedures for the treated seeds were carried out as reported by Worrall *et al.*, (2012). Approximately 100 seeds of each cultivar were soaked in 10ml aqueous solution containing 25 µl of 3mM JA (prepared from 1.19 molar (M) stock in absolute

ethanol) at low temperature of 4°C for a period of 24 hours. For control, equal numbers of seeds were soaked in 10ml aqueous solution containing 25 µl of 100% ethanol. All the incubated seeds were put through three rounds of washing with deionized water, for approximately 15 minutes and sown in compost.

Plant growing conditions and soil drench

Seeds of Moneymaker tomato cultivar (*Solanum lycopersicum* L.) were grown within the Lancaster Environment Centre glasshouse, which as programmed has low and high temperature ranges of $20 \pm 4^\circ\text{C}$ and $30 \pm 4^\circ\text{C}$ respectively. The 600W Phillips Plantaster light bulbs in the glasshouse supplied $200 \pm 50\mu\text{mol m}^{-2}\text{s}^{-1}$ of photosynthetic photon flux at 14h photoperiod to supplement low natural photoperiods. The seeds were sown in 0.3L pots filled with high nutrient compost (Levington M3), and seeds were watered two times a day (10am and 4pm) from a hose connected tap within the glasshouse. For soil- drench treatment, untreated seeds were sown directly into pots and watered, until seedlings emerged and grew to second leaf stages. A 0.1mM JA concentration was prepared from the 1.19M JA stock and administered to soils at a volume of 30mls per pot. Control seedlings were soil-treated with 30mls of 0.09% aqueous ethanol solution per pot. At 3 weeks of age, the plants were repotted alongside the seed treated plants into 0.9L pots containing new compost. However, roots of soil – drenched seedlings were carefully rinsed to remove compost that may contain chemicals before repotting. This was to avoid any further influence or effect the chemicals might have on subsequent plant growth. Growth parameters were measured at fourth leave stage

Results

The effect of exogenous application of JA by seed treatment and soil drenching on height of Moneymaker tomato (*Solanum lycopersicum* L.) is showed in figure 1 below. Tomato plants that received 3mM JA seed treatment showed a reduction in their mean height and are significantly different ($p < 0.001$) to control soil treated plants . Also, tomato plants of JA soil drench treatment showed a significant reduction in their mean height ($p = 0.001$) compared to the control soil drenched plants (Tukey's *post hoc* test)

JA seed primed Moneymaker tomato plants showed significant increase in their mean root weight compared to the control (Figure 2, $p=0.026$). However, there was no significant difference in mean dry weight between JA soil drenched plants and their controls (Figure 2, $p=0.110$). Although, significant increase in mean root weight was found in soil-drenched plants compared to control seed treated plants (Figure 2, $p=0.002$), (Tukey's *post hoc* test).

Tukey's *post hoc* test revealed no significant difference in mean stem dry weight between 3mM JA seed primed plants and ethanol seed treated plants (control) (figure 3, $p=0.388$). However, JA soil drenched plants showed significant increase in their mean dry stem weight compared to that of the control soil drenched plants (figure 3, $p=0.034$) (Tukey's *post hoc* test).

Plants of JA seed treatment did not show significant difference in their mean leaf weight (figure 4, $p=0.129$) compared to the control, but a significant increase was found in JA soil drenched plants than the control soil drenched plants (figure 4, $p=0.011$). Also, JA soil drenched plants showed a significant increase in their mean dry weight compared to those of control seed treatment (figure 4, $p=0.026$).

Increase in total mean dry weight of Moneymaker tomato plant (*Solanum lycopersicum* L.) was detected in those of JA seed treatment (figure 5, $p=0.018$) compared to control seed treatment (Tukey's *post hoc* test). Similarly, there is significant increase in mean dry weight of JA soil drench plants than the control soil drenched plants (figure 5, $p=0.001$) using Tukey's *post hoc* test.

Discussion

The data showed a significant reduction in height of JA seed treated plants and JA soil drench plants compared to their controls. This stunting is possibly the consequence of suppressing other phytohormone pathways by exogenous JA application for priming defense against necrotrophic

pathogens, which includes *B. cinerea* (Denance *et al.*, 2013). However, the data showed increase in stem dry weights and leaf dry weight of the primed plants. This is equivalent to the findings of Sorial *et al.*, (2010), which showed how Jasmonic acid application on sweet basil, resulted to a reduction in the plant height but increased the stem and leaf dry weight of the plant. The data showed that the increase in all the mean dry weights of all the measured primed plant parts were only statistically significant in JA soil- drench plants, except for the root weight when compared to control. This indicates that 30 application JA as soil- drench might have inhibited root growth Maciejewska and Kopcewicz, (2003) showed that exogenous JA or MeJA application on Japanese morning glory tends to reduce its root growth, and further increase in the Jasmonate concentration further increases the root reduction of the plant. In contrast, the data showed that JA seed treated plants showed increase in mean dry measurements, which were not statistically significant except for root dry weights. This further suggests that the significant root dry weight of JA seed treated plant compared to the control might be due to absence of direct contact between the plant hormone and the plant root.

In addition, a significant increase in total mean plant dry weight was observed in both JA seed treated and JA soil drenched plants. This might be because the exogenous application JA have induced some physical changes that led to increase in the dry weight of the primed plants such as increase in the number of trichomes per unit surface of the plant (Boughton, 2005). Link to the fact that JA application is known to increase trichome as though to inhibit the movement of the attacking herbivore on the plant (War *et al.*, 2013). The increase in dry weight could also be due some defense responses that led to cell wall strengthening. Ahmad *et al.*, (2013) also showed how exogenous application of plant growth regulators like SA, and JA at early stage of maize plant (*Zea mays*) development gave notable improvement in its subsequent growth stages.

136




Figure 1. Mean plant height of 4 weeks old Moneymaker tomato plant (*Solanum lycopersicum* L.) of 3mM JA seed treatment, control seed treatment, JA soil drench or control soil drench. The measurement was carried out using a 100cm wooden ruler. Values show means of 10 plant replicates for each treatment and vertical bars represent ± 1 SE.

146

147

148

149

150

Root dry weight




Figure 2. Mean root dry weight of 4 weeks old Moneymaker tomato plants (*Solanum lycopersicum* L.) of 3mM JA seed treatment, control seed treatment, JA soil drench or control soil drench. The measurement was carried out using a weighing balance. Values show means of 10 replicate plants for each treatment and vertical bars represent ± 1 SE.




Figure 3. Mean stem dry weight of 4 weeks old Moneymaker tomato plants (*Solanum lycopersicum* L.) of 3mM JA seed treatment, control seed treatment, JA soil drench or control

soil drench. The measurement was carried out using weighing balance. Values show means of 10 plant replicates for each treatment and vertical bars represent ± 1 SE.




Figure 4. Mean leaf dry weight of 4 weeks old Moneymaker tomato plants (*Solanum lycopersicum* L.) of 3mM JA seed treatment, control seed treatment, JA soil drench or control soil drench. The measurement was carried out using a weighing balance. Values show mean replicates of 10 plants for each treatment and vertical bars represent ± 1 SE.

Plant dry weight




Figure 5.

Mean plant dry weight of 4 weeks old Moneymaker tomato plants (*Solanum lycopersicum* L.) of 3mM JA seed treatment, control seed treatment, JA soil drench or control soil drench. The measurement was carried out using weighing balance. Values are mean replicates of 10 plants for each treatment and vertical bars represent ± 1 SE.

Conclusion

It was observed that Jasmonic Acid (JA) at 3mM application by seed treatment or soil drench increased the most of the growth parameters studied, except in plant height where there was decrease observed leading to the stunting of growth.



References.

- Abd El Rahman T., El Oirdi M., Gonzalez-Lamothe R., Bouarab K. (2012). Necrotrophic pathogens use the salicylic acid signaling pathway to promote disease development in tomato. *Molecular Plant- Microbe Interactions*, **25**: 1584-1593.
- Ahanger R., A., Bhat H., A., Bhat T., A., Ganie S., A., Lone A., A., Wani I., A., Gania S., A., Haq S., Khan O., A., Junaid J., N., Bhat T., A. (2013). Impact of climate change on plant disease. *International journal of Modern Plant and Animal Sciences*, **1**: 105-115.

- Ahmad I., Shahzad M., A., Afzal I., Farooq M., Wahid A. (2013), Growth improvement in spring maize through exogenous application of ascorbic acid, salicylic acid and hydrogen peroxide. *International Journal of Agriculture and Biology*, **15**: 95-100.
- Ahmad S., Gordon-Weeks R., Pickett J., Ton J. (2010). Natural variation in priming of basal resistance: from evolutionary origin to agricultural exploitation. *Molecular Plant Pathology*, **11**: 817–827.
- Baarlen V., A., N., Woltering P., Staats E., J., M., (2007), Histochemical and genetic analysis of host and non-host interactions of Arabidopsis with three Botrytis species: an important role for cell death control. *Molecular Plant Pathology*, **8**:41-54.
- Bari R., Jones J., D., G. (2009). The role of plant hormones in plant defence. *Plant Molecular Biology*, **69**:473–48.
- Bechtold U., S., Karpinski S., Mullineaux P., M. (2005). The influence of the light environment and photosynthesis on oxidative signalling responses in plant-biotrophic pathogen interactions. *Plant Cell And Environment*, **28**: 1046-1055.
- Behdani M., Pooyan M., Abbasi S. (2012). Evaluation of antifungal activity of some medicinal plants essential oils against *B. cinerea*, causal agent of postharvest apple rot, in vitro” *International Journal of Agriculture and Crop Sciences*, **14**: 1012-1016.
- Boughton A., J., Hoover K., Felton G., W., (2005), Methyl-jasmonate application induces increased densities of glandular trichomes on tomato, *Lycopersicum esculentum*. *Journal of Chemical Ecology*, **31**: 2211–2216.
- Denance N., Sanchez-Vallet A., Goffner D., Molina A. (2013). Disease resistance or growth: the role of plant hormones in balancing immune responses and fitness cost. *Frontiers in Plant Sciences*, **155**: 1-12.
- Koornneef A., Pieterse C., M., J. (2008). Cross talk in defense signaling. *Plant Physiology*, **146**: 839–844.
- Maciejewska B., Kopcewicz J. (2003). Inhibitory effect of methyl jasmonate on flowering and elongation growth in *Pharbitis nil*. *Journal of Plant Growth Regulation*, **21**: 216-223.
- Sorial M., E., El- Gamal S., M., Gendy A., A. (2010). Response of sweet basil to jasmonic acid application in relation to different water supply. *Bioscience Research*, **7**: 39-47.
- War A., R., Hussain B., Sharma H., C. (2013). Induced resistance in groundnut by jasmonic acid and salicylic acid through alteration of trichome density and oviposition by *Helicoverpa armigera* (Lepidoptera: Noctuidae). *AOB Plants*, **5** : plt053; doi: 10.1093/aobpla/plt053.

- 244 Worrall D., Geoff H., Moore J., P., Glowacz M., Croft P., Taylor J., E., Paul N., D.,Roberts M.,
245 R. (2012). Treating seeds wit activators of plant defense generates longlasting priming
246 resistance to pests and pathogens. *NewPhytologist*, **193**: 770-778.
247
- 248 Zhang Y., Chen K., Zhang S., Ferguson I. (2003). The role of salicylic acid in postharvest
249 ripening of kiwifruit. *Postharvest Biological Technology*, **28**: 67–74.
250