

Original Research Article

STUDIES ON HETEROSIS FOR HEAT TOLERANCE IN TOMATO

ABSTRACT

An investigation was carried out with a view to characterizing, evaluating and estimating heterosis for yield and ^{its} yield attributes under high temperature conditions at the experimental field of Horticulture Department, Faculty of Agriculture during May to August 2013. The better parent heterosis was estimated for ten characters. Considering better parent heterosis, the cross C41  P5 (203.22%) exhibited highest positive heterosis for fruit yield plant⁻¹ and lowest from the hybrid of C51  C1 (32.61%). Again the hybrid C41  C51 showed the highest heterobiotic effect of 0.72 for days to first flower. Out of 10 hybrids, C51  C71 showed the highest positive heterosis (85.12) on number of fruits plant⁻¹. The cross FP5  C71 exhibited the highest significant and positive heterosis (69.31) on individual fruit weight which indicate suitable combinations for improving fruit weight. The positive and the highest heterosis for pericarp thickness was recorded from the cross combination C11  C71(56.25), while the cross combination of C51  C11 and C51  71 exhibit the positive and significant heterosis for the length and fruit width, respectively.

 **Keywords:** Hybrid, heterosis, tomato, *Solanum lycopersicon* L.

INTRODUCTION

Tomato (*Solanum lycopersicon* L.) belongs to the family Solanaceae having the chromosome number of 2n= 24. It is one of the most popular and nutritious vegetable crop all over the world including Bangladesh (Ahmed, 2002). It is a self-fertilized annual crop. Tomato is grown in winter in Bangladesh, as the temperature is congenial at that period for optimum growth and yield. But it has great potentiality to grow in summer also. Due to its palatability and vitamin content its demand remains high throughout the year, while its production is too low than requirements especially in summer season. In summer' availability of fresh tomato is less due to agro-ecological constraints. Cultivation of summer tomato are affected due to excessive rainfall, temperature, wind storm, etc. during the monsoon season. Therefore, special thrust is needed to increase summer vegetables production and thereby also increase cash flow to the farm family. Heterosis or hybrid vigour is an important biological phenomenon refers to the manifested superiority of the F1 hybrid resulting from the cross of genetically dissimilar homozygous parents. Now days, heterosis breeding is one of the most efficient tool to exploit the heterotic response for several traits. Many hybrids of the crops like bean, cucumber, eggplant, onion, watermelon,

cabbage, and radish are frequently grown on large scale (Singh *et al.*, Bhatt *et al.*, 1999). Substantial amount of heterosis for many traits in tomato have been reported by several authors. Hence, attempt can be made to exploit heterosis for better yield and quality of tomato. Though heterosis for a number of economic characteristics of tomato have been reported by several researchers, but its feasibility and utilization for commercial production is still lacking which need to be initiated in this country. Recently Bangladesh Agricultural Research Institute (BARI) has strengthened the programme for year round tomato variety development and already succeeded to develop some heat tolerant hybrids (Anon., 2003). But research is continuous process and attempt can be made to develop new hybrid combinations which might be superior over the existing hybrid varieties. Therefore, present investigation was to undertaken to study the heterosis for yield and yield attributes.

44

45 MATERIALS AND METHODS

46 Description of the Study Area

47 The experiment was conducted at the experimental field of Horticulture Department, Sylhet
48 Agricultural University, Bangladesh during May to August 2013. It is located 275 km
49 north-east of Dhaka. The study area is located on latitude 24°N to 33.67° and longitude
50 91°54' to 95.88° E. The soil of experimental site belongs to the AEZ-20: Eaturan Surma
51 Kushiyara Floodplain. Soil was brown hill soil in texture and highly acidic (pH4.83) in
52 nature. The climate of the experimental site was subtropical characterized by heavy rainfall
53 during May to October and scanty during rest of the year. During the experimental period
54 the monthly average rainfall (cm) and air temperature (°C) from May to August were 34.2
55 cm, 26.9 cm, 18.9 cm and 26.55°C, 29.85°C, 29.59°C, 29.1°C. The mean monthly relative
56 humidity ranged from 46 to 80% with the peak occurring between July and August.

57 Experimental Design and Planting

58 Seeds of ten cross combinations of tomato were selected in this study. Seeds were sown in
59 three different seed beds on May 15, 2013. After seed sowing the seed bed was given
60 partial shade with black net. The seed bed was also covered by a fine white net (60 × 60
61 mesh) to protect seedlings from insect attack. Eight days old seedlings were transplanted in
62 polythene bag for hardening of the seedlings. Twenty five days old seedlings were
63 transplanted in the experimental field on June 10, 2013. The experiment was laid out in
64 Randomized Complete Block (RCB) Design with three replications. The unit bed was made

almost 15 cm in height with a view to prevent the crop from moist condition during heavy rain. The land was previously cultivated with county bean. The unit plot size was 2.3 m × 4.8 m which was again divided into two beds and each bed there were 12 plants row⁻¹ and 24 plants plot⁻¹. Plants were spaced at 60 cm × 40 cm between plant to plant and row to row, respectively. The unit plot and blocks were separated by 50 cm and 75 cm drain, respectively. The land was fertilized with 15 t well decomposed cowdung 300 kg urea, 200 kg TSP and 150 kg MoP hectare⁻¹, respectively. Half of the quantity of cow dung and the entire amount of TSP were applied during land preparation. The remaining cowdung and half of MoP were applied before 5 days of planting. The whole of urea and half of MP were applied in 3 equal splits as top dressing at 15, 30 and 50 days after transplanting. Light irrigation of each seedling was given just after transplanting for better establishment of the plants. To protect the crop from heavy rain polytunnel was provided over the crop during entire cropping period. The height of the polytunnel at the middle of the bed was 6.0 feet while it was 4.5 feet at both sides. The structure of the polytunnel was made of bamboo where transparent polythene used over the structure. Stacking was provided to keep the plant erect. Weeding, irrigation, ~~P~~^pruning, mulching and other intercultural operations were done as and when necessary.

82

83 **Estimation of heterosis**

84 For estimation of heterosis in each character the mean values of 10 F₁'s have been
85 compared with better parent (BP) for heterobeltosis. Percent heterosis was calculated as –

86 For better parent heterosis,

$$\begin{aligned} & (F_1 - BP) \\ 88 \quad H (\%) &= \frac{\text{-----}}{BP} \times 100 \\ 89 \end{aligned}$$

90 Where, F₁ = Mean of F₁ generation

91 BP = Mean of better parent

92 The significance test for heterosis was done by using the following formulae:

93 Critical difference (CD) for heterosis over better parent

$$94 \quad CD = \sqrt{2MSE/r} \times t$$

95 Where, MSE= Error mean square

96 r = Number of replication

97 t = Tabulated value of t at error df at 1 % and 5% level of significance

98 CD was used to compare the observed differences among different treatments. If the
99 difference was greater than CD value, it was considered as significant or vice-versa.

100 Results and discussion

101 Analysis variance (Table 1) revealed that the days to first harvesting, the estimated
102 heterosis over better parent revealed that out of 10 cross combinations negative heterosis
103 for days to first harvesting was not found. Among the 10 cross combinations, 5 hybrids
104 showed significant and positive heterosis for days to first flowering (Table 2). The highest
105 positive heterosis for this trait was observed in the cross combination C41 $\times$ C71 (11.67%).
106 Heterobeltiotic for this trait ranged from 0.72% to 11.67%. In case of harvesting, the
107 estimated heterosis over better parent revealed that out of 10 cross combinations negative
108 heterosis was found in C41 $\times$ C51. ^(-3.03%). ~~The negative heterosis observed in the cross~~
109 ~~combination C41 $\times$ C51 (-3.03%).~~ Heterobeltosis for this trait ranged from -3.03% to
110 7.70%. There was no significant negative heterosis, 8 hybrids showed the positive heterosis
111 for this traits. Heterosis for this trait ranged from 0% to 85.12%. The highest positive
112 heterosis was found in cross C51 $\times$ C71 (85.12%). Shoudhury and Khanna (1972), Jamwal
113 *et al.* (1984) and Legon *et al.* (1984) also reported heterobeltosis for tomato fruits per plant
114 under high temperature growing environment. Nine cross combinations out of 10 cross
115 combinations exhibited significant positive heterotic value for fruit weight over better
116 parent. Heterosis for this trait ranged from 2.25% to 69.31%. The highest significant
117 positive heterosis was found in cross combination FP5 $\times$ C71 (69.31%). Heterosis for this
118 trait ranged from 28.26% to 203.22%. The highest positive heterosis was found in cross
119 combination C41 $\times$ FP5 (203.22%). Shoudhury and Khanna (1972) reported 73.77% and
120 Sidhu and Singh (1993) reported 71.7% heterobeltosis for yield per plant in tomato.

121
122
123
124

125 **Table 1. Analysis of variance for hybrids on different characters of tomato**

Source of variation	DF	Days to first flowering	Days to harvest	No. of fruits plant ⁻¹	Individual fruit weight	Fruit length	Fruit width	TSS (%)	Locule fruit ⁻¹	Pericarp thickness
Replication	2	0.62	5.49	15.96	9.21	0.04	0.04	0.04	0.02	0.01
Genotype	14	11.93**	10.16**	122.97**	207.60**	0.91**	1.02**	0.72**	2.07**	0.06**
Error	28	1.81	1.58	0.24	1.93	0.02	0.01	0.03	0.36	0.01

126

127 **Table 2 Percent heterosis over better parent of 10 tomato cross combinations for**
128 **different characters**

129

Hybrids	Days to first flowering	Days to harvest	No. of fruits plant ⁻¹	Individual fruit weight	Pericarp thickness
C41 × C51	0.72	-3.03	46.08**	24.03**	13.33
C41 × C11	4.38	4.45	67.10**	45.24**	43.76**
C41 × FP5	8.76**	5.27	59.22**	29.78**	26.66
C41 × C71	11.67**	7.70	38.80**	46.30**	60**
C51 × C11	2.17	1.14	0.43	2.25	23.52
C51 × FP5	1.45	3.24	43.60**	18.51**	17.70
C51 × C71	4.28	2.84	85.12**	55.23**	23.52
C11 × FP5	8.76**	2.03	0	25.76**	18.75
C11 × C71	10.08**	7.28	17.28**	64.84**	56.25**
FP5 × C71	10.95**	2.03	13.46**	69.31**	46.16
CD 5%	2.25	2.1	0.52	2.32	0.13
CD 1%	3.03	2.83	1.10	3.13	0.17

130

** indicates significant at 1% level of probability.

131 In case of fruit length, eight cross combinations showed significant negative heterosis.

132 Heterosis for this trait ranged from -36.36% to 6.70%. The only positive heterosis was
133 found in cross combination C51 × C11 (6.70%). In case of fruit width, the cross

134 combination C51 × FP5 and C51 × C71 showed significant positive heterosis (4.02%) and ^{one} !

135 cross combinations out of 10 cross combinations showed significant negative heterosis.

136 Heterobeltosis for this trait ranged from -26.10% to 5.31%. Heterobeltosis for total soluble

solids revealed that only one cross combinations (C41 × FP5) out of 10 cross combinations showed significant negative heterosis. Heterosis for this trait ranged from -35.78% to 8.70%. The highest negative heterosis (-35.78%) was found in cross combination C41 × C11 × FP5 (8.7%) and FP5 × C71 (2.01%) showed positive heterosis over FP5 and the two cross combinations, positive better parent heterosis for TSS was observed in two crosses. It was C11 × FP5 (8.7%) and FP5 × C71 (2.01%). Heterobeltosis for locules per fruit, the estimated heterosis over better parent revealed that 3 cross combinations out of 10 crosses showed significant negative heterosis. Heterosis for this trait ranged from -33.33% to 16.67%. The highest negative heterosis (-33.33%) was found in cross combination C41 × FP5 and C41 × C71. Heterobeltosis for pericarp thickness of fruit, the estimates of heterosis over better parent revealed that 3 cross combinations out of 10 crosses showed significant positive heterosis. Heterosis for this trait ranged from 13.33% to 60%. The highest positive heterosis (60%) was found in cross combination C41 × C71 followed by C11 × C71 (56.25%).

Table 3. Percent heterosis over better parent of 10 tomato cross combinations for fruit characteristics

Hybrids	Fruit length	Fruit width	TSS (%)	Locule fruit ⁻¹	Fruit yield plant ⁻¹
C41 × C51	-17.65**	-2.52	0.79	-33.32	183.33**
C41 × C11	-16.34**	5.31	-1.25	-22.23*	141.30**
C41 × FP5	-25.69**	-17.07	-35.78**	-33.33*	203.22**
C41 × C71	-17.64**	-1.64	0*	-33.33*	97.06**
C51 × C11	6.70**	-4.018*	2.03	16.67	32.61**
C51 × FP5	-4.54	4.018*	-0.61	8.32	119.44**
C51 × C71	-21.27**	4.018*	2.01	-8.32	155.55**
C11 × FP5	-6.14*	-26.10*	8.7**	-22.23	28.26**
C11 × C71	-21.27**	-26.10**	1.41	-22.23	93.48**
FP5 × C71	-36.36**	-24.39**	2.01*	-11.10	100**
CD 5%	0.23	0.17	0.27	0.99	0.007
CD 1%	0.311	0.225	0.36	1.33	0.101

** indicates significant at 1% level of probability, * indicates significant at 5% level of probability

154 Considering above results in discussion, the hybrid combinations, C41 × FP5, C41 × C51
155 and then for further evolution alas with available heat tolerant variety to recommend form
156 level production.

157 REFERENCES

- 158 Ahmed SU. Genetics of fruit set in Tomato (*Lycopersicon esculentum* L) under hot-humid
159 condition. Ph. D. Thesis. Dept. of Hort, Bangabandhu Sheikh Mujibur Rahman
160 Agriculture University, Gazipur. 2002; pp. 1-236.
- 161 Anonymous. Developed vegetables varieties and technologies. Olericulture Division, HRC,
162 BARI, Joydebpur, Gazipur. 2003; pp. 25-38.
- 163
- 164 Bhatt RP, Biswas VR and Kumar N. Studies of heterosis for certain characters in tomato
165 (*Lycopersicon esculentum* Mill.) under mid hill condition. Progressive Hort. 1999;
166 31(1-2):41-43.
- 167 Chaudhury RC and Khanna RK. Exploitation of heterosis in tomato yield and components.
168 South Indian Hort.1972; 20: 59-65.
- 169 El Ahmadi A and Stevens MA. Reproductive responses of heat-tolerant tomatoes to high
170 temperatures. J. Amer. Soc. Hort. Sci. 1979; 104:686-691.
- 171 Jamwal RS, Rattan RS and Saini SS. Hybrid vigour and combining ability in tomato. South
172 Indian Hort. 1984; 32(2): 69-74.
- 173 Legon MC, Diaz N and Garcia PG. Performance of tomato hybrids and their parents in
174 summer. Centro Agricola. 1984; 11(1): 35-44.
- 175 Rashid MM. Shabji Baggan(In Bengali). Bangla Academy, Dhaka. 1993; pp. 192-219.
- 176 Sidhu AS and Singh S. Studies on heterosis and divergence in tomato. Plant breeding
177 abstracts. 1993; pp. 018-064.