

Effect of Nitrogen Fertilizer Rates on Panicle Length and Yield Component of Some Improved Rice Varieties (*Oryza sativa* L.)

Abstract

Rice (*Oryza sativa*) is a principal staple food crop in Kenya. However, its production is still low due to inherently low and declining soil fertility. This has resulted into food and nutritional insecurity and low living standards. The situation has been compounded by the ever escalating fertilizer prices which has made it unaffordable to most small holder farmers. Although some studies have been done on fertilizer application on some rice varieties, the Mwea upland rice (MWUR) varieties were bred under low fertilizer input environment while other authors have indicated that the New Rice for Africa (NERICA) varieties give high yields under low input conditions. The objective of the research was to establish the yield performance of eight promising improved rice varieties under four nitrogen (N) levels and identify the variety that gives high yields at relatively low N fertilizer rates. Field experiments were carried out at Alupe in Western Kenya under rainfed upland condition. The experiment layout was split plot factorial in a Randomized Complete Block Design with three replicates. The main plot treatments were four rates of nitrogen fertilizer which were; 0 (control), 40 kg ha⁻¹, 80 kg ha⁻¹ and 120 kg ha⁻¹ applied as urea in two equal splits, sub-plots consisted of four MWUR and four NERICA varieties. The parameters measured included panicle length, yield at 14% moisture content, filled grain ratio percentage and yield components. The measured parameters increased significantly with increase in the level of nitrogen fertilizer. Varietal difference was significant and MWUR varieties recorded higher panicle length and yield component at low nitrogen levels (0N and 40N) as compared to the NERICAs with MWUR 1 and 2 recording higher values. The NERICAs out-yielded the MWUR varieties at higher nitrogen levels. NERICA 4 recorded highest yield among the NERICA varieties regardless of the N level. Results from our study suggested that MWUR 1 and 2 and NERICA 4 were more tolerant to low nitrogen soil as compared to MWUR 3 and 4 and NERICA 1, 10 and 11, because of higher panicle length, higher filled grain ratio percentage and higher yield component and may be suitable for soils deficient in nitrogen.

Keywords: *Oryza sativa* L., Nitrogen, Panicle length, yield components

Introduction

Rice (*Oryza sativa* L.) is the most important food crop, being the primary food source for more than one third of the world's population and is grown in 11% of the world's cultivated area. Upland rice has high production potential in Kenya and can be grown by most farmers at low input level [1]. Kenya has a potential to bring 1,000,000 ha under upland rice production [2]. However, currently the area under rainfed rice is 13,000 ha [3] hence the national rice development strategy (NRDS) focus is expanding and increasing yields by exploiting existing opportunities on a sustainable basis [4]. Nitrogen (N) is one of the most important nutrients for plant growth and a major factor that limits agricultural yields [5,6]. It is required in adequate amount at early, mid tillering, panicle initiation and at ripening stage for better grain development. It has played a major role in the global food production over the past 60 years [7]. Nitrogen (N) requirements of rice crop are met from both the soil and fertilizers. Usually due to acute N deficiency in most rice soils, fertilizer N must be applied to meet the crop demand [8] and yet excessive N application would lead to increased production cost and negative physiological effects due to excessive growth. This might block agricultural sustainability and cause environmental pollution.

The amount of N required and management of the N varies depending on variety, soil conditions, cultural practices, crop rotations, and other factors. The majority of smallholder farmers who form the bulk of agriculture in Kenya are faced with low soil fertility problems. Rice yields decline due to low input system, for example, rice yields of less than 2.0 tons/ha against a potential of 5.0 tons/ha [3] has been reported. The yield gap is attributed mainly to effects of drought and low soil fertility, the most prominent of which is nitrogen deficiency. As a result, rice farmers are unable to harness the potential of rainfed rice to meet their food and nutritional demand. Lack of identification of improved varieties adaptable to low fertility soils in rainfed conditions and the ever escalating fertilizer prices have made this important input unaffordable to most small-holder farmers who are resource poor hence low rice production [9]. However, some

varieties can perform well under low nitrogen input [10]. There is therefore need to evaluate the response of the MWUR viz a viz the NERICA varieties to identify the superior rice varieties that are adaptable to low nitrogen levels. Identification of low nitrogen adaptable rice varieties is thus crucial in sustaining production in areas where low soil fertility prevails.

[11] reported that nitrogen is the most important nutrient that limit crop production. They argued that this loss can be mitigated by exploiting varieties well adapted to low soil N conditions in order to increase farm yields. According to [12], selection of the most appropriate level of nitrogen fertilization is a major concern of economic viability of crop production and the impact of agriculture. Given the importance of nitrogen fertilization on the yield in grain from the rice plant, it is necessary to know what the best dose is for each variety as well as its influence on components of yield. Yield studies that will help in identifying varieties that have either specific or general adaptation to low nitrogen which can be exploited for varietal recommendation hence address the soil fertility problem in the region. The identified low nitrogen adaptable varieties would have a substantial impact on rice productivity in terms of soil nutrient supply, nutrient use efficiency, improved food security and livelihood/income of rural households in the region. Therefore advanced farming systems research can contribute to the nutritional well-being and food security of the communities. Incorporating a high yielding variety, which is tolerant to low soil nitrogen is important. Studies have shown that application of N fertilizer to rice leads to increase in panicle length, spikelet number, and grain yield [10,13]. However, the response of rice varieties to increasing N levels has not been studied or documented in Kenya currently. This information is of importance in any rice breeding programme in Kenya because there is little knowledge on the response of improved rice varieties to increasing levels of N fertilizer [9]. The main objective of this project was to establish and document data on the yield performance of eight promising high-yielding improved rice varieties to various nitrogen levels as well as to identify the variety that gives high yields at relatively low N rates. These objectives were set against the hypotheses that improved rice varieties in Kenya are not responsive to increasing N rates as the existing NERICA varieties and that they are not low nitrogen use efficient.

MATERIALS AND METHODS

Site Description

Field studies were conducted to assess the effect of nitrogen fertilizer rates on Panicle length and yield component of eight improved rice varieties under rainfed upland conditions between August 2012 and April 2013 at Alupe farm of Lake Basin Development Authority (LBDA) in Western Kenya. Geographically, the experimental area is located at latitude 0° 30' 0 N; longitude 34° 7' 50 E and at the elevation of the 1170 m above sea level. Precipitation ranged between 680.5 – 860 mm and temperature ranged from 16-34°C during the growing period. The soils at Alupe have been characterized as Ferralo-orthic Acrisol with pH of 5.0 as determined in H₂O (1:1) [14]. Specific soil chemical characteristics of the location were determined according to the analytical procedures of [15] and are presented in Table 1. Soil samples were collected at depth of 20 cm using a soil auger from all corners of the plot and between diagonals and the middle part. This was done at the start of the experiment to assess the initial chemical status of soil in each of the plots. The soil was thoroughly mixed together to form a composite sample which was then air dried and passed through 2.0 mm sieve for soil texture and sub samples ground through 0.5 mm sieve for chemical analyses. The field was ploughed and harrowed to a suitable tilth before sowing.

Experimental Design and Treatments

The experiment layout was split plot factorial arrangement in a Randomized Complete Block Design (RCBD) with three replicates. The main plot was fertilizer N source and different rice varieties as sub-plots. The main plot treatments were four rates of nitrogen fertilizer which were; 0 (control), 40 kg ha⁻¹, 80 kg ha⁻¹ and 120 kg ha⁻¹ applied as urea in two equal splits i.e. 14 days after sowing (DAS) and at panicle initiation (46 DAS) as recommended by [16]. Sub-plots consisted of eight rice varieties i.e. four MWUR rice varieties namely Mwea upland rice 1, 2, 3 and 4 (MWUR1-4) that were bred and selected by KARI-Mwea under low fertilizer input environment and four New Rice for Africa (NERICA) varieties namely 1, 4, 10 and 11 that have been released to farmers in Kenya. Drill sowing was done in 5 m x 1.2 m plots, 20 cm between rows and within rows at a depth of 3 cm and four seeds per hill. The seedlings were thinned

to 1 plant per hill 15 days after sowing. In between the plots, trenches of 0.5 m depth and 1 m in width were constructed, to avoid seepage between plots. A basal dose of triple super phosphate (TSP) was used as the source of phosphorus, applied at the rate of 30 kg P ha⁻¹. The fertilizer granules were placed at the bottom of the hole and covered with about 1 cm of soil before sowing the seeds to avoid fertilizer burn to the rice seeds during dissolution of triple superphosphate. The rice seeds were covered with roughly 2 cm of soil. The sowing date coincided with onset of the long rainy season. Normal cultural trial practices such as weeding and bird scaring were carried out.

Table 1. Chemical characteristics of soils at LBDA Alupe farm (0 - 20 cm depth)

Soil chemical properties	
pH _{1:1 soil:water}	4.69
ECe (Electrical conductivity)	0.18
Total N g kg ⁻¹	0.73
Available P (ppm)	0.14
Extractable-Zn (ppm)	1.75
Extractable-Cu (ppm)	5.66
Extractable-Mn (ppm)	48
Extractable-Fe (ppm)	479

Sampling and Measurements

Panicle lengths

This was determined using a metre rule at physiological maturity. Measurements were done from the panicle base to the tip of five hills in each plot.

Yield components

At physiological maturity, a 1 m⁻² quadrat was placed in each plot leaving out the border rows and border plants. The plants within the quadrats were harvested to determine yield components and rice grain yield.

The number of grains per 5 g, and filled grains per panicle were determined and yield calculated according to [1]. The yield was extrapolated in kilograms per hectare as follows:

$$\text{Yield (Kg ha}^{-1}\text{)} = \text{Number of panicles m}^{-2} \times \text{Number of grains/panicle} \times \% \text{ filled grains} \times (\text{Weight of 1,000 grains} \div 1,000) \times 10,000$$

Yield at 14% Moisture Content

This was determined from an area of 1m^{-2} by measuring the moisture content of the grains immediately after harvesting using a grain moisture tester (model number AF 34086, Japan) and converting the yield of the grains to 14% moisture content and extrapolating to Kg ha^{-1} as described by [1].

Filled Grain Ratio Percentage

Grains harvested from the 1 m^{-2} in all the plots were put in different buckets of water where poorly filled grains together with empty grains floated, while the well filled grains settled at the bottom. Filled grains were then separated from empty and poorly filled grains. The grains were dried and later weighed but each was handled separately. The percentage of filled grains was calculated according to [1].

Statistical data analysis

Factorial analysis of variance (ANOVA) was carried out on the data for the variables measured to determine the significance of the effects of nitrogen fertilizer levels on the rice varieties using a statistical computer package (SAS software version 9.2) [17]. Significant means were separated using least significance difference test (LSD) at 5% level for comparing the treatment means. Linear regression curve was used to show and define the nature and strength of the relationships between nitrogen levels, rice varieties and grain yield.

RESULTS

Panicle Length

Nitrogen levels had a highly significant effect on panicle length. Increase in nitrogen level caused a significant increase in panicle length ($P \leq 0.05$). The plants subjected to 0N had the shortest panicle

lengths (Figure 1). Varietal differences were also significant ($P \leq 0.05$). At 40 N, MWUR 1 had significantly higher panicle length than MWUR 2, 3 and 4 and the NERICAs ($P \leq 0.05$). NERICA 4 recorded significantly higher panicle length at 0N as compared to NERICA 1, 10 and 11. At 80N, MWUR 1 and NERICA 4 recorded slightly higher panicle length as compared to NERICA 1, 10, 11, MWUR 2 and 4. However, there was significant difference in panicle length between MWUR 3 and NERICA 1, 10, 11, MWUR 2 and 4. There was no significant interaction between the varieties and treatments ($P > 0.05$).




Fig. 1. Effect of N rate on panicle length of eight rice varieties grown at four levels of nitrogen treatment (Means of three replicates $\pm$ SE). LSD (0.05) $V = 1.06$, $T = 0.75$.

Yield at 14% Moisture Content

Nitrogen application showed a highly significant ($P \leq 0.05$) effect on yield at 14% moisture content. Plants treated with 120N and 80N recorded significantly ($P \leq 0.05$) higher yield at 14% moisture content than plants treated with 40N and 0N. The varietal difference was also significant ($P \leq 0.05$) with MWUR 1 and 2 registering significantly higher values at 0N as compared to MWUR 3, 4 and the NERICAs. At 40N, MWUR 1 and 2 recorded significantly higher yield at 14% moisture content compared to the NERICAs however, the difference in yield

was not statistically significant in MWUR 3 and 4. The results also revealed a significant variety and treatment interaction.




Fig. 2. Effect of N rate on yield at 14% moisture content of eight rice varieties grown at four levels of nitrogen treatment (Means of three replicates $\pm$ SE). LSD (0.05) V = 15.67, T = 11.07.

Filled Grain Ratio (%)

Generally nitrogen application caused an increase in filled grain ratio percentage in all the NERICAs and MWURs rice varieties. Plants fertilized with 120N and 80N had significantly higher filled grain ratio percentage than plants treated with 0N and 40N ($P \leq 0.05$). The varietal difference was highly significant ($P \leq 0.05$). MWUR 1 and 2 recorded significantly ($P \leq 0.05$) higher filled grain ratio percentage at 0N and 40N compared to MWUR 4 and the NERICAs. At 80N and 120N, NERICA 4, 10 and 11 had slightly higher filled grain ratio percentage compared to MWUR varieties. There was a significant ($P \leq 0.05$) interaction between variety and nitrogen level.




Fig. 3. Effect of N rate on filled grain ratio (%) of eight rice varieties grown at four levels of nitrogen treatment (Means of three replicates $\pm$ SE). LSD (0.05) V = 1.2, T = 0.86.

Yield Component

Yield component significantly ($P \leq 0.05$) increased with the increase in nitrogen rate in all the NERICAs and MWURs rice varieties with plants subjected to 120N recording the highest yield while plants subjected to 0N recorded the lowest yield. The varietal difference was significant ($P \leq 0.05$) with MWUR 1 having significantly higher yield component at 0N and 40N compared to MWUR 3, 4 and NERICA 1, 4, 10 and 11. Generally MWUR varieties registered higher yield components at 0N and 40N, while NERICAs varieties recorded higher yields at 80N and 120N. NERICA 4 had significantly ($P \leq 0.05$) higher yield component at 0N in comparison to NERICA1, 10 and 11. There was a positive and significant correlation between yield component and filled grain ratio percentage (Figure 5). Yield component and panicle length also recorded a positive and significant correlation (Figure 6).




Fig. 4. Effect of N rate on yield component of eight rice varieties grown at four levels of nitrogen treatment (Means of three replicates $\pm$ SE). LSD (0.05) V = 99.68, T = 70.48.




Fig.5. Relationship between yield component and field grain ratio (%) of rice as affected by different varieties and nitrogen levels




Fig.6. Relationship between yield component and panicle length of rice as affected by different varieties and nitrogen levels.

Discussion

The results showed that panicle length was significantly influenced by different nitrogen application treatments. Plants fertilized with higher nitrogen levels recorded significantly higher panicle lengths than plants fertilized with lower N levels. These results are in line with those reported by [18,19,20]. The longer panicles obtained in treatments receiving higher nitrogen rates were probably due to better nitrogen status of plant during panicle growth period because nitrogen takes part in panicle formation as well as panicle elongation and for this reason, panicle length increases with the increase of N-fertilization. When absorbed during the reproductive and ripening phases, nitrogen promotes development of the panicle and stimulates nutrient absorption and assimilation as evidenced by an increase in size and number of filled grain ratio percentage [21]. According to [22], the increase in panicle length due to adequate N nutrition is explainable in terms of possible increase in nutrient mining capacity of plant as a result of better root development and increased translocation of carbohydrates from source to growing points in well-fertilized plots. The varietal difference was also significant with MWUR 1 recording higher panicle length at low nitrogen levels compared to other rice varieties. This may imply that MWUR 1 is tolerant to low nitrogen and its cell division and elongation is not greatly affected by nitrogen deficit as compared to MWUR 2, 3, 4 and NERICA 1,4,10 and 11. NERICA 4 registered higher panicle length regardless of the nitrogen level as compared to NERICA 1, 10 and 11. This might imply that among the NERICA varieties under study, NERICA 4 is tolerant to low nitrogen conditions in the soil but can also thrive under adequate nitrogen levels.

Application of N fertilizers significantly increased field grain ratio percentage and yield at 14% moisture content. Field grain ratio percentage and yield at 14% moisture content increased in all the rice varieties with an increase in nitrogen level. These results are in accordance with those of [13,23,24]. The increase in yield at 14% moisture content with increasing N rates may be due to the high number of panicles per square meter, which is significantly associated with high grain yield in rice [23]. According to [13], the improvement in yield owing to the application of N-fertilizers might be brought about by the beneficial effect of nitrogen on nutrient uptake and physiological growth. Nitrogen nutrition influences the content of photosynthetic pigments, the synthesis of the enzymes involved in the carbon reduction and the formation of the membrane system of chloroplasts, thus the increase in growth and yield owing to the application of

N-fertilizers may be attributed to the fact that these nutrients being important constituents of nucleotides, proteins, chlorophyll and enzymes, are involved in various metabolic processes which have direct impact on vegetative and reproductive phases. Varietal difference was significant with MWUR rice varieties having higher filled grain ratio percentage and higher yield at 14% moisture content at low nitrogen levels as compared to NERICA varieties, while the NERICA varieties registered higher values at higher nitrogen levels. This may imply that the MWUR rice varieties are tolerant to low nitrogen and may out-yield the NERICAs under low nitrogen soils while the NERICAs might perform better than the MWUR varieties under optimal nitrogen levels.

Nitrogen fertilizer application had a statistically significant effect on the yield component of the studied rice varieties. There was an increase in yield component with increase in each successive level of nitrogen fertilizers. Similar results were also reported by [12,25,26]. The higher yield component under high nitrogen levels might be attributed to the vigorous and healthy growth and development of more productive tillers and leaves ensuring greater resource utilization as compared to plant that were deficient in N. Increases in yield components are associated with better nutrition, plant growth and increased nutrient uptake [27]. The crop at no nitrogen level produced the lowest grain yield which was significantly lower than all other nitrogen levels tested. This might be due to lack of supply of nitrogen into the soil to meet the required nutrients for physiological processes, which in turn lowered the yield. According to [28], the application of nitrogen improves various crop parameters like panicle length, more productive tillers and grain yield thus resulting in higher yields. There was a significant positive relationship of yield component with panicle length and filled grain ratio percentage in all the varieties. This implies that panicle length is a significant yield component for rice yield. This high capacity for sink formation was associated with their higher capacity for N acquisition and biomass production. Panicle length contributes to increment of rice grain yield indirectly by increasing the number of panicles per m² and number of spikelets per panicle

Varietal difference was significant and MWUR varieties recording superior yield components at low nitrogen levels (0N and 40N) as compared to the NERICAs with MWUR 1 and 2 recording higher values

while the NERICAs out-yielded the MWUR varieties at higher nitrogen (80N and 120N). NERICA 4 and 11 recorded higher yield components at low nitrogen level as compared to the other NERICA varieties. The results may imply that the MWUR varieties are tolerant to low nitrogen as compared to the NERICAs and that MWUR 1 and 2 are more adaptable to low nitrogen as compared to MWUR 3, 4 and the NERICA varieties. NERICA 4 may be more adaptable to low nitrogen as compared to the other NERICA varieties that were under study. MWUR varieties were bred as low-management rice plant types targeted for resource-limited, smallholder production systems and their performance at 0N and 40N thus confirms the significant role they may play in African smallholder systems. The observed greater yield in NERICA 4 compared to NERICA 1, 10 and 11 at 0N and 40N could be attributed to its higher potential for panicle production and greater ability to partition photosynthates more efficiently to the grain as indicated by higher filled grain ratio percentage, compared to the other varieties. According to results by [10,13] NERICA 4 was tolerant to low N because of the greater yield observed especially at zero and low N. This was attributed to its high potential for panicle production and ability to partition photosynthates more efficiently to the grain as indicated by a higher harvest index compared with the other cultivars. [29] reported that NERICA 4 produced significantly higher yield and was better weed competitor than CG 14, ITA 150, WAB 56-104, and the other NERICA cultivars (NERICA 1, 10, and 11) in the dry savannas of Nigeria.

Conclusion

This study has shown that nitrogen levels affected panicle length and yield components of all the MWURs and NERICAs varieties studied. There was appreciable varietal difference in response to low soil nitrogen conditions. Generally MWUR varieties studied were more adaptable to low nitrogen conditions as compared to NERICA varieties. The study therefore accepts the hypothesis that MWUR varieties which had been bred to low-N management are well responsive to low N environments and are good for resource-limited, smallholder production systems. They exhibit superior yielding capacity under low nitrogen fertilized soils. They may be important in the small-holder farming systems in Kenya. The NERICA varieties recorded higher yield at high nitrogen levels as compared to MWUR varieties, though,

NERICA 4 gave higher yield as compared to other NERICA varieties regardless of the nitrogen level. The decision on the appropriate N rates would depend on economic conditions of the farmer. NERICA 4 would fit in both low and high input fertilizer farming systems. The study suggest that MWUR 1 and 2 and NERICA 4 were more tolerant to low nitrogen as compared to MWUR 3 and 4 and NERICA 1, 10 and 11 because of higher panicle length, higher filled grain ratio percentage and higher yield component and may be suitable for soils deficient in nitrogen.

References

1. Ministry of Agriculture (MOA). Guidelines to rice production in Kenya. Agricultural information resource centre, Nairobi; 2011.
2. Sikuku PA, Onyango JC and Netondo GW. Yield component and gas exchange responses of NERICA rice varieties (*Oryza sativa* L.) to vegetative and reproductive stage water deficit. Global Journal of Science Frontier Research 2012;7(3): 49 - 59.
3. Ministry of Agriculture (MOA). Economic Review of Agriculture. Central Planning and Project Monitoring Unit. Ministry of Agriculture, Government of Kenya; 2010.
4. Rosemary, A.E., Bibiana, M.W., Njuguna, N., Dominic, M.K., Daniel, A. Rice value chain, study report of Kenya. Ministry of Agriculture and Kenya Agricultural research Institute. 2010;36-37.
5. Dai XQ, Zhang HY, Spiertz JH, Yu J, Xie GH, Bouman B. Crop response of aerobic rice and winter wheat to nitrogen, phosphorus and potassium in a double cropping system. Nutrient cycling in Agroecosystems. 2009;86(3):301-315.
6. Xia L, Zhiwei S, Lei J, Chenggan R, Wang M and Chuangen L. High/low nitrogen adapted hybrid rice cultivars and their physiological responses. African Journal of Biotechnology 2012;10(19):3731-3738.
7. Simonne E, Liu G, Gazula A, Hochmuth B, Landrum L, Gast D, Davis L, Laughlin W, Randal R, Bauer M, Vann C, Saft C and Toro E.. Yield response of overhead irrigated snap bean to nitrogen rates. *Proc. Fla. State Hort. Soc.* 2012;125:174-181.
8. Choudhury AT and Kennedy IR. Nitrogen Fertilizer Losses from Rice Soils and Control of Environmental Pollution Problems. Soil Science and Plant Analysis. 2005;36:1625–1639.

- 336 9. Kimani JM, Tongoona P, Derere J and Nyende A. Rice grain yield improvement under low soil
337 phosphorus and nitrogen conditions for two Mega-environments. African Crop Science Conference
338 Proceedings. 2011;10:587-597.
- 339 10. Oikeh S, Toure A, Sidibe A, Niang A, Semon M, Sokei Y and Mariko S. Response of upland NERICA
340 rice cultivars to nitrogen and plant density. Archives of Agronomy and Soil Science 2009;55:301-314.
- 341 11. Liu D and Shi Y. Effects of different nitrogen fertilizer on quality and yield in winter wheat. Advance
342 Journal of food Science and Technology. 2013;5(5):646-649.
- 343 12. Islam MS, Hasanuzzaman M, Rokouzzamanc M and Nahar K. Effect of split application of nitrogen
344 fertilizer on morpho-physiological parameters of rice genotypes. International Journal of Plant Production.
345 2009;3(1): 52-61.
- 346 13. Kamara AY, Ekeleme F, Omoigui L, Oikeh OS, Chikoye D and Tegbaru A. Responses of upland rice
347 cultivars to nitrogen fertilizer in the Savannas of Nigeria. Agronomy. 2010;102(1):333-339.
- 348 14. Fertilizer Use Recommendation Project (FURP). Final report annex1: fertilizer trial documentation
349 (ferdoc). Ministry of Agriculture, Nairobi, Kenya;1987.
- 350 15. International Institute of Tropical Agriculture (IITA) Automated and semi automated methods for soil
351 and plant analysis. Manual series No. 7 IITA Ibadan Nigeria;1989.
- 352 16. Oikeh SO, Nwilene F, Diatta S, Osineme O, Toure A, Okeleye K.. Responses of upland NERICA rice
353 to nitrogen and Phosphorus in forest agroecosystems. Agronomy. 2008;100(3):735-741.
- 354 17. SAS Institute. SAS/STAT Users guide. Release 9.1. ed. SAS institute Inc. Cary, NC, USA; 2003.
- 355 18. Haque MA, Khan H, Haque ME, Islam MS and Islam MF. Effect of nitrogen application on
356 morphological characters and yield attributes of fine *aman* rice. Agroforestry and Environment. 2012;6(1):
357 67-70.
- 358 19. Rafey A, Khan PA and Srivastava VC. Effect of N on growth, yield and nutrient uptake of upland rice.
359 Indian Journal of Agronomy. 1989; 34(1);133-135.
- 360 20. Ali A, Choudhry MA, Malik MA and Ahmad R. Effect of various doses of nitrogen on the growth and
361 yield of two wheat cultivar. Pakistan Journal of Biological Science. 2000;3(6): 1004-1005.

21. Awan TH, Ali RI, Manzoor Z, Ahmad M and Akhtar M. Effect of different nitrogen levels and row spacing on the performance of newly evolved medium grain rice variety, KSK-133. *Animal & Plant Sciences*. 2011;21(2): 231-234.
22. Pramanik K, and Bera AK. Effect of Seedling Age and Nitrogen Fertilizer on Growth, Chlorophyll Content, Yield and Economics of Hybrid Rice (*Oryza sativa* L.). *International Journal of Agronomy and Plant Production*. 2013;4(5): 3489-3499.
23. Fageria NK and Baligar VC. Lowland response to nitrogen fertilization. *Soil Science and Plant Analysis* 2001;32:1405-1429.
24. Meena SL, Surendra S, Shivay YS and Singh S. Response of hybrid rice (*Oryza sativa*) to nitrogen and potassium application in sandy clay loam soils. *Indian Journal of Agricultural Science*. 2003;73(1): 8-11.
25. Bhuyian LR, Islam N, Mowla G. Rice response to N application with different irrigation schedules. *Rice Research Newsletter*. 1990;15(6): 18.
26. Hach CV and Nam NH. Responses of some promising high-yielding rice varieties to nitrogen fertilizer. *Omonrice*. 2006;14: 78-91.
27. Thakur RB. Performance of summer rice (*Oryza sativa*) to varying levels of nitrogen. *Indian Journal of Agronomy*. 1993;38(2):187–190.
28. Chaturvedi I, Effect of nitrogen fertilizers on growth, yield and quality of hybrid rice (*Oryza sativa* L.). *Central European agricultural* 2005;6(4):611-618.
29. Ekeleme F, Kamara AY, Oikeh SO, Omoigui LO, Amaza P and Chikoye D. Response of upland rice cultivars to weed competition in the savannas of West Africa. *Crop Protection*. 2009;28:90-96.