

Original Research Article

Growth Response to Nutrient Management during a 3 Year Study of *Miscanthus x giganteus*

Abstract

To accelerate the acclimation of novel crops for bio-energy feedstock supply chain, a pilot study focused on *Miscanthus Giganteus* was conducted in the piedmont area of North Carolina to determine region specific productivity. The main aim of the study was to determine the growth response in biomass accumulation during a 3 year establishment period under different nutrient management practices. Growth response was tested through nutrient application at (5) different fertilizer rates: 0 NPK kg ha⁻¹ (T₁), 67 NPK kg ha⁻¹, (T₂), 135 NPK kg ha⁻¹ (T₃), 202 NPK kg ha⁻¹ (T₄), and 269 NPK kg ha⁻¹ (T₅). The experimental study took place at NC A&T State University research farm located in Greensboro, NC (Guilford County). “Nutrient Management” *P* (<.0001/<.0001), “Harvest Year” *P* (<.0001/<.0001), and the “Interaction Effect” *P* (0.0002/0.01) were significant factors affecting (*Fresh/Dry*) matter accumulation observed during 3 consecutive years. Fresh matter was shown to dramatically increase in biomass accumulation with fertilizer treatment T₃ during 2013 (17.57±1.74 t ha⁻¹), 2014 (38.51±0.8 t ha⁻¹), and 2015 (45.43±2.91 t ha⁻¹) harvests as compared to the control treatment T₁. Dry matter followed a similar trend in which yield was shown to significantly increase at treatment application T₃ during 2014 (23.54±0.8 t ha⁻¹), and 2015 (36.15±3.05 t ha⁻¹), as compared to the dry matter yields recovered from treatment application T₁ during 2014 (11.02±1.6 t ha⁻¹), and 2015 (26.76±0.64 t ha⁻¹). Treatment T₃ has produced significantly higher biomass than T₁ & T₂.

Keywords: Biomass, Giant Miscanthus, Growth Response, North Carolina.

Abbreviations: (DOY) Day of Year, (FY) Fresh Yield, (DY) Dry Yield.

Highlights

- *Miscanthus Giganteus* is well adapted to the climatic conditions of the piedmont region of N.C.
- Hyper accumulation of above ground biomass can be achieved in a short period under resource constraints.
- There is a positive growth response due to nutrient application at different rates of NPK.

1. Introduction

Miscanthus is native to tropical and sub-tropical regions of Africa and southern Asia [1]. *Miscanthus* has a unique advantage over traditional C₃ biomass crops (wheat, rice, and potatoes) due to the enhanced capacity of the C₄ photosynthetic pathway which produces large quantities of biomass [2]. Primarily, 3 species are referred to high biomass yielders for *Miscanthus*; *M. giganteus*, *M. sacchariflorus*, and *M. sinensis* [3]. Our research goal was to explore the productivity and resources requirement to produce a viable, non-invasive, high yielding grass species *M. giganteus*. *M. giganteus* used in this study is a genetically altered crop which is produced as a sterile hybrid between *M. Sinensis* and *M. Sacchariflorus* with characterization of being a low input, high biomass yielding, rhizomatous perennial grass, with a growing interest in bio-materials industry.

Miscanthus is an undomesticated crop which under optimized production practices could increase productivity [4]. Potential by-products from pyrolysis of residues could produce bio-liquid, gas, and biochar [5]. In 2007, an estimated 10,000ha of *Miscanthus* biomass was dedicated for heat generation in power stations in the UK [6]. If marketability of *Miscanthus* continues to expand economic competitiveness could equal that of traditional crops cultivated on arable land [7]. By-products from *Miscanthus* biomass could increase its production value in bio-refinery (chemical, manufacturing and power generation) systems. *Miscanthus* has been sold as horse bedding due to high absorbency and bio-degradability characteristics [8]. Another alternative use is as a bio-degradation product in mushroom cultivation [9]. Lastly, by-products recovered through *Miscanthus* operations could help increase dimensional stability to improve the quality of fiberboard [10].

To increase economic viability of *Miscanthus* biomass production, growers focus on nutrient management strategies which increase yields. However, excessive application of fertilizers leads to environmental pollution and reduced profit margins. *Miscanthus* has gained interest over the past few years due to low input requirement, 3m year⁻¹ growth rate, with estimated Drymatter Yield (DY) between 20-25 t ha⁻¹. [11]. Average lifespan of well-established *Miscanthus* is estimated between 15–20 years [12]. However, maturity is reached in the third year with DY estimated between 10-30 t ha⁻¹ at autumn harvest [13]. Increased biomass accumulation during the first six years of establishment has been noted however, noticeable reductions were observed after seven consecutive production years [6]. DY of over 37 t ha⁻¹ during first year establishment have also been reported [14]. Nitrogen loss from fertilizer application has been reported to be comparable to cereal production [15]. Studies conducted have shown that to produce 1.5 kg m⁻² above ground DY it would require 9.2, 1.3 and 20.4 g m⁻² of N-P-K [16]. Parallel and, stable growth responses to nutrient application were not found for *M. Giganteus* in previously conducted experiments.

A significant increase in biomass production has been noted when N application reached upwards to 110 kg ha⁻¹ of N [13]. A (62%) increase in additional biomass accumulation was reported when fertilizer rates increased from 100 to 200 kg N ha⁻¹ [14]. However, results have been reported of smaller incremental DY increase of only (0.05%) when N fertilizer application was increased from 0 to 180 kg ha⁻¹ of N [17]. DY estimates vary dramatically between

consecutive annual harvest's, reports of first year biomass was estimated between (0.7- 2 t ha⁻¹), second year (7.9-15.5 t ha⁻¹) and in the third production year ranged between (17.4-24.5 t ha⁻¹) [18]. Few studies have been conducted on *M. Giganteus* in North Carolina. Little available information on the production of *M. x Giganteus* using a blanket application of N-P-K was found during the literature review.

1.1 Objectives

- 1.) Assessment of growth response of *M. x giganteus* under differential nutrient applications.
- 2.) Determine biomass productivity response between annual harvests during three consecutive harvest years.

2. Methodology

2.1 Experimental Site and Trial establishment

The study was conducted on an Enon Sandy Clay Loam (Taxonomic class: Fine, mixed, active, thermic Ultic Hapludalfs) with (2-4%) slopes and (6-10%) slopes, located in Greensboro, North Carolina (North Carolina Agricultural and Technical State University Research Farm, coordinates: latitude: 36.06733°, longitude: -79.73447, altitude 223m. The current climate hardiness zone is classified as zone 7_B.

2.2 Site Preparation and Nutrient Management

Miscanthus variety "Freedom" rootstock was obtained from REPREEVE Renewables, LCC in Greensboro, USA and was grown using five different fertilizer treatments. The experimental plot layout was a 2x5 factorial design replicated four times. A total of twenty 4m x 12m plots were established with 3m alley. Furrows were opened using a plow and rhizomes were placed 0.60m apart and covered with soil. Five different fertilizer rates (0 NPK kg ha⁻¹ (T₁), 67 NPK kg ha⁻¹ (T₂), 135 NPK kg ha⁻¹ (T₃), 202 NPK kg ha⁻¹ (T₄), and 269 NPK kg ha⁻¹ (T₅)) were Fertilizer was applied in split applications through broadcasting at pre-determined amounts of 17-17-17 N-P-K (N-P₂O₅-K₂O) fertilizer. Pest management was done by applying Biceps in (2013), and a (Fungicide) Diathane was applied in (2014); weeds were manually removed during each growing season to reduce competition pressure. Major phenological developments are listed in (Table 1).

Table 1. Provides a summary of principal cultivation practices and major developmental phases. Day of Year (DOY), and Growing Degree Days (GDD) both production seasons. Asterix indicates non- quantifiable parameter. Growing degree days was calculated as GDD= [(Temp max°C + Temp min°C)/2]. Thermal Base Temperature (-10°C) was determined by a previously determined formula [19].

Event	2013		2014		2015	
	DOY	GDD	DOY	GDD	DOY	GDD
Planting Date:	147	*	*	*	*	*
Germination Date:	170	303	99	3	86	9
Herbicide Application:	163	211	99	3	*	*

Fertilizer Application ₁ :	164	227	126	157	118	155
Fertilizer Application ₂ :	196	681	198	1,114	146	424
Harvest Date:	326	1,835	329	2,185	26	2,318
Tasseling Started:	266	1,597	255	2,005	268	2,393
Vegetative State Ended:	304	1,835	298	2,206	331	2,452
Predominately Brown Color:	312	1,846	322	2,204	352	2,468

111

112 2.3 Biomass Harvesting and Sub-Sample Quantification Measurements

113 Plants were harvested leaving stubs approximately 15 cm from the soil surface. Biomass from 1
 114 M² quadrant area was collected from each plot from a randomly chosen location, and the entire
 115 sample was immediately weighed on a Mettler Toledo SB16000 scale. To reduce analysis
 116 sensitivity to edge effects, samples were not taken adjacent to outside rows or located within
 117 close proximity to borders. A subsample (6 stalks) was collected from each quadrant to measure
 118 moisture content. Fresh biomass samples were dried at 70°C to determine the moisture content of
 119 each sample, using a Fischer thermotemp oven over a 24hrs duration. Moisture content was
 120 determined and expressed as a percentage of fresh weight [20].

121 2.4 Statistical Analysis

122 Growth response was analyzed using PROC MIXED (Mixed Procedure, SAS v9.2, SAS
 123 Institute, Cary, NC, U.S.A.), using (4) replications with (5) fertilizer rates “Nutrient
 124 Management”, and (3) harvest years “Harvest Year” as fixed effects with the corresponding
 125 interaction between “Nutrient Management” x “Harvest Years”.

126 3. Results

127 *Miscanthus* biomass production was studied for 3 growing seasons from 2013-2015 and the data
 128 was tabulated for fresh weight, moisture loss and dry weight. Yield data and statistical analysis
 129 are presented in the tables below.

130 Table 2.) Results of Fit Statistics and Type 3 Tests of Fixed Effects using PROC MIXED for
 131 biomass accumulation characteristics of the individual main effects “Nutrient Management” and
 132 “Harvest Window” and the Interaction Effect for *Miscanthus* collected annually for 3
 133 consecutive years. Comparisons for means were pooled across “Nutrient Management” and
 134 “Harvest Window”.

Type 3 Tests of Fixed Effects		
Effect	Fresh Matter Accumulation	Dry Matter Accumulation
Block	0.40	0.47
Nutrient Management	<.0001**	<.0001**
Harvest Year	<.0001**	<.0001**
Nutrient Management x Harvest Year	0.0002*	0.01*
-2 Res Log Likelihood	153.9	141.6
AIC	155.9	143.6
AICC	156.1	143.8

BIC 157.2 144.9

†=* indicates a significant deviation between Main Effects “Nutrient Management” and “Harvest Year” at 0.001, and ** indicates highly significant deviation 0.0001 between aforementioned pairwise comparison.

To determine the optimum dose of fertilizer to aid in efficient nutrient management guidelines for *Miscanthus*, we evaluated a range of fertilizer applications (T₂, T₃, T₄, and T₅) with a control treatment T₁ which received no nutrient application. During the first establishment year we found that pairwise comparisons between treatment applications T₃, T₄, and T₅ produced a significant increase in fresh matter yield compared to the control treatment T₁ (Table 3). In the following growing season (2014) the same trend in growth response was evident, however, during (2015) only treatment comparisons T₃ and T₅ produced significantly greater fresh matter yields compared to the control treatment (T₁) (Table 3). When evaluating yield response between treatment plots and observed fresh matter yields produced we found that pairwise comparisons between treatment applications T₂ and T₃ did not differ significantly during the first production year (2013), however, during (2014), and (2015) treatment T₃ produced significantly higher fresh matter yields as opposed to treatment T₂ (Table 3). When evaluating treatment application T₄ to yields achieved by treatment application T₃ we observed significantly comparable fresh matter yields retrieved during the first and second growing season (2013) (Table 3). However, in latter production years (2015) treatment comparisons T₄ and T₃ were significantly different in fresh matter accumulation in which treatment T₃ produced significantly higher fresh matter yield accumulation (Table 3). Yield response between treatment applications T₄ compared to T₅ produced comparable yields during the first two production years (2013-2014), however, yields were significantly higher for treatment application T₅ during the 2015 growing season (Table 3).

Table 3.) Differences of pairwise comparisons using Tukey Post Hoc Analysis for Fresh Matter Accumulation of the interaction effect between “Nutrient Management” and “Harvest Year” for *Miscanthus* collected during the 2013, 2014, and 2015 growing seasons. Individual Nutrient Management yields obtained at each fertilizer rate during each individual harvest year.

Pairwise Comparisons	2013 (<i>p</i> Value)	2014 (<i>p</i> Value)	2015 (<i>p</i> Value)
T ₁ vs.T ₂	0.49	0.002**	0.06
T ₁ vs.T ₃	0.03*	<.0001**	<.0001**
T ₁ vs.T ₄	0.04*	<.0001**	0.36
T ₁ vs.T ₅	0.03*	<.0001**	0.0007**
T ₂ vs.T ₃	0.14	<.0001**	0.002**
T ₂ vs.T ₄	0.17	0.0001**	0.32
T ₂ vs.T ₅	0.14	0.01**	0.06
T ₃ vs.T ₄	0.91	0.06	0.0002**
T ₃ vs.T ₅	0.99	0.001**	0.17
T ₄ vs.T ₅	0.91	0.10	0.007**
Nutrient Management	2013 (t ha ⁻¹)	2014 (t ha ⁻¹)	2015 (t ha ⁻¹)
T ₁	12.52±1.14	16.24±2.39	33.55±0.74
T ₂	13.70±1.19	23.87±2.27	37.91±4.22

T ₃	17.57±1.74	38.51±0.80	45.43±2.91
T ₄	17.32±2.74	33.98±1.28	36.32±2.63
T ₅	16.84±0.03	30.86±2.51	42.29±3.39

†=* indicates a significant deviation between Main Effects “Nutrient Management” and “Harvest Year” at 0.001, and ** indicates highly significant deviation 0.0001 between aforementioned pairwise comparison.

3.2 Dry Matter Accumulation

During the (2013) growing season applying synthetic fertilizer from treatment application (T₂-T₅) did not increase dry matter yield as compared to treatment T₁ (Table 4). In the following production year (2014) all plots that received synthetic fertilizer (T₂-T₅) produced significantly greater yield response compared to treatment application T₁. However, during the next growing season (2015) we found that only treatment applications T₅ and T₃ produced a significant increase in dry matter yield (Table 4). When determining differences between increased fertilizer applications between synthetic fertilizer additions we found that treatments T₂ and T₃ produced comparable yields during the first production year (2013). However, during (2014) and (2015) a significant deviation in yield was observed in which T₃ produced higher dry matter yield compared to treatment application T₂ (Table 4). Treatment application T₃ and T₄ produced similar dry matter yields during the first and second production year (2013-2014). However, in (2015) yields retrieved from plots applied with treatment T₃ were significantly greater compared to yields recovered from treatment application T₄ (Table 4). No significant difference was determined between the application of treatments T₄ and T₅ during the first two production years, however, during (2015) T₅ produced a significantly greater yield as opposed to treatment T₄ (Table 4).

Table 4.) Differences of Tukey Post Hoc Analysis for Dry Matter Accumulation of the interaction effect between “Nutrient Management” and “Harvest Year” for *Miscanthus* collected during the 2013, 2014, and 2015 growing season. Individual Nutrient Management yields obtained at each fertilizer rate during each individual harvest year.

Pairwise Comparisons	2013 (<i>p</i> Value)	2014 (<i>p</i> Value)	2015 (<i>p</i> Value)
T ₁ vs.T ₂	0.70	0.01*	0.07
T ₁ vs.T ₃	0.08	<.0001**	<.0001**
T ₁ vs.T ₄	0.15	<.0001**	0.15
T ₁ vs.T ₅	0.11	0.001**	0.0004**
T ₂ vs.T ₃	0.18	0.0002**	0.002*
T ₂ vs.T ₄	0.29	0.01*	0.71
T ₂ vs.T ₅	0.22	0.26	0.03*
T ₃ vs.T ₄	0.77	0.11	0.001**
T ₃ vs.T ₅	0.92	0.004*	0.30
T ₄ vs.T ₅	0.84	0.15	0.01
Nutrient Management	2013 (t ha ⁻¹)	2014 (t ha ⁻¹)	2015 (t ha ⁻¹)
T ₁	9.16±0.80	11.02±1.60	26.76±0.64

T ₂	9.57±1.22	15.66±1.31	30.16±3.52
T ₃	12.37±0.86	23.54±0.80	36.15±3.05
T ₄	11.84±1.64	20.5±1.07	29.95±2.07
T ₅	11.71±0.22	18.24±1.51	34.26±2.88

†=* indicates a significant deviation between Main Effects “Nutrient Management” and “Harvest Year” at 0.001, and ** indicates highly significant deviation 0.0001 between aforementioned pairwise comparison.

4. Discussion

4.1 Comparison with Previous Research

Through the literature the growth response exhibited by *Miscanthus* to nutrient management is often debated. Due to the inconsistency of the reported findings we chose to compare our results with experiments that were conducted in a similar climatic zone. An experiment located in Lexington, KY focused primarily on Nitrogen addition to *Miscanthus* with a harvest window of 185 days growth, however, our experiment during first year growth used a 179 days. Maughn et al. [21] used N application rates 0, 60, and 120 kg ha⁻¹ for a duration of 185 days produced 15.2, 17.9, and 17.6 t ha⁻¹ of (DY) for the respective treatments while we found at 179 day duration dry yields of 9.16, 9.57, and 12.37 t ha⁻¹ for treatment applications T₁, T₂, and T₃. Researchers from Pulawy, Poland found that during 1st and 2nd year growth at application of 75-50-75 NPK kg ha⁻¹ was found to yield 12.05 t ha⁻¹ and 19.95 t/ha of (DY) when data was pooled between two growing sites [22]. The most relatable treatment to application 75-50-75 would be the 67-67-67 or T₂ treatment applied during our study. We observed first year yields of (9.57 t ha⁻¹) and (15.66 t ha⁻¹) when evaluating biomass produced during first and second establishment years. Researchers in North Carolina conducted a multi-year experiment growing *Miscanthus*, they found a more dramatic increase in yields that received nil-fertilizer application during 1st production year (2.61 t ha⁻¹) and during the 2nd production year (15.78 t ha⁻¹) [23]. Our results found that first and second year yields treated with T₁ were estimated at (9.16 t ha⁻¹ and 11.02 t ha⁻¹) in the following production year. The same researchers also compared yields during 1st and 2nd production years using fertilizer rate of 135-147-0 N-P-K kg ha⁻¹ and the corresponding (DY) was estimated at (3.02 t ha⁻¹) and (15.26 t ha⁻¹) [24]. Our treatment T₃ produced 12.37 t ha⁻¹ during the first production year (2013) and 23.54 t ha⁻¹ in the following production year (2014), this increase in biomass production could be due to better establishment of rhizomes from 2013 to 2014 and thereby put out more biomass.

5. Conclusion

Over the duration of this three year experimental study, we explored growth response due to nutrient management practices involving differential levels of synthetic fertilizer application. We found that dry matter accumulation increased from 2013 to 2014 by (38%), and from 2014 to 2015 by (43%). During 2014, and 2015 we found that treatment application T₃ produced significantly higher dry matter yields compared to the control treatment which increased yields by (53%), and (25%). We also found that dry matter yield increased by (33%) and (16%) during 2014 and 2015 growing seasons when treatment T₃ was applied as opposed to treatment T₂.

Depending on the project study outcomes, significant yield for *Miscanthus* can be achieved using 135-135-135 (kg ha⁻¹) of NPK versus no nutrient supplementation. However, application of increased doses of fertilizers did not produce significantly different yield over T₃.

References

- [1] Swaminathan K, Alabady M, Varala K, De Paoli E, Ho I, Rokhsar D, & et al. Genomic and small RNA sequencing of *Miscanthus* × *giganteus* shows the utility of sorghum as a reference genome sequence for Andropogoneae grasses. *Genome Biol* 2010;11(2).
- [2] Danalatos N, Archontoulis V, & Mitsios I. Potential growth and biomass productivity of *Miscanthus giganteus* as affected by plant density and N-fertilization in central Greece. *Biomass and Bioenergy* 2007;31(2): 145-152.
- [3] Clifton-Brown J, Chiang, Y. & Hodkinson R. *Miscanthus*: genetic resources and breeding potential to enhance bioenergy production. *Genetic improvement of bioenergy crops*. New York:Springer; 2008, p.273-294.
- [4] Robson P, Farrar K, Ga A, Jensen E, Clifton-Brown J, & Donnison I. Variation in canopy duration in the perennial biofuel crop *Miscanthus* reveals complex associations with yield. *Journal of experimental botany* 2013;ert104.
- [5] Melligan F, Dussan K, Auccaise R, Novotny E, Leahy J, Hayes M, & Kwapinski W. Characterization of the products from pyrolysis of residues after acid hydrolysis of *Miscanthus*. *Bioresource technology* 2012;108:258-263.
- [6] Christian D, Riche A, & Yates N. Growth, yield and mineral content of *Miscanthus* × *giganteus* grown as a biofuel for 14 successive harvests. *Industrial crops and products* 2008;28(3): 320-327.
- [7] Jørgensen U, & Schwarz K. Why do basic research? A lesson from commercial exploitation of *Miscanthus*. *New Phytologist* 2000;148(2):190-193.
- [8] Fowler P, McLauchlin A, & Hall L. *The potential industrial uses of forage grasses including Miscanthus*. Bangor, UK: BioComposites Centre, University of Wales; 2003
- [9] Shrestha P, Szaro T, Bruns T, & Taylor J. Systematic search for cultivatable fungi that best deconstruct cell walls of *Miscanthus* and sugarcane in the field. *Applied and environmental microbiology* 2011;AEM-02996.
- [10] Velásquez J, Ferrando F, & Salvado J. Effects of kraft lignin addition in the production of binderless fiberboard from steam exploded *Miscanthus sinensis*. *Industrial Crops and Products* 2003;18(1): 17-23.
- [11] El Hage R, Brosse N, Chrusciel L, Sanchez C, Sannigrahi P, & Ragauskas, A. Characterization of milled wood lignin and ethanol organosolv lignin from *miscanthus*. *Polymer Degradation and Stability* 2009;94(10):1632-1638.
- [12] Lewandowski I, Clifton-Brown J, Scurlock J, & Huisman W. *Miscanthus*: European experience with a novel energy crop. *Biomass and Bioenergy* 2000;19(4): 209-227.
- [13] Lewandowski I, & Schmidt U. Nitrogen, energy and land use efficiencies of *miscanthus*, reed canary grass and triticale as determined by the boundary line approach. *Agriculture, ecosystems & environment* 2006;112(4):335-346.
- [14] Ercoli L, Mariotti M, Masoni A, & Bonari E. Effect of irrigation and nitrogen fertilization on biomass yield and efficiency of energy use in crop production of *Miscanthus*. *Field Crops Research* 1999;63(1): 3-11.

- 265 [15] Christian D, Poulton P, Riche A, Yates N, & Todd A. The recovery over several seasons
 266 of ¹⁵N-labelled fertilizer applied to *Miscanthus giganteus* ranging from 1 to 3 years old.
 267 Biomass and Bioenergy 2006;30(2):125-133.
- 268 [16] Beale C, & Long S. Seasonal dynamics of nutrient accumulation and partitioning in the
 269 perennial C₄ grasses *Miscanthus* × *giganteus* and *Spartina cynosuroides*. Biomass and
 270 Bioenergy 1997;12(6):419-428.
- 271 [17] Schwarz H, Liebhard P, Ehrendorfer K, Ruckebauer P. The effect of fertilization on yield
 272 and quality of *Miscanthus sinensis* Giganteus. Industrial Crops and Products 1994;2(3):
 273 153-159.
- 274 [18] Schwarz H. *Miscanthus sinensis* ‘giganteus’ production on several sites in Austria.
 275 Biomass and Bioenergy 1993;5(6):413-419.
- 276 [19] Clifton-Brown C, Neilson B, Lewandowski I, & Jones B. The modelled productivity of
 277 *Miscanthus* × *giganteus* (GREEF et DEU) in Ireland. Industrial Crops and Products
 278 2000;12(2):97-109.
- 279 [20] Heaton, E, Dohleman, F, & Long P.. Meeting US biofuel goals with less land: the
 280 potential of *Miscanthus*. Global Change Biology 2008;14(9):2000-2014.
- 281 [21] Maughan M, Bollero G, Lee D, Darmody R, Bonos S, Cortese L, & Voigt T. (2012).
 282 *Miscanthus* × *giganteus* productivity: the effects of management in different
 283 environments. GCB Bioenergy, 4(3), 253-265.
- 284 [22] Borzecka-Walker M, Faber A, & Borek R. (2008). Evaluation of carbon sequestration in
 285 energetic crops (*Miscanthus* and coppice willow). Int. Agrophysics, 22, 185-190.
 286 Haines S. (2011). Nitrogen and Phosphorus Fertilizer Effects on Establishment of
 287 *Miscanthus* × *giganteus* in North Carolina.