

# The Relationship between Body Conformation, Testicular Traits and Serum Testosterone Levels in Pre-pubertal Male Boer Goat Crosses

Chukwuemeka Okere\* Latoya Keith and Olga Bolden-Tiller

Department of Agricultural and Environmental Sciences, Tuskegee University, Tuskegee, AL, 36088

## ABSTRACT

In the present study, the relationship among body conformation traits, scrotal circumference and serum testosterone concentrations were investigated in pre-pubertal male Boer goat crosses at the Caprine Research and Education Unit, Tuskegee University, Tuskegee, AL. Body conformation traits (chest girth -CG, height at withers-HTW, body length -BL, body condition scores -BCS, body weight -BW, shoulder width -SW), and scrotal circumference -SC were monitored at 3 week intervals for 12 weeks. Also, blood samples were collected and a calibrated IMMULITE 1000 assay system was used for the quantitative measurement of total serum testosterone levels (TT). Although, results show a non-significant relationship between TT levels and many body conformation traits, serum testosterone levels were lowly and positively correlated to BL ( $r = 0.19$ ), BW ( $r = 0.19$ ). Whereas, CG ( $r = 0.13$ ), HTW ( $r = 0.11$ ) and SC ( $r = 0.18$ ) were non-significant. SC was moderately correlated to BL ( $r = 0.30$ ;  $P = 0.001$ ) and SW ( $r = 0.33$ ;  $P < .001$ ) and strongly correlated with BW (0.61), CG ( $r = 0.53$ ), HTW ( $r = 0.41$ ) respectively. In addition, SC and BW increased ( $P < .01$ ) linearly from week 6 through 12. It is speculated that the week 9 peak of TT levels obtained in this study represents the attainment of sexual maturity (puberty) in Boer goat male crosses. Based on the results of this study, we hypothesized that SC measurements, when used in conjunction with TT levels and body conformation traits can be a valuable breeding soundness evaluation tool for selecting or culling breeding Boer goat sires at an early age by limited resource producers.

**Keywords:** Boer goats, body conformation, testicular traits, serum testosterone

## 1. INTRODUCTION

Meat goat breeds available for production in the U.S include: South African Boer, New Zealand Kiko, Myotonic, Savannah, and Spanish goats [1]. The Boer goat from South Africa is a breed developed for meat production that evolved from selection pressures placed on indigenous goats of the region by farmers [2]. Genetic progress in the meat goat industry requires the identification of superior meat goat sires. The most accurate way to test a sire's genetic worth is to perform breeding

\* Tel. : (334) 727-8904; fax: (334) -727-8552.  
E-mail address: cokere@mytu.tuskegee.edu.

24 soundness evaluation (BSE) and also generate progeny from the animal [3]. BSE  
25 predicts the potential fertility of a buck. It is based on an examination that includes  
26 tests for physical soundness, **scrotal circumference**, semen quality, and in some  
27 cases, serum testosterone profiles, and libido/mating ability.

28         The age and weight at which puberty occurs vary greatly among breeds of  
29 goats and the level of nutrition during development; however, research with various  
30 breeds [4, 5] suggests that a practical indication of imminent puberty is when scrotal  
31 circumference is between 25 and 27 cm. In bulls, there is a positive genetic  
32 correlation between a sires' scrotal circumference, the scrotal circumference of his  
33 sons, and the pregnancy rate of his daughters [6]. This indicates that bulls with a  
34 larger scrotal circumference are likely to sire sons with larger scrotal circumference,  
35 and daughters likely to reach puberty at younger ages.

36         Testosterone is a steroid hormone produced by cells of the testes, and is  
37 considered to be the primary circulating androgen that regulates testicular function.  
38 Testosterone levels are useful in selection of young sires, and in characterizing  
39 sexual maturity in different breeds of **sires** [7]. It is important to evaluate  
40 testosterone levels to determine the development of the reproductive system [8].

41         The relationship between body measurements and serum testosterone in  
42 10-12 month old Dorper rams was reported by **Fourie et al. (2005)**. **Serum**  
43 **testosterone did not have a significant correlation with most of the body parameters**  
44 **measured [9]**. However, serum testosterone concentration in rams was positively  
45 correlated to masculinity (**muscle score**), and scrotal circumference. Scrotal  
46 circumference was positively and significantly correlated with body weight, body  
47 length, chest dept, shoulder height, shoulder width, hindquarter width, canon bone  
48 length, masculinity wedge shape, selection index and age. **Rams with a wedge**  
49 **shape scores between five and seven were significantly heavier than those with**  
50 **type scores between two and four**. Collectively, **Fourie et al. (2005)** concluded that  
51 there is a significant and positive relationship between scrotal circumference and  
52 serum testosterone, but no significant correlation between linear shape type score  
53 and serum testosterone levels.

54         Testicular development in relation to age, body weight, semen  
55 characteristics and testosterone was reported in Kivircik ram lambs [10]. All

measurements of testis, body weight, and serum testosterone concentrations were positively correlated with each other. A significant positive correlation was found at seven to eight months of age between all testicular measurements, semen volume and motility [10]. Scrotal circumference could provide useful estimate of testicular growth, as its correlations with other testicular measurements were the highest, and this information could be used as selection criteria for ram lambs at an early age.

A procedure that would link external body conformation and testicular traits with serum testosterone or semen quality may provide a good guide to breeding soundness evaluation in meat goat sires [11]. Since limited resource producers may not be in a position to test libido and/or ejaculate qualities of males before using them for breeding, a procedure that would link external testicular measurements with serum testosterone may provide a good guide to breeding soundness evaluation, especially, where bucks are reputed to have exceptionally high libido [12, 13].

There is no evidence in literature to suggest that this approach has been used in selecting potential Boer meat goat sires for superior breeding. Therefore, the aim of this study is to identify the relationship between certain body conformation, testicular traits and serum testosterone levels of pre-pubertal male Boer crosses.

## 2. MATERIAL AND METHODS

### 2.1. Animal Management:

Twenty five pre-pubertal male Boer goat (Boer x Kiko) crosses ( $110.11 \pm 20.1$  days old), and singled sourced from a local meat goat producer were used in this study. The Tuskegee University Animal Care and Use Committee approved animal care, handling and sampling procedures. Upon arrival the animals were given an overall health check by an attending veterinarian at Tuskegee University's School of Veterinary Medicine. The animals were treated with Panacur (*fenbendazole*) to control the development and reproduction of internal parasites and quarantined for three weeks at the Tuskegee University's Caprine Research Facility. Each animal was housed for the entire experimental period in individual 1.8 x 2.1m indoor pens after the quarantine period. Throughout the experimental period, animals were maintained on a daily diet that consisted of a high energy concentrate that was

given at 2lbs/day. The animals were also allowed *ad libitum* access to hay, water and mineralized salt blocks.

## **2.2. Body Conformation Measurements**

The body conformation measurements recorded at 3-week intervals for 12 weeks (wk 0/31 d, wk 3/152 d, wk 6/173 d, week 9/194 d, and wk 12/215 d of age). Body weight, (BW, kg) was recorded using a MTIAHS500 Sheep and Hog Scale System). Whereas body condition score (BCS) was scored on a subjective scale of 1 = emaciated to 5 = obese). A measuring tape was used to determine shoulder width (the horizontal distance between the processes on the left shoulder to those on the right shoulder blade), chest girth (the width around the chest just behind the front legs), body length (the distance from the sternum to the aitch bone), hip width (the distance between the left and right femur bones). The height at wither (the vertical length from the thoracic vertebrae to the ground) was determined with the aid of a metric ruler.

## **2.3. Testicular Measurements**

The scrotal circumference was determined in each animal by pulling the testicles firmly into the lower part of the scrotum, grasping the neck of the scrotum with one hand, squeezing and pulling down. Thereafter, the circumference was measured with the aid of a measuring tape and recorded every 3 weeks for 12 weeks as the largest diameter of the scrotum.

## **2.4. Blood Collection**

Blood samples were collected at intervals of three weeks for 12 weeks from each animal. The blood was collected via jugular venipuncture into 10ml heparinized vacutainer tubes, and placed on ice immediately after collection. The serum was separated by centrifugation at 3000 rpm at 4°C for 10 minutes and aliquoted into separate vials which were kept frozen at -4° C until testosterone assay.

## **2.5. Serum Testosterone Assay**

A calibrated IMMULITE 1000 system (developed at Meharry Medical College, Nashville, Tennessee) was used for the quantitative measurement of total serum testosterone. IMMULITE 1000 Total Testosterone is a solid-phase, enzyme-labeled, competitive chemiluminescent immunoassay. The solid-phase, a

122 polystyrene bead enclosed within an IMMULITE Test Unit, is coated with a  
123 polyclonal rabbit antibody specific for testosterone. Serum samples and alkaline  
124 phosphatase-labeled testosterone were simultaneously introduced into the Test  
125 Unit, and incubated for approximately 60 minutes at 37°C with intermittent agitation.  
126 During this time, testosterone in the sample competed with alkaline phosphatase  
127 labeled testosterone for antibody-binding sites on the bead. Unbound material was  
128 then removed by a centrifugal wash. Substrate is then added, and the Test Unit is  
129 incubated for another 10 minutes.

130 The chemiluminescent substrate, a phosphate ester of adamantyl dioxetane,  
131 undergoes hydrolysis in the presence of alkaline phosphatase to yield an unstable  
132 intermediate. The continuous production of these intermediate results in the  
133 sustained emission of light, thus improving precision by providing a window for  
134 multiple readings. The bound complex - and thus also the photon output, as  
135 measured by the luminometer - is inversely proportional to the concentration of  
136 testosterone in the sample.

## 137 **2.6 Statistical Analysis**

138 Descriptive statistics [14] was performed on the data to determined individual  
139 buck differences (means and standard deviations) in selected body conformation,  
140 testicular traits, and serum testosterone profiles. Also, data was subjected to  
141 analysis of variance using the GLM procedures [14]; correlation coefficients (r) were  
142 established between various body, testicular parameters and serum testosterone  
143 profiles.

## 144 145 **3. RESULTS AND DISCUSSION**

146  
147 The means and standard deviations for body conformation, testicular traits  
148 and serum testosterone levels (TT) are shown in Tables 1.

149

150 **Table 1. Body Conformation, Testicular Traits, Serum Testosterone in Pre-pubertal**  
 151 **Male Boer Goat Crosses.**

Parameters	Week 0 (N= 25)	Week 3 (N=23)	Week 6 (N= 23)	Week 9 (N=23)	Week 12 (N= 23)
Body Condition Score (BCS, 1-5)	3.40 ± 0.50 <sup>b</sup>	3.21± 0.51 <sup>ab</sup>	3.08± 0.41 <sup>ab</sup>	2.91± 0.59 <sup>a</sup>	3.13± 0.69 <sup>ab</sup>
Body Length (BL, cm)	58.62 ± 2.41 <sup>a</sup>	61.95± 3.21 <sup>b</sup>	63.72± 2.26 <sup>bc</sup>	65.25± 2.34 <sup>cd</sup>	67.13± 2.75 <sup>d</sup>
Body Weight (BW, kg)	27.66 ± 2.54 <sup>a</sup>	27.52± 2.20 <sup>a</sup>	31.95± 2.64 <sup>b</sup>	34.72± 2.98 <sup>c</sup>	37.46± 3.44 <sup>d</sup>
Chest Girth (CG, cm)	66.34± 3.22 <sup>ab</sup>	64.37± 1.95 <sup>a</sup>	67.82± 2.58 <sup>bc</sup>	68.91± 2.21 <sup>cd</sup>	70.79± 2.35 <sup>d</sup>
Height at Withers (HTW, cm)	58.01± 1.90 <sup>a</sup>	59.97± 2.27 <sup>b</sup>	61.20± 1.86 <sup>bc</sup>	62.29± 2.13 <sup>c</sup>	64.58± 2.39 <sup>d</sup>
Hip Width (HW, cm)	42.16± 3.12 <sup>a</sup>	41.95± 2.28 <sup>a</sup>	43.63± 1.65 <sup>ab</sup>	42.18± 1.68 <sup>a</sup>	45.06± 2.06 <sup>b</sup>
Scrotal Circumference (SC, cm)	22.87± 2.42 <sup>a</sup>	22.54± 1.77 <sup>a</sup>	23.22± 0.86 <sup>ab</sup>	24.53± 1.43 <sup>b</sup>	26.05± 1.35 <sup>c</sup>
Shoulder Width (SW, cm)	41.14± 2.54 <sup>a</sup>	40.40± 1.84 <sup>a</sup>	41.73± 1.52 <sup>a</sup>	41.29± 1.92 <sup>a</sup>	44.06± 2.11 <sup>b</sup>
Testosterone (TT, ng/ml)	5.625± 6.03 <sup>a</sup> (N= 20)	8.74±11.26 <sup>a</sup> (N= 17 )	7.48±8.25 <sup>a</sup> (N= 23)	27.48±14.42 <sup>b</sup> (N= 23)	10.74±14.40 <sup>a</sup> (N=23)

152 \*Means ± Standard Deviations

153 <sup>a,b,c,d</sup> Means in each row with the same superscript are not significantly different ( $P =$   
 154 .05).

155 The average SC at week 0 was (22.87±2.42cm), with a slight decline into week 3  
 156 (22.54±1.77cm). However, SC increased linearly from week 6 through week 12  
 157 (23.22±0.86cm, 24.53±1.43cm and 26.05±1.35cm), respectively. There was no  
 158 significant difference for SC between weeks 0, 3, and 6; and week 6 and week 9  
 159 ( $P = .15$ ), respectively. In addition, SC and BW increased correspondingly from  
 160 week 6 through week 12 (23.22±0.86cm; 31.95 ±2.64kg) (24.53±1.43cm; 34.72  
 161 ±2.98kg) and (26.05±1.35cm; 37.46 ±3.44kg), respectively. The average TT level at  
 162 week 0 or 131 d of age was 5.625±6.03ng/ml, with a slight increase into week 3 or  
 163 152 d of age (8.74±11.26ng/ml,  $P = .09$ ). Also, there was a decline ( $P = .11$ ) in TT  
 164 level at week 6/173 d of age (7.48±8.25ng/ml), and week 9/194 d of age showed an  
 165 all time high (27.85±14.42ng/ml), followed by a decline at week 12/215 d of age  
 166 (10.74±14.40ng/ml). For weeks 0, 3, 6 and 12 there was no significant difference

167 ( $P=.19$ ) for TT levels. However, TT for week 9 was significantly different from week  
168 0, 3, 6 and 12 ( $P=.05$ ), respectively. It is speculated that the week 9 peak of serum  
169 testosterone levels obtained in this study represents the attainment of sexual  
170 maturity (puberty) in Boer male crosses. Bezerra et al. (2009) reported mean  
171 testosterone levels varying from ( $0.259 \pm 0.172$  ng/ml to  $4.613 \pm 2.892$  ng/ml and 0  
172  $.521 \pm 0.311$  to  $3.417 \pm 2.021$  ng/ml) in Boer goats starting from 1 month to 8 months  
173 of age during dry and rainy seasons. In young Saneen and British Alpine goats,  
174 testosterone levels are marked by an initial decline followed by a peak at the time  
175 when male reach sexual maturity [16, 17].

176 The correlation coefficients (r) for body conformation, testicular traits and  
177 serum testosterone levels are presented in Table 2. TT levels recorded a low to  
178 moderate correlation with most of the body conformation traits measured in this  
179 study. Whereas coefficients for: SW vs. TT and TT vs. CG were non-significant.  
180 Fourie et al. (2005) found similar results in Dorper rams, where serum testosterone  
181 concentration was positively correlated to masculinity ( $r = 0.15$ ;  $P=.05$ ) and scrotal  
182 circumference ( $r = 0.23$ ;  $P<.05$ ), which is slightly higher than values reported in the  
183 current study. Species difference (ram vs. bucks) could explain the discrepancy in  
184 the TT vs. SC correlation values.

185 According to Coulter and Foote (1976), SC is an important trait that is closely  
186 associated with the testicular growth and sperm production in males of all meat  
187 animals. Thus, selecting males based on their SC would result in larger testes,  
188 potentially with the capacity to produce more semen [18]. Being a highly heritable  
189 component of fertility, it is important to include SC during animal evaluation for  
190 breeding soundness [19]. In the current study, SC was positively and significantly  
191 ( $P<.001$ ) correlated to BL ( $r = 0.30$ ), BW ( $r = 0.61$ ), CG ( $r = 0.53$ ), HTW ( $r = 0.41$ ) and  
192 SW ( $r = 0.33$ ). Similar results were reported by Fourie et al. (2005) who found  
193 positive and significant correlations between SC and BW ( $r = 0.38$ ), BL ( $r = 0.34$ )  
194 and SW ( $r = 0.27$ ) in Dorper rams. Adeyinka and Mohammed (2006) reported  
195 positive correlations between TT and BW ( $r = 0.30$ ;  $r = 0.43$ ), SC ( $r = 0.42$ ;  $r = 0.52$ ),  
196 and BW and SC ( $r = 0.93$ ;  $r = 0.88$ ) in young Boer bucks. Ugwu (2009) reported a  
197 highly significant positive relationship between SC and testis weight of West African  
198 Dwarf bucks. Testis weight is known to be highly correlated ( $r = 0.93$ ) with testicular



sperm reserves [13] and males with larger testes tend to produce more sperm [21]. It follows that a good measurement of scrotal circumference would be a reliable predictor of sperm producing capacity.

Overall, results from this study indicate that there are significant and positive correlations between few body conformation traits and serum testosterone levels. However, a low to moderate correlation was found between scrotal circumference and serum testosterone level in pubertal male Boer crosses. In general, correlations between serum testosterone levels and body conformation in this study were either low or negative. However, scrotal circumference was found to have a higher correlation with body weight than serum testosterone level in pubertal male Boer crosses. Based on the results of this study, it is hypothesized that monitoring scrotal circumference, serum testosterone levels and body conformation traits is a useful tool for selecting superior breeding Boer goat sires at an early age.

Table 2. Correlation Coefficients (r) for Body Conformation, Testicular Traits and Serum Testosterone in Pubertal Male Boer Crosses

	BCS	BL	BW	CG	HTW	HW	SC	SW
BL	-0.0819							
BW	0.1677	0.6715***						
CG	0.1739	0.4941***	0.8280***					
HTW	-0.1447	0.5914***	0.6631***	0.4979***				
HW	0.1290	0.3635***	0.4589***	0.3612***	0.3675***			
SC	0.0946	0.3026***	0.6117***	0.5262***	0.4111***	0.1721		
SW	0.0891	0.2771**	0.4994***	0.4671***	0.3897***	0.4539***	0.3291***	
TT	-0.1952*	0.1853*	0.1866*	0.1279	0.1124	-0.1299	0.1776	-0.0851

BL= Body Length, BW= Body Weight, CG= Chest Girth, HTW= Height at Withers, HW= Hip Width, SC= Scrotal Circumference, SW= Shoulder Width, TT = Testosterone Level

\*= Significant if  $P \leq .05$

\*\*= Significant if  $P \leq .01$

\*\*\*= Significant if  $P < .001$



The present finding of positive correlation for SC with testosterone in the pubertal bucks may indicate the importance of monitoring SC and/or serum testosterone levels in testicular development and the onset of puberty in male Boer goat crosses. Studies in other animal species [9, 13, 14, 15] have increasingly shown the role of testosterone in male reproductive functions. Further studies on the activity of testosterone receptors in the caprine testis around and after puberty are needed to better understand the functional role of testosterone.

#### 4. CONCLUSION

1. There was an intricate relationship among testosterone concentrations, scrotal circumference, and various body conformation traits.
2. There were significant and positive correlations between body length; body weight and serum testosterone levels. However, low to moderate correlation was found between scrotal circumference and serum testosterone level in pubertal male Boer crosses.
3. Scrotal circumference was found to have a higher correlation with body weight than serum testosterone level in pubertal male Boer crosses.
4. The present results could assist the development and implementation of selection or culling criteria for breeding Boer goat sires at an early age.

#### ACKNOWLEDGEMENTS

Authors would like to thank Ms. Davi-Anne Phillip for editorial assistance. Also, Mr. Mel Jones and Mr. Danny Williams of the Caprine Unit, and George Washington Caver Agricultural Experimental Station, Tuskegee University, Tuskegee, AL for providing financial and technical support.

#### REFERENCES

1. Browning R, Payton, Donnelly B. Leite-Browning ML, Pandya P, Hendrixson AC, Byars M. Evaluation of three meat goat breeds for doe fitness and reproductive performance in the southeastern United States. In:

- 256            Proceedings of 8th world Congress on Genetics Applied to Livestock  
257            Production. Belo Horizonte, Brazil: 2006; August 13 – 18.
- 258            2. Casey NH vanNiekerk WA. The Boer goat. I. Origin, Adaptability,  
259            Performance Testing, Reproduction, and Milk Production. Small Ruminant  
260            Research, 1988; 1(3): 291-302.
- 261            3. Ott RS. 1986. Breeding soundness examination of bulls. In: Morrow DM (ed).  
262            Current Therapy in Theriogenology. Pp 125.136. WB Saunders,  
263            Philadelphia, 1986.
- 264            4. Ford D, Okere C, Bolden – Tiller O. Libido Test Scores, Body Conformation  
265            and Testicular Traits in Boer & Kiko Goat Bucks. Journal of Agricultural and  
266            Biological Sciences; 2009, 4 (5): 1-8.
- 267            5. Keith L, Okere C, Solaiman S, Bolden-Tiller O. Accuracy of Predicting Body  
268            Weights from Body Conformation and Testicular Morphometry in Pubertal  
269            Boer Goats. Research Journal of Animal Science. 2009; 3 (2): 26-31.
- 270            6. Elmore RG, Bierschwal CJ, Martin CD, Youngquist RE. A summary of 1127  
271            breeding soundness examinations in beef bulls. Theriogenology 1975;  
272            3:209-218.
- 273            7. Eloy AMX Santa Rosa JS. Perfis Plastimaticos de Testosterona Durante a  
274            Puberdade de Machos Caprinos da Raca Moxoto. Pesquisa Agropecuaria  
275            Brasileira, 1998; 33: 1645-1652.
- 276            8. Chakraborty PK, Stuart LD Brown JL. Puberty in the Male Nubian Goat:  
277            Serum Concentrations of LH, FSH and Testosterone from Birth through  
278            Puberty and Semen Characteristics at Sexual Maturity. Animal  
279            Reproduction Science. 1989; 20: 91-101.
- 280            9. Fourie PJ Schwalbach LM Neser FWC Greyling JPC. Relationship between  
281            Body Measurements and Serum Testosterone Levels of Dorper Rams.  
282            Small Ruminant Research 2005; 56:75-80.
- 283            10. Elmaz O, Cirit U. Demir H. Relationship of Testicular Development with  
284            Age, Body Weight, Semen Characteristics and Testosterone in Kivircik  
285            Ram Lambs. South African Journal of Animal Science 2007; 37(4): 269-  
286            274.

- 287 11. Okere C, Bradley P, Bridges ER, Bolden-Tiller O, Ford D, Paden A.  
288 Relationships among Body Conformation, Testicular Traits and Semen  
289 Output in Electro-ejaculated Pubertal Kiko Goat Bucks. Journal of  
290 Agricultural and Biological Sciences. 201; 16 (8): 43 - 48.
- 291 12. Adeyinka IA, Mohammed ID. Relationship between Live Weight and Linear  
292 Body Measurements in Two Breeds of Goats of Northern Nigeria. Journal of  
293 Animal and Veterinary Advances, 2006; 5 (11): 891-893.
- 294 13. Ogwuegbu SO, Oko BO, Akusa MO and Arie TA. Gonadal and  
295 Extragonadal Sperm Reserves of Maradi (Red Sokoto) Goat. Journal of Bull  
296 and Animal Health Production in Africa, 1985; 33:139-141.
- 297 14. Statistix 7, 2000. Version 7 for Windows. Analytical Software, Analytical  
298 Software, P.O. Box 12185, Tallahassee , FL 32317 USA.  
299 [www.statistix.com](http://www.statistix.com)
- 300 15. Bezerra, FQG, Aguiar Filho CR, Freitas Neto LM, Santos Junior ER,  
301 Chaves RM, Azevedo, EMP. Santos MHB, Lima PF, Oliveira MAL. Body  
302 Weight, Scrotal Circumference and Testosterone Concentration in Young  
303 Boer Goat Males Born During the Dry of Rainy Seasons. South African  
304 Journal of Animal Science, 2009; 39 (4): 301-305.
- 305 16. Macmillan KL. Hafs HD. The Reproductive Tract of Holstein Bulls from  
306 Birth to Puberty. Journal of Animal Science, 1969; 28:233-239.
- 307 17. Ahmad, N, Noakes, DE. Wilson CA. Secretory Profiles of LH and  
308 Testosterone in Pubescent Male Goat Kids. Small Ruminant Research,  
309 1996; 21:51-56.
- 310 18. Coulter GH, Foote RH. Effect of Season and Year of Measurement on  
311 Testicular Growth and Consistency of Holstein Bulls. Journal of Animal  
312 Science, 1976; 42:434-438.
- 313 19. Bailey TL, Monke D, Hudson RS, Wolfe DF, Carson RL. Riddle MG.  
314 Testicular Shape and its Relationship to Sperm Production in Mature  
315 Holstein Bulls. Theriogenology, 1996; 54:881-887.
- 316 20. Ugwu, SOC. Relationship between Scrotal Circumference, *In Situ* Testicular  
317 Measurements and Sperm Reserves in West African Dwarf Bucks. African  
318 Journal of Biotechnology, 2009; 8(7):1354-1357.

319 21. Okwun, OE, Igboeli, G, Ford, JJ, Lunstra, DD. Johnson, LJ. Number and  
320 Function of Sertoli Cell, Number and Yield of Spermatozoa and Daily  
321 Sperm Production in Three Breeds of Boars. Journal of Reproduction  
322 Fertility, 1996; 107: 137-149.

323

324

325