



Original Research Article

HEIGHT-DIAMETER RELATIONSHIP MODELS FOR TEAK (*Tectona grandis*) PLANTATION IN NIMBIA FOREST RESERVE, NIGERIA

ABSTRACT

The study on height-diameter modelling was carried out on Teak (*Tectona grandis*) plantation in Nimbia Forest Reserve. Two fit methods (Chapman-Richards and Weibull) were used to model height-diameter relationship. The teak plantation is divided into four forest beats as management unit, therefore, stratified random sampling was employed to select five plots of 20 x 20 m from each forest beat (as stratum), thereby having a total of 20 plots. Stump diameter (Dst), diameter at breast height (Dbh), diameter at middle and top positions (Dm and Dt) of trees, and tree height for all the selected trees were measured. Average Dbh measured 15.00cm and the mean tree height was 6.77m. Pseudo coefficient of determination (Pseudo R^2) and residual mean square error (RMSE), goodness-of-fit statistics were considered as model selection criteria. Chapman-Richards function produced the best goodness-of-fit statistics for Teak height-diameter modelling. The height-diameter models require additional site factors for better models; hence the need for establishing permanent sample plots (PSP) in order to get additional information from re-measurements of the plots.

INTRODUCTION

Growth and yield models are generally used to predict temporal and future growth of forests. In forestry, it is important to be able to make accurate future predictions of the mean values of growth variables based on repeated measurements through time made on units grouped hierarchically. In many forest management practices, decisions are based on yield projections that crucially depend on projections of plot level averages of tree height, basal area, and other morphometric variables [1].

Knowledge of diameter at breast height (Dbh) and tree height is fundamental to both developing and applying many growth and yield models. One of the greatest challenges in model development is measuring tree variables and data gathering. In forest measurement, Dbh of a tree can be measured quickly, easily, and accurately, but the measurement of total tree height is relatively complex, time consuming and expensive [2]. Furthermore, some site conditions and tree composition especially in tropical forests may prevent accurate height measurements on all trees measured for Dbh as it may not be possible to unambiguously observe a given tree, or reach an appropriate vantage point. Therefore, with many permanent and temporary sample plot systems, Dbh is conventionally measured for all trees sampled, but height is measured for only a sub-sample of trees selected across the range of diameters observed [3]. Height-diameter relationship models are then used to estimate the heights of trees measured only for diameter. A number of height-diameter models have been developed using only Dbh as a predictor variable for estimating total or merchantable tree height (e.g. [4]; [2]; [5]; [6]; [7]). However, the relationship between the diameter and its height varies among stands [8] and depends on the growing environment and stand conditions [9].

Teak, though native of India, Myanmar (Burma), Thailand and Laos, grows nowadays in the whole inter-tropical region (excluding deserts areas of Africa) [10]. In the international market, Teak has been recognized for its high quality timber. Its high durability, good dimensional stability and aesthetic qualities make it a very profitable option for both public and private forestry schemes [11]. As a sustainable supply of timber from natural forests shrinks and the demand continues to increase, the general trend in the future of teak growing will be towards increasing production and utilizing more plantation grown teak [12]. This suggests a need for enhanced knowledge ranging from diverse aspects of teak plantations establishment as well as silviculture, management, utilization and the ecological aspects of both plantations and natural stands [13].

52 MATERIALS AND METHODS

53 Study Area

54 Nimbia forest reserve is located in the Northern Guinea Savanna zone of Nigeria but
55 in the derived savanna zone at the eastern part of Jema'a Local Government Area of Kaduna
56 state, 70km south east of Jos, along Jos-Kafanchan road. It lies between latitudes $9^{\circ}29'$ and
57 $9^{\circ}32'N$ and longitudes $8^{\circ}30'$ and $8^{\circ}36'E$ with an elevation of about 594m above sea level
58 [14].

59 Nimbia forest reserve has an undulating topography. The eastern end of Nimbia forest
60 reserve is the last part of Assop escarpment. It descends in a series of steps with long level
61 stretches interrupted by steep boulder-strewn descents. It descends more gently west wards to
62 Jama'a-Jagindi plains. The forest reserve drains south-west into the Gimi River and west
63 towards the stream that forms the western boundary. There is no permanent stream that flows
64 through the reserve. The northern and southern parts of the reserve are bounded by Lioc
65 stream and Gimi River respectively.

66 Nimbia forest reserve is within the Jema'a platform and is underlain predominantly by
67 igneous and metamorphic rocks. The greater part of the reserve is underlain by the Newer
68 Basalt of the late Tertiary and Quaternary periods which is composed of olivine, amygdalites
69 and zeolites.

70 There are different soil types within the plantation, but most commonly found are the
71 red deep freely draining loams and dark-brown loamy soils. According to [15] soils of
72 Nimbia area as indicated on the soil map of Nigeria  as belonging to the unit 15F and are
73 classified as Typical Dystrusteps (USDA)/ Dystric Cambisols (FAO ). The soils also belong
74 to the Nimbia series which developed from weathered olivine basalt and classified as
75 Eutrophic Brown soil by [16].

76 The climate is determined by altitude and its location in relation to the seasonal
 77 migration of the inter-tropical convergence zone. The position of Nimbia with respect to
 78 altitude (594m above sea level) induces orographic rain and has an annual rainfall of between
 79 1500mm – 2000mm spread over a period of seven months (April - October) while the dry
 80 months are five months (November - March). Minimum temperatures range between 17°C
 81 and 22°C (December - March) and the maximum ranges from 28°C to 35°C (August -
 82 March). Relative humidity is between 30 and 36% in the dry season and 95% in the rainy
 83 season.

84 The natural vegetation of Nimbia ranges from the Southern Guinea Savanna to a dry
 85 type of rain forest, which consists mostly of savannah woodland with tall grasses and
 86 broadleaved trees usually with short boles (e.g. *Milicia excelsa*, *Guiera senegalensis*, *Parkia*
 87 *biglobosa*, *Daniela olivera* and *Vitex doniana*). Over the years the reserve has been over
 88 exploited which resulted in the introduction of fast growing exotic species specifically
 89 *Tectona grandis* and *Gmelina arborea*. The trees were planted in rows of 3 x 3m, both inter
 90 and intra row spacing attaining a height of about 13.6m and diameter of about 41cm
 91 averagely at the age of 20years [14].

92 **Sampling and Data Collection**

93 A reconnaissance survey of the Teak plantation was carried out with an attempt to
 94 locate the boundaries of the plantations and get acquainted with the general conditions of the
 95 forest reserve. With the assistance of the forest officers and forest guards in the reserve, the
 96 selected plots and stands were located by the use of prismatic compass, GPS, 50-meter tape,
 97 cutlass, and ranging poles. Nimbia forest reserve is divided into four different forest bits and
 98 each bit is divided into compartments as management units. Stratified Random Sampling
 99 Technique was employed, considering each forest beat as a stratum. Each stratum was further

100 divided into 20 x 20 m sampling units out of which five plots were randomly selected,
101 thereby having a total of 20 plots for the study.

102 Within each randomly selected sample plot, preference was given to the enumeration
103 of healthy trees with more typical growth forms; dead trees and trees with abnormal form
104 were avoided. This is because in developing growth and yield models for effective
105 management of forest plantations, only such trees that meet grade, soundness and size
106 requirements for commercial logs or poles are considered relevant. While this sampling
107 guideline appeared to introduce a bias in favor of better-formed trees, it is justified because
108 only healthy trees with good form are of commercial value and therefore, require volume
109 computation. The following measurements were carried out on each selected tree:

- 110 1. Stump diameter (35 cm above ground level) and diameter at breast height (at 1.3 m
111 above ground) using the girth-diameter tape (in cm);
- 112 2. Stem diameter at the middle and upper positions of the tree using the Spiegel relaskop
113 (in cm)
- 114 3. Merchantable height of selected trees using the Spiegel relaskop up to 7 cm upper
115 diameter limit (in meters);
- 116 4. Total height of dominants and co-dominant trees using the Spiegel relaskop (in
117 meters).

118 **Data Analysis**

119 Data collected were organized and screened (removing the outliers) for analysis.
120 Descriptive statistical analysis was further carried out in order to summarize the data. All
121 analysis carried out were conducted using SAS statistical package version 9.3, licensed to the
122 University of British Columbia, Vancouver, Canada.

123

124

Fitting of Height-diameter Models

For the construction of height-diameter models, the first step was to plot the data from field measurements of height against the Dbh in a scatter form. This gives an outlook of how the relationship between the two variables would be. Different height-diameter models (Exponential, Chapman-Richards, Gompertz, and Modified logistic) were reviewed before choosing the two functions used for this study and out of them the Chapman-Richards was found to be the most widely used, but the decision to add another model was to enable selection of the better one. Considering the model flexibility; region and species of this study, the Chapman-Richards and Weibull functions were selected for the height-diameter modelling due to their simplicity, accuracy, and precision. PROC NLIN statistical analysis method in SAS package was used to fit the two functions and the model form was presented in Table 1.

Table 1: Models selected for height-diameter

Model	Expression	References
Chapman-Richards	$H = 1.3 + \alpha (1 - \exp^{-\beta Dbh})^\gamma$	[17]; [18]
Weibull	$H = 1.3 + asym. (1 - \exp[(-b * Dbh)^c])$	[19]

RESULTS AND DISCUSSION

Summary Statistics of Measured Parameters

For efficient and accurate growth and yield modelling, field data must be biologically and statistically valid before using them to develop models. In this study the data used were carefully obtained from the field and subjected to biological validation and the results indicated a normal distribution pattern as tree tapers from bottom to the top, so does the diameter decreases from bottom to the top as indicated in Table 2. This shows that the environmental factors are favorable for the normal growth and development of the species over time. Trees that are bigger in size also produce more volume compared with those with

smaller diameters and this further confirms the biological validity of the data. The results of the descriptive statistics (Table 2) were found compatible with the works of [14]; [20]; [21]; and [22].

Table 2: Summary statistics of measured parameters

Statistics	Dst(cm)	Dbh(cm)	Dm(cm)	Dt(cm)	Height(m)	BA(m ²)	Volume(m ³)
Mean	19.1	15.0	11.7	8.2	6.8	0.019	0.0963
Min	11.1	8.6	7.5	5.0	3.4	0.006	0.0178
Max	42.7	31.8	27.5	15.0	12.5	0.080	0.6968
SD	4.41	3.61	2.98	1.93	1.22	0.01	0.07
SE	0.16	0.13	0.11	0.07	0.05	0.00	0.00

Height-diameter Relationship

Different approaches were adopted by numerous researchers in developing height-diameter models. Chapman-Richards and Weibull functions are flexible and easier to work with, as such were considered for this study. Figure 1 shows the scatter relationship between height and Dbh using the actual field data before model fitting. From the scatter plot it shows non-uniformity of the height distribution especially as the trees grow taller, this is as a result of less competition at the upper level of the stands. Consistent increment in height with the increase in diameter was noted up to the 20 cm in diameter. Similar relationships were observed by [23] and [24]. They all believe the changes to be as a result of genetic materials and environmental factors that influence growth and development of the trees.

Both Chapman-Richards and Weibull have three parameters in the fit models and the iterative search of the parameter estimates in SAS software revealed a relatively higher Standard Error (SE) for Weibull estimates compared with the Chapman-Richards (Table 3) and this could be as a result of using asymptote in Weibull function for estimating height from diameter. The two functions used for modelling the relationship between height and diameter are nonlinear in nature, as such the R^2 value from ordinary least squares (OLS) cannot be used in comparison of the better model, therefore, Pseudo R^2 was used together

with Residual Mean Square Error (RMSE) as the basis for selecting the better model. From the goodness-of-fit statistics (Table 4) it is evident that, the Pseudo R^2 values are low for both Chapman-Richards and Weibull functions. This is a confirmation that height-diameter relationship may be influenced by other factors and the height-diameter modelling will yield more convincing result if such factors as; stand level density, spacing, age, and some climatic variation are added to the model. However, such information cannot be obtained in a single measurement, there is a need for a permanent sample plots that will be used for re-measurements over time. [23] developed height-diameter models with the inclusion of covariates and reported a better goodness-of-fit statistics compared with the models that exclude covariates. [5] developed mixed-effects models for estimating height from diameter; they first derived the models from each stand data as fixed effect before combining the stand attributes as mixed-effects. The resulting statistics revealed that mixed-effects models performed better in terms of goodness-of-fit than the fixed-effect models. Similar results were obtained by [2].

It is difficult to select a single model for all the species since they possess different attributes and the goodness-of-fit statistics are very close to each other for all the species. Like many studies ([25]; [2]; [5]; and [24]) that suggest the use of Chapman-Richards over other functions for height-diameter modelling, results of this study also revealed Chapman-Richards as being a better model for modelling Teak species height-diameter relationship considering the lower SE values. To account for the variation between different stand conditions, [24] adopted the methods of incorporating stand variables into height-diameter model as described by [26].




Figure 1: Height-diameter relationship for teak

Table 3: Height-diameter parameter estimates

Model	Parameters	Estimates	SE
Chapman-Richards	β_1	0.4729	0.15
	β_2	1.0624	0.18
	β_3	0.9706	0.01
Weibull	asymptote	9.6740	1.05
	b	11.912	2.37
	c	0.9303	0.17

β_1 , β_2 and β_3 are asymptote; rate and shape parameters, b and c are scale and shape parameters

Table 4: Goodness of fit statistics

Model	Pseudo R^2	RMSE
Chapman-Richards	0.3359	0.9946
Weibull	0.3353	0.9950

*RMSE = Residual Mean Square Error

CONCLUSION

This study revealed that the growth performance of trees in terms of tree taper has followed a normal distribution pattern, with diameter being reduced from bottom to the top of the trees. Goodness-of-fit test for height-diameter relationship shows lower Pseudo R^2 value which indicates weaker relationship between height and diameter without inclusion of other site factors. Height growth tends to decline as the trees grow bigger in size, which is as a result of lesser competition at the upper level of the stand distribution. Chapman-Richards function appears to be a better model for Teak height-diameter modelling. Other stand variables like density, spacing, age and management practices should be incorporated in order to improve the accuracy and reliability of height-diameter models.

REFERENCES

- [1] Hall, D.B. and R.L. Bailey (2001). Modeling and prediction of forest growth variables based on multilevel nonlinear mixed models. *Forest Science*, 47(3): 311-321.
- [2] Sharma, M. and J. Parton (2007). Height-diameter equations for boreal tree species in Ontario using a mixed-effects modeling approach. *Forest Ecology and Management*, 249(2007): 187-198.
- [3] Huang, S., S.J. Titus, T.W. Lakusta, and R.J. Held (1994). Ecologically based individual tree height-diameter models for major Alberta tree species. Alberta Environmental Protection. Land and Forest Service. Forest Management Division. Report #2. 27pp.
- [4] Fang, Z. and R.L. Bailey (1998). Height-diameter models for tropical forest on Hainan Island in southern China. *Forest Ecology Management*, 110:315-327.
- [5] Jiang, L. and Y. Li (2010). Application of nonlinear mixed-effects modeling approach in tree height prediction. *Journal of Computers*, 5(10): 1575-1581.
- [6] Ahmadi, K., S.J. Alavi, M.T. Kouchaksaraei, and W. Aertsen (2013). Non-linear height-diameter models for oriental beech (*Fagus orientalis* Lipsky) in the Hyrcanian forests, Iran. *Biotechnol. Agron. Soc. Environ.*, 17(3): 431-440.
- [7] Oyamakin, S.O., A.D. Fajemila, and S. Abdullateef (2013). Parameter estimation of height-diameter relationships of *Gmelina arborea* Roxb. (Family Verbenaceae). *Agriculture and Biology Journal of North America*, 4(4): 468-475.

- 231 [8] Calama, R. and G. Montero (2004). Interregional nonlinear height-diameter model with
232 random coefficients for stone pine in Spain. *Canadian Journal of Forest Research*, 34:
233 150-163.
- 234 [9] Sharma, M. and S.Y. Zhang (2004). Height-diameter models using stand characteristics
235 for *Pinus banksiana* and *Picea mariana*. *Scand. J. For. Res.*, 19: 442-451.
- 236 [10] Bermejo, I., I. canellas, and A. San Miguel (2004). Growth and yield models for teak
237 plantation in Costa Rica. *Forest Ecology and Management*. 1(31).
- 238 [11] Tewari, D.N. (1992). *A Monograph on Teak (Tectona grandis)*. International book
239 distributors. Hehra Durii: India.
- 240 [12] Pandey, D. and C. Brown (2000). La Taca: Una vision global. *Unasyuva*. 201(51): 5-17.
- 241 [13] Krishnapillay, B. (2000). Silviculture and management of teak plantations. *Unasyuva*.
242 201(51): 5-17.
- 243 [14] Adegbehin, J.O. (2002). Growth and yield of *Tectona grandis* (Linn. f) in the Guinea
244 and Derived Savanna of Nothern Nigeria. *International Forestry Review*, 4(1): 66-76.
- 245 [15] FDALR (1990). The reconnaissance soil survey of Nigeria. A technical report.
- 246 [16] D'hoore, J.L. (1964). *Soil Map of Africa*. Scale 1:500,000 Explaratory Monograph.
247 Lagos: C.C.T.
- 248 [17] Richards, F.J. (1959). A flexible growth function for empirical use. *Journal of*
249 *Experimental Biology*, 10(2): 290-300.
- 250 [18] Chapman, D.G. (1961). Statistical problems in dynamics of exploited fisheries
251 population. In: Neyman, J. (Ed.), *Proceedings of 4th Barkeley Symposium on*
252 *Mathematical Statistics and Probability*, 4:153-168
- 253 [19] Yang, R.C., A. Kozak, and J.H.G. Smith (1978). The potential of Weibull-type functions
254 as flexible growth curve. *Canadian Journal of Forest Research*, 8: 424-431.
- 255 [20] Shamaki, S.B., S.O. Akindele, and A.D. Isah (2011). Development of volume equations
256 for teak plantation in Nimbia Forest Reserve in Nigeria using Dbh and height. *Journal*
257 *of Agriculture and Environment*, 7(1): 71-78.
- 258 [21] Shamaki, S.B. and S.O. Akindele (2013). Volume estimation models from stump
259 diameter for for teak plantation in Nimbia Forest Reserve, Nigeria. *Journal of*
260 *Environmental Science and Water Resources*, 2(3): 89-94.
- 261 [22] Shamaki, S.B. and M. Ibrahim (2013). Volume estimation models using dbh and height
262 for *Gmelina arborea* (Roxb.) plantation in Nimbia Forest Reserve, Nigeria. *Biological*
263 *and Environmental Sciences Journal for the Tropics*, 10(4): 14-19.

- 264 [23] Guimaraes, M.A.C., N. Calegario, L.M.T. Carvalho, and P.F. Trugilho (2009). Height-
 265 diameter models in forestry with inclusion of covariates. *Cerne*, 15(3): 313-321
- 266 [24] Krisnawati, H., Y. Wang, and P.K. Ades (2010). Generalized height-diameter models for
 267 *Acacia mangium* Willd. Plantations in South Sumatra. *Journal of Forestry Research*,
 268 7(1):1-19.
- 269 [25] Liu, Z. and F. Li (2003). The generalized chapman-Richards function and applications to
 270 tree and stand growth. *Journal of Forestry Research*, 14(1): 19-26.
- 271 [26] Staudhammer, C. and V. LeMay (2000). Height prediction equations using diameter and
 272 stand density measures. *Forestry Chronicle*, 76: 303-309.