

Short Research Article**Geoengineering Biodiversity: Study to access feasibility of geoengineering techniques on biodiversity****Abstract**

As Earth continues to experience increased global warming, biological species on Earth are pushed to constantly modify and adapt to the changing Earth's climate in order to survive. This environmental pressure might push some of the more fragile species toward the brink of extinction thus human interventions are deemed necessary to minimize the current and future impacts of climate change. Current interventions include mitigation through international policies and regional laws by reducing anthropogenic outputs into Earth's climate system as well as conservation efforts to slow down extinction rate. Here, we will discuss how geoengineering can be added into one of these human interventions to reduce impacts of climate change on Earth's biodiversity. Geoengineering might provide immediate and simple solutions to current climate problems, however, only few researches have been conducted to study impacts and feasibility of geoengineering on life on Earth. In this paper, discussion on the feasibility and impacts of geoengineering on biodiversity will be assessed for two main techniques of geoengineering; carbon dioxide removal (CDR) and solar radiation management (SRM).

Keywords: *Biodiversity – Geoengineering – Climate Change – Carbon Dioxide Removal – Solar Radiation Management*

Introduction

Climate change has been a continuously ongoing issue that governments and international bodies have yet to find the feasible permanent solutions. Since the last few decades, Earth has experienced some of the worst climate variability, which not only affected humanity, but also the resiliency of other biological species on Earth. Diffenbaugh and Field (2013) argued that over the past few decades, biological species on Earth experienced increase pressure to undergo modifications in their genotype as well as phenotype through a process of adaptive radiation in order to continue surviving to compensate extreme climate variability on Earth.

Intergovernmental Panel on Climate Change (IPCC) reported that with current trajectory of global annual and cumulative emission of greenhouse gases (GHGs), Earth will experienced an increase of 1.4°C to 5.8°C in global mean temperature within the next few decades (2). One study indicated even higher global mean temperature rises of 4°C to 8°C towards the year 2100 (3). In addition, IPCC projected that current carbon dioxide (CO₂) concentration, which often use as indicator, will also observed increase within the next decades, ranging between 478-1099 parts per million (ppm) by the year 2100 (2). Projected increase of CO₂ concentration will not

only add burden on current GHGs concentration in the atmosphere but is deemed sufficiently enough to cause major ocean acidification leading to acceleration of another great extinction of major marine species as well as incur significant losses of ocean resources, which may be absolute detrimental to countries that are highly-dependent on these resources (4, 5).

[Fig.1] – Refer to supplementary materials

Figure 1 illustrates the global average temperature trend since 1890. An overall rising trend is observed throughout the dataset and through history only one of Earth warmest year have been attributed to ocean-atmospheric coupling feedbacks such as El-Nino Southern Oscillation (ENSO) (7). Kahn (2015) further added that majority of Earth warmest years on records were due to anthropogenic global warming. Additional data from Japan's meteorological agency support Kahn's observations that Earth four out of five warmest recorded years are within the last 10 years and does not corresponded with ENSO or other natural related forcing (7).

To address the rising trend of temperature and CO₂ concentration, major international bodies and some regional governments have directed mitigation procedures to reduce current and future anthropogenic burden on climate and biodiversity. These include implementation of international policies, protocols, conservation efforts, and regional laws directed towards managing localized anthropogenic activities and ensuring biodiversity resiliency (8,9).

Suggested mitigation procedures have shown some level of effectiveness throughout the implementation stage (8,9). However, with continuous trending in global economies, increasing population size, and increasing need for additional resources can provide barrier to the successfulness of the suggested mitigation procedures in the future (8,9). The Royal Society's report on geoengineering (2009) stated that unless all mitigations procedures are deemed reliable and able to exhibit a highly successful implementation, further interventions are needed to address the increase impacts of climate change on humanity as well as biodiversity.

Geoengineering

The concept of geoengineering has been around since 1965 when then President of United States, Lyndon Johnson, through his scientific advisors, suggested in altering the earth's surface using reflective particles in order to reflect additional sunlight, which at the time caused by the increase in carbon dioxide concentration in the atmosphere (23). As we continue to experience highly variable climate, increasing global warming, and further biodiversity loss, human manipulation of climate is deemed feasible for Earth to be habitable and perhaps resilience as we progress towards the future.

Geoengineering of climate can be divided into two main components consisting of carbon dioxide removal (CDR) techniques as well as solar radiation management (SRM) techniques (8). These two techniques posed unique types of interventions in which CDR intervened by removing greenhouse gases particularly CO₂ from the atmosphere and SRM intervened by modifying Earth's albedo through offset of GHGs and the use of reflectors (8, 9). Table 1 illustrates main methods of CDR and SRM techniques.

[Table 1] – Refer to supplementary material

For this particular article, we will be selecting some of both CDR and SRM methods of which would likely to give impactful result on biodiversity resiliency. In addition, we will also evaluate the feasibility of each method chosen and rate them on how well these methods help in the conservation of biodiversity. We hypothesized that with human control climate through these selected methods, much of Earth's biodiversity of both terrestrial and marine will be conserved and the feasibility of the next great extinction can be reduced. Methods chosen are (1) ocean fertilizations and (2) afforestation from CDR technique as well as (1) enhanced albedo and (2) aerosol injection from SRM technique.

CDR Option 1: Ocean Fertilization

Ocean fertilization uses nutrients such as iron, phosphorus, and nitrogen to encourage the growth of phytoplankton (enhancement of algal bloom) to sequester carbon from the atmosphere (9). The mechanism involved a simple photosynthesis processes of which phytoplankton will retained carbon within their cells during photosynthesis for processing food and when they die, remaining carbon embedded within their cells will be confined and sequestered in the deep ocean (9). Few studies had approached ocean fertilization method and most have suggested that dissolved nutrient, particularly iron, into the ocean has led to some extent of carbon dioxide removal from the atmosphere (9-11). It is suggested that a ton input of iron into the ocean is able to sequester roughly 30,000 to 110,000 tons of CO₂ (9). Although ocean fertilization helps to reduce atmospheric CO₂ concentration, other studies have also found that by increasing carbon sequestration in the deep ocean, the process can contribute to the increase of ocean acidification of which in return, could result in the disruption of marine biodiversity through coral bleaching and decalcification of shelled molluscs (5, 9, 12, 24). In addition, McCauley et al. (2015) argued that intentional chemical introduction into the ocean may accelerate the extinction of several sensitive marine biodiversity. Another study by Blain et al. (2007) further added that effective carbon sequestration through phytoplankton bloom should not be used as a method to the geoengineering CRD technique due the method being highly unsustainable in the long run.

CDR Option 2: Afforestation

Afforestation involves planting trees seedling on open land to naturally sequester CO₂ from the atmosphere (9). The method is deemed as a prime geoengineering technique and a great climate change mitigation strategy as forest can stored huge amount of CO₂ up to 2.2 to 9.5 metric tons per acre per year (9). Several studies indicated that afforestation is one of the important indicators to project future CO₂ concentration and can be one the most cost-effective strategies for any climate mitigation methods (9, 13, 14). These studies also include benefits of afforestation such as erosion control, wildlife habitat reclamation, and can be used for recreational purposes (9, 13, 14). Furthermore, in Woziwoda and Kopec (2014) study, they observed that afforestation had been fairly effective in conservation of plant species diversity over an abandoned exposed peatlands, thus in essence conserving local biodiversity whilst minimizing CO₂ release into the atmosphere from these peatlands. Although afforestation tends to be a best solution to mitigate climate change, there are drawbacks associated with the method. These drawbacks may include accidental CO₂ released due to drought or forest fires, lands reclamation for crop-based industries, and land reclamation for residential areas for growing human population (9). Other than that, there are existence of economically driven interests of certain parties on how much will afforestation might cost them their livelihood and source of income (25). Biodiversity-wise, possible drawbacks may include displacement and extinction of original species due to introduction of newer species through afforestation of which can grow and populate afforested area faster than original species (15).

SRM Option 1: Enhanced Albedo

Enhanced albedo consists of increasing reflectivity of solar forcing through brightening of Earth's surfaces and cloud whitening using addition of cloud-condensation nuclei (CCN) (9). This method utilized the concept of solar radiation reflectivity of which by increasing Earth's albedo, more incoming solar radiation will be reflected back to space thus limiting further temperature increase (9). The Royal Society (2009) reported that in order to radiatively cools Earth of approximately 4W/m², we need to modify current surface albedo so that current total solar radiation reflection increases from ~107 W/m² to ~111W/m². For this, a novel idea was proposed to increase surface albedo by using vegetated surfaces through genetically modifying plants' genomes in order to change their surface appearances into brighter or almost reflective in nature (8). Furthermore, multiple studies conducted had shown that albedo-engineered crops are able to influence regional climate variability by influencing local solar radiation forcing (16, 17). Figure 2 illustrates global agriculture distribution and cooling associated with albedo-engineered crops. Diagrams modeled on the situation whereby all major agriculture crops are albedo-engineered and it is suggested that the potential cooling during summer solstice will be approximately 1°C throughout North America and midlatitude Eurasia (17). Lenton and Vaughan (2009) added that by modifying grassland and cropland albedo, radiative forcing could be reduced up to -0.83W/m². Although model presented is viable, impacts of this method on biodiversity are alarming. Albedo-engineered crops might change the region landscape thus altering ecological behavior surrounding

new crops production (16). In addition, Rigdwell et al. (2009) argued, bio-geoengineering crops can cause extreme drought in part of subtropical countries as well as modify localized soil contents in certain regions around the globe.

SRM Option 2: Aerosol Injection

Aerosol injection is a deliberate attempt to introduce reflective chemical droplets into the atmosphere (9). Implementation of this method requires aircraft to disperse droplets into the stratospheric layer of the atmosphere where strong mixing of gases occurs (8, 9). These droplets are of the same chemical species produced during volcanic eruptions, particularly of hydrogen sulphide (H₂S) and sulphur dioxide (SO₂) (8, 9). Wigley (2006) states that volcanic eruption, drawing example of Mount Pinatubo 1991 incident, provide great scenario of which scientists can study the feasibility of aerosol injection into the stratosphere to modify global climate. Illustrated by Figure 1, it is observed that during the eruption of Mount Pinatubo, global temperature has dropped for approximately two years before regaining back previous rising momentum. Wigley (2006) further added that this eruption not only contribute to a period of global cooling, but is deemed stable as climate has not been seriously affected by eruptive particles. Figure 3 illustrates the possibility of reduction of incoming solar radiation based on numerical model in which three possibilities of geoengineering manipulations are studied depending on how frequent aerosol injection is introduced into the atmosphere (4). By using this model, the Royal Society (2009) estimated that aerosol injection able to reduce incoming solar radiation by 1.84%. As for biodiversity, geoengineering climate using aerosol injection may provide huge array of problems for life on Earth. Multiple studies indicated that the adverse effects of aerosol injection include increase of ocean acidification, disruption of regional precipitation, and increase frequency of acid rains (4, 8, 19). Other studies focusing on evaluating biodiversity to historical volcanic eruption have also found that species tend to go extinct after major volcanic eruptions and subsequently replaced by other species that able to adapt and survive, thus completely changing the region ecosystem (19, 20).

[Table 2] – Refer to supplementary material

Discussion

Geoengineering has been mostly targeted to tackle climate change, however, it is also best to say that geoengineering can be an essential tool to tackle other dependent of climate change such as biodiversity. Each method presented of either CDR or SRM techniques has their own strengths and weaknesses depending on how they are implemented globally and how they affected biodiversity. In term of accessing feasibility of these four methods with biodiversity, most of methods presented are somewhat ignoring the impacts they have on biodiversity itself as they focus solely on the objectives to reduce global warming. Nonetheless, to

be neutral, it can be argued that since there are only few researches that had been dedicated on finding biodiversity implications through geoengineering techniques, this may limit the factual as well as theoretical evidences of each method used in CDR and SRM techniques against biodiversity.

Out of all four methods discussed, only afforestation of CDR technique has so far fit with stated hypothesis. It has been argued that, afforestation has not only helps to geoengineer climate through CO₂ sequestration but the method may also helps in the conservation of biodiversity, thus making biodiversity resilience towards the future and become more sustainable (9). Lang et al. (2014) stated that afforestation has not only benefiting the wood industries but it is a win-win system of which biodiversity is enriched through carbon sequestration, nutrient retention, and groundwater recharge. In addition, afforestation method is one of the most affordable ways to geoengineered the climate and reclaimed used lands with estimated cost of USD65 to USD200 per acre depending on species planted and regional locations (8, 9). However, it is worth to keep in mind that despite afforestation may helps in conserving biodiversity as a whole, some biological species within the afforestation areas can be disturbed and displaced thus altering original ecosystem interaction (15, 22). Ridgwell et al. (2009) argued that as long changes can make impacts on climate (and biodiversity), sacrificing few for that change is deemed necessary. So far, safety risks associated with afforestation are generally kept at minimal level (8). Future suggestion for geoengineering research is to include necessary section on biodiversity against impacts of individual methods proposed in the geoengineering techniques, both CDR and SRM.


References

1. Diffenbaugh, N. S., & Field, C. B. (2013). Changes in Ecologically Critical Terrestrial Climate Conditions. *Science*, 341, 486-492. doi:10.1126/science.1237123
2. Intergovernmental Panel on Climate Change. (2001). *Climate Change 2001: Impacts, Adaptation, and Vulnerability*. In J. J. McCarthy, O. F. Canziani, N. A. Leary, D. J. Dokken, & K. S. White. (Eds.), Scenario of the 21st Century (pp. 175-179). Cambridge, United Kingdom: Cambridge University Press.
3. Zickfeld, K., Morgan, M. G., Frame, D. J., & Keith, D. W. (2010). Expert Judgements about transient climate response to alternative future trajectories of radiative forcing. *Proceeding of the Natural Academy of Science*, 107(28), 12451-12456. doi:10.1073/pnas.0908906107
4. Wigley, T. L. (2006). A combined mitigation/geoengineering approach to climate stabilization. *Science*, 314(5798), 452-454. doi:10.1126/science.1131728

- 174 5. McCauley, D.J., et al. (2015) Marine defaunation: Animal loss in the global ocean. *Science*, 347 (6219):247, 1255641-1
175 doi:10.1126/science.1255641
- 176 6. Global Average Surface Temperature Anomalies. (2015). In *Japan Meteorological Agency*. Retrieved March 20, 2015.
- 177 7. Kahn, B. (2015). 2014 Officially Hottest Year on Record. In *Scientific American*. Retrieved March 20, 2015.
- 178 8. The Royal Society. (2009). *Geoengineering the climate: Science, governance and uncertainty* (pp. 1-81). London, United
179 Kingdom: Author.
- 180 9. Congressional Research Service. (2013). *Geoengineering: Governance and Technology Policy* (pp. 1-39). Washington, DC:
181 Author.
- 182 10. Coale, K.H., et al. (1996). A massive phytoplankton bloom induced by an ecosystem-scale iron fertilization experiment in the
183 equatorial Pacific Ocean. *Nature*, 383, 495-501. doi:10.1038/383495a0
- 184 11. Blain, S., et al. (2007). Effect of natural iron fertilization on carbon sequestration in the Southern Ocean. *Nature*, 446, 1070-1075.
185 doi:10.1038/nature05700
- 186 12. Ericson, J.A., et al. (2012). Combined effects of two ocean change stressors, warming and acidification, on fertilization and early
187 development of the Antarctic echinoid *Sterechinus neumayeri*. *Polar Biology*, 35, 1027-1034. doi: 10.1007/s00300-011-1150-7
- 188 13. Sheng-wei, S., Peng-fei, H., Ping, Z., & Cheng-lin, M. (2015). The Impact of Afforestation on Soil Organic Carbon Sequestration
189 on the Qinghai Plateau, China. *PLoS ONE*, 10(2), 1-18. doi:10.1371/journal.pone.0116591
- 190 14. Pan, Y., et al. (2011). A Large and Persistent Carbon Sink in the World's Forests. *Science*, 333, 988-993. doi:
191 10.1126/science.1201609
- 192 15. Woziwoda, B., and Kopeck, D. (2014). Afforestation or natural succession? Looking for the best way to manage abandoned cut-
193 over peatlands for biodiversity conservation. *Ecological Engineering*, 63, 143-152. doi:10.1016/j.ecoleng.2012.12.106
- 194 16. Knorr, W., and Schnitzler, K-G. (2006). Enhanced albedo feedback in North Africa from possible combined vegetation and soil-
195 formation processes. *Climate Dynamics*, 26, 55-63. doi: 10.1007/s00382-005-0073-9
- 196 17. Ridgwell, A., Singarayer, J.S., Hetherington, A.M., and Valdes, P.J. (2009). Tackling Regional Climate Change by Leaf Albedo
197 Bio-geoengineering. *Current Biology*, 19(2), 146-150. doi:10.1016/j.cub.2008.12.025
- 198 18. Lenton, T.M., and Vaughan, N.E. (2009). The radiative forcing potential of different climate geoengineering options.
199 *Atmospheric Chemistry & Physics Discussions*, 9, 2559-2608. doi: 10.5194/acp-9-5539-2009, 2009
- 200 19. Barash, M.S. (2011). Causes of Mass Extinction of Sea Organisms at the Paleozoic–Mesozoic Boundary. *Doklady Earth Sciences*,
201 438(2), 750-753. doi: 10.1134/S1028334X11060286
- 202 20. Jozsef, P., and Kocsis, A.T. (2014). Volcanism of the Central Atlantic magmatic province as the trigger of environmental and
203 biotic changes around the Triassic-Jurassic boundary. *Special Paper - Geological Society of America*, 505, 245-261. doi:
204 10.1130/2014.2505(12)
- 205 21. Lang, A.C., et al. (2014). Mixed afforestation of young subtropical trees promotes nitrogen acquisition and retention. *Journal of*
206 *Applied Ecology*, 51(1), 224-233. doi: 10.1111/1365-2664.12157

22. Souza, A.F., Ramos, N.P., Pizo, M.A., Hubel, I., and Crossetti, L.O. (2013). Afforestation effects on vegetation structure and diversity of grasslands in southern Brazil: The first years. *Journal for Nature Conservation*, 21(1), 56-62. doi: 10.1016/j.jnc.2012.10.002
23. Bielo, D. (2010). What is geoengineering and why is it considered a climate change solution. In *Scientific American*. Retrieved on 20th October 2017.
24. Gazeau, F., et al. (2013). Impacts of ocean acidification on marine shelled molluscs. *Marine Biology*, 160(8), 2207-2245. doi:10.1007/s00227-013-2219-3
25. Ryngaert, C. (2016). Climate change mitigation techniques and international law: Assessing the externalities of reforestation and geoengineering. *Ratio Juris*, 30(3), 273-289. doi:10.1111/raju.12154

229 **Supplementary Materials**



230

231 **Fig 1.** Annual Global Average Temperature from 1890 to 2014. Recent temperature anomalies are calculated using deviation of
 232 baseline (1980-2010) average. Arrows (red and blue) represent five warmest years on record (1998, 2005, 2010, 2013, 2014). Red
 233 arrow indicates warm year due to 1998 ENSO. Orange arrow indicates cooling after Mount Pinatubo eruption in 1991. Black line
 234 represents actual temperature anomaly based on yearly record. Blue line represent average 5 years running mean of temperature
 235 anomaly, and red line represent long-term trend linear anomaly of annual global average temperature. Dataset provided by Japan
 236 Meteorological Agency (6).

237

Techniques	CDR	SRM
Methods	▪ Ocean fertilizations ▪ Carbon capture and sequestration ▪ Afforestation ▪ Enhanced weathering	▪ Enhanced albedo (surface and cloud) ▪ Aerosol injection (preferably hydrogen sulphide, H₂S and sulphur dioxide, SO₂) ▪ Space reflector

Table 1. Suggested methods of CDR and SRM techniques of geoengineering. Table summarized complete methods outlined by The Royal Society’s report on geoengineering. Congressional Research Service’s report (2013).





Fig 2. Global crop distribution and cooling associated with albedo-engineered crops. **(Left)** Diagram illustrates Earth’s surfaces model if covered by albedo-engineered agriculture crops. **(Right)** Diagram illustrates temperature anomaly taken during December-January-February (DJF) and June-July-August (JJA) if global agriculture surface are covered with albedo-engineered crops. Note that patches above the Northern Hemisphere are corresponding to sea-ice feedback during winter and summer solstices. (17)

246



247

248 **Fig 3.** Reduction of incoming solar radiative forcing based on different geoengineering scenarios. LOW GEO indicates no direct
 249 human intervention on modification of climate. HIGH GEO indicates scenario of which released of H₂S and SO₂ through aerosol
 250 injection are similar to eruption of Mount Pinatubo for every 2 years. MID GEO indicates mid-level intervention by human whereby
 251 aerosol injections are introduced over longer interval period of time. (4)

252

253

254

255

Methods	Summary	Pro	Cons
Ocean fertilization (CDR)	- Done through addition of iron nutrient into the ocean to encourage growth of phytoplankton. As phytoplankton photosynthesize CO₂, carbon will be retained in their cells which will be sequestered as carbon in Deep Ocean when they die. - Estimated 30,000 to 110,000 tons of carbon removed from air.	Reduction of CO₂ in the atmosphere	- Ocean Acidification - Marine defaunation - Low effectiveness level
Afforestation (CDR)	- Done through planting tree seedlings. - Regarded as prime sequestration strategy of which forest is able to store carbon for longer time periods. - Able to modify local climate at large scale afforestation.	- Reduction of CO₂ in the atmosphere - Affordable	- Forest fire, drought will release CO₂ back into atmosphere. - Loss of original diversity. - Cost-effective
Enhanced albedo (SRM)	- Done through increasing reflectivity or certain surfaces. - Using reflective paints (white) on building surfaces. - Modification of plant to augment albedo. - Cloud whitening through dispersion of cloud-condensation nuclei (CCN).	- Limit temperature increase through enhanced solar radiation reflectivity. - Low safety risks	- Cost of implementation - Marginally effective - Displacement of species
Aerosol injection (SRM)	- Done through injection of aerosol particularly H₂S and SO₂ into the stratosphere. - Help to either direct solar radiation back to space or to absorb heat.	Limit temperature increase through enhanced solar radiation reflectivity or by absorbing	- Ocean acidification. - Irregular precipitation - Acid rain - Species extinction

		heat. ▪ Very effective	
--	--	---------------------------	--

257 **Table 2.** Methods explanation of each selected CDR and SRM techniques of geoengineering. Discussion includes small summary and
258 point of pros and cons of each method. Unless specified, points of pros and cons refer to how well methods feasible to geoengineering
259 (1-20)
260