

**EVALUATION OF HEAVY METAL CONTAMINATION OF SOILS ALONGSIDE
AWKA- ENUGU ROAD, SOUTHEASTERN NIGERIA**

ABSTRACT

Fourteen top soil samples were collected from old Awka-Enugu road passing through Obioma, Nsude, Ngwo and Abor communities. Samples were taken from a depth of 10-15cm after making a hole with a pick axe. Total concentrations of cadmium (Cd), copper (Cu), nickel (Ni), lead (Pb), zinc (Zn), cobalt (Co), and chromium (Cr) were determined using a Buck 210 VGP atomic absorption spectrophotometer (AAS). Soil contamination was evaluated with pollution index (Pi) and index of enrichment factor (EF) concept. The enrichment factor of copper (Cu) was observed to be generally low across the study area. This implies that Cu toxicity in the area is not significant. However, the enrichment and pollution indices of cadmium (Cd) and lead (Pb) were observed to be high within the study area. The study area was observed to be highly contaminated by cadmium and lead but moderately contaminated with respect to nickel, zinc and chromium. The enrichment and pollution trends of the metals in descending order shows $Cd > Pb > Ni > Zn > Cu > Cr > Co$. In summary, the total heavy metal concentrations of soils studied were slightly higher than previously reported levels for similar soils which suggest some degree of heavy metal pollution.

Keywords: Pollution index, Enrichment factor, Heavy metals, Contamination, Health Risk Level.

1.0 INTRODUCTION

Surfaces such as pavements and road surfaces in urban environments usually contain an accumulation of fine solid particles on them. These particles deposited on these surfaces are commonly referred to as street dust, road deposited sediments (RDS) or road dust. Road deposited sediments are usually a complex mixture of fine solid particles and pollutants obtained from a wide range of urban and industrial sources and processes. Their sources are either as a result of human activities (vehicle tyre and body wear, vehicular exhaust emissions brake-lining materials, road salt, building and construction materials or soil material particles, litter obtained from plant and leaf and atmospheric deposition [1-5]. Roads are known as the second largest non-point source of creating pollution in urban environments. Road and highway surfaces are impervious, and serve as temporary sinks for various types of pollutants that are washed off during rainfall to the surrounding environments [6-8]. Road surfaces receive varying amount of heavy metals by the process of atmospheric deposition, sedimentation, impaction and interception (Li et al., 2001). In urban environments, the top soils and road deposited sediments (RDS) or road dust are indicators of heavy metal contamination from atmospheric deposition. Industries, traffic, mining activities, smelters and construction are some of the main anthropogenic sources of this heavy metal pollution. The traffic source includes vehicles (tyre wear, brake linings, fuel combustion, etc.) while road infrastructure (pavement wear, corrosion of galvanized steel crash barriers, etc) contribute the rest. It has been reported that pollutants such as arsenic, cadmium, chromium, copper, nickel, lead and zinc generally have high concentrations on road surfaces, roadside soils and

particulate matter due to heavy traffic, and most often affect the environmental air quality (Culbard et al.,1988).

Roads in the study area play a major role in stimulating socio-economic progress and development. Major activities of the inhabitants of the study area include sales of vehicle parts, repairs and scavenging of useful motor parts from scrapes. Wastes emanating from such activities could lead to contamination of surficial materials including soils. Contamination by heavy metals in any environment is a major concern because of their toxicity, threat to life and the ecosystem. Automobile mechanic workshop wastes have been reported by earlier scholars for the elevated concentrations of cadmium (Cd), chromium (Cr), copper (Cu), lead (Pb), nickel(Ni), and zinc(Zn) in soil profiles in the vicinity of automobile waste dumps in Nigeria [11]. Remarkable metal contamination especially lead pollution has also been attributed to automobile emissions in the roadside ecosystems [12]. Most researchers have reported the influence of the traffic load on heavy metal contents in top soil and their variability with distance [13-14].

Viard *et al* [7] reported that the deposition and the levels of metals in surficial soils decrease with increasing distance from the highways. They study also revealed that the impact of the highway was clear up to 320m at least for lead, but the most important contamination was observed near the road. Nabuloa *et al* [16] reported that leaves of roadside crops can bio-accumulate trace metals at high concentrations causing serious health risk to consumers of such products. Hence, accurate evaluations of the heavy metal concentrations of polluted soils are required to assess the potential ecological risk of affected areas. The potential health risks and high levels of heavy metal pollution measured in a number of cities of the world have therefore generated a lot of interest in heavy metal pollutions in road deposited sediments. Roadways and automobiles are presently considered as the largest sources of heavy metals in the urban environment. The objectives of this study are to assess the heavy metal contamination in roadside top soils along the old road passing through Obioma- Nsude – Ngwo and Abor areas of Enugu State, Southeastern Nigeria and to compare the observed levels of pollution in the area with established standards of unpolluted soils. It is envisaged that the study would provide background data for subsequent investigation of the environment, biological dose response and effect on human health.

1.1.BACKGROUND GEOLOGY OF THE STUDY AREA

The study area lies within longitudes $7^{\circ}20'$ – $7^{\circ}22'E$ and latitudes $6^{\circ}21'$ – $6^{\circ}30'N$ (Fig 1). The Night Mile Corner, Ngwo-Uno area, has been the centre of industrial and commercial activities in recent times, particularly those industries that require viable and efficient water supply systems. This is because the study area has a high groundwater potentials when compared to other surrounding areas which is often attributed to the local hydrogeological characteristics of the area. The study area is underlain by the Ajali Formation which is known for its huge aquiferous potentials. The Ajali Formation is made of well sorted, massive, coarse –grained, white sandstones with occasional intercalations of thin beds of white shales and clays. Rivers Ekulu, Iva, Oroho and Nyaba take off from the base of the Ajali Formation on the Enugu escarpment descending through the underlying Mamu Formation into Enugu town (Fig 1). The Mamu Formation trends northeast (N-E) of the study area (Fig 1) and comprises of alternations of sandstones, dark-blue or grey shales, sandy shales and mudstones with local coal seams at various horizons [17].




Figure 1. Location and geological map of the study area

2.0 MATERIALS AND METHOD

Fourteen (14) top soil samples were systematically collected along Awka-Enugu old road passing through Obioma- Nsude - Ngwo - Abor communities which stretch about 17 kilometers (Fig 1). The sampled soil layer partly contains humus with completely decomposed non-identifiable organic materials. Sample numbers 4, 5, 6, 7, 9, 10, 11 and 13 were collected from the edge of the road while sample numbers 1, 2, 3, 8, 12, and 14 were collected farther away off the road (3-150m). Each sample was taken from a depth of 10-15cm after making a hole with a pick axe from the surface. Site locations were recorded and geo-referenced using a standard Garmin eTrex global positioning system (GPS) instrument. The sample coordinates, names and descriptions are presented in table 1. Soils collected varied in texture, colour and composition from clay to sand with some samples having gray, red and brown colours. A good vegetation cover was noticed with *Imperata cylindrica* (tall grasses) being dominant. Soil pH which is a measure of the acidity or basicity considered as a master variable was measured by the use of an electronic pH-meter (Table 2).

The soil samples were air-dried, disaggregated and mechanically sieved using 100-mesh screen. Samples are considered acceptable if 100% passes through 100-mesh screen [18] and also to ensure homogeneity prior to digestion[19]. Digestion of soil samples were carried out according to standard procedures as described by [20] and [21], where 0.2 grams of each sample was weighed and dried using crucibles. The samples were digested in acid mixtures of 5 ml nitric acid, 3 ml perchloric acid and 2 ml hydrofluoric acid [20]. They were placed on hot plates for ten minutes and later transferred to a fume chamber where they were allowed to cool. The cooled solutions were made up to 250 ml in volumetric flasks with de-ionized water. The obtained solutions were later used for the measurement of heavy metals including

113 Lead (Pb), Copper (Cu), Chromium (Cr), Nickel (Ni), Cadmium (Cd), Zinc (Zn) and Cobalt
114 (Co) using the Buck 210 VGP atomic absorption spectrophotometer.

115 3.0 RESULTS INTERPRETATION AND DISCUSSION

116 Heavy metal concentrations are expressed in milligram per kilogram (mg/kg) of dry weight
117 of the soil samples. The summary of parameters of the analytical data of the soil samples are
118 shown in table 2. The enrichment factor (EF) and pollution index (PI) are presented in tables
119 3 and 4 respectively while table 5 shows the range and mean values of heavy metal contents
120 (mg/kg) of soil samples and related established background values from [22-25] and [26] for
121 non-polluted soils.

122 **Table 1: Location of soil samples**

Sample Location	Latitude N(degrees)	Longitude E (degrees)	Remarks
Obioma -1	06°29'748''	07° 25' 186''	Fine grained, friable loose sand.
Obioma -2	06°21'803''	07° 25' 034''	Well sorted sand weathered to reddish-brown coloured clayey soil.
Obioma -3	06°22'299''	07° 25' 016''	Heavily leached dark loam soil.
Nsude-4	06°22'748''	07° 24' 189''	Fine grained friable sand and ironstone weathered to reddish laterite with concretions .
Nsude-5	06°23'334''	07° 24' 109''	Weathered brown/reddish ironstones, quartz pebbles which favours extensive growth of vegetation.
Nsude-6	06°24'497''	07° 24' 133''	Light brown, fine grained and well sorted sand.
Ngwo-7	06°25'406''	07° 26' 181''	Loam soil with thick vegetation cover with tall grasses.
Ngwo-8	06°25'019''	07° 26' 267''	Ironstone rubbles weathered to gray and brown coloured laterite.
Ngwo-9	06°26'536''	07° 26' 439''	Loose friable, gray coloured well sorted sand.
Ngwo-10	06°27'023''	07° 26' 422''	Brown weathered ironstones, quartz pebbles.
Abor-11	06°27'496''	07° 25' 350''	Heavily leached/weathered to reddish-brown coloured laterite and loam with scanty vegetation cover.
Abor-12	06°28'474''	07° 25' 414''	Silty and dark coloured sand
Abor-13	06°21'803''	07° 25' 034''	Ironstone boulders weathered to red coloured soil, favours growth of tall grasses.
Abor-14	06°29'748''	07° 25' 186''	Unconsolidated weathered sand on a flat terrain dotted with shrubs

123 **Table 2. Concentrations (mg/kg) of Metals in Soil Samples**

Sample Location	Concentration of Metals in mg/kg							
	Cd	Cr	Cu	Ni	Pb	Co	Zn	pH
Obioma-1	6.12	1.37	2.87	6.36	4.12	0.62	7.00	6.0
Obioma-2	5.25	1.50	2.50	1.50	0.87	0.00	7.00	6.0
Obioma-3	4.75	0.00	0.12	10.37	3.50	0.00	2.87	5.0
Nsude-4	4.62	4.00	0.12	6.75	1.75	0.00	3.62	6.0
Nsude-5	2.50	3.00	1.00	2.62	2.25	1.12	4.37	6.0
Nsude-6	1.12	0-00	0.75	10.1	1.87	0.62	1.87	6.0

Ngwo-7	1.12	0.00	0.12	4.00	4.75	0.37	4.00	6.0
Ngwo-8	0.72	1.12	1.87	0.87	3.25	0.00	6.00	5.0
Ngwo-9	0.25	3.75	2.25	11.12	0.50	0.37	5.00	6.0
Ngwo-10	0.00	3.62	0.87	11.00	2.25	0.87	0.00	5.0
Abor-11	1.50	0.00	3.75	6.62	1.37	0.25	3.50	6.0
Abor-12	1.62	2.12	0.62	11.8	2.62	0.37	2.62	6.0
Abor-13	1.12	5.12	1.12	1.62	3.50	0.87	7.25	6.0
Abor-14	0.37	2.37	2.00	0.75	3.75	0.25	6.25	6.0

124

Table 3. Enrichment Factors (EF) of Soil Samples.

Sample Location	Enrichment Factors (EF) of Metals						
	Cd	Cr	Cu	Ni	Pb	Co	Zn
Obioma-1	84.04	0.41	4.37	4.44	15.43	1.27	4.00
Obioma-2	120.22	0.76	6.36	1.74	5.43	1.27	6.67
Obioma-3	113.20	0.00	0.97	12.56	22.75	0.00	2.85
Nsude-4	175.86	3.38	0.50	13.06	18.16	0.00	5.74
Nsude-5	38.50	1.02	1.71	2.05	9.45	2.58	2.80
Nsude-6	19.04	0.00	1.41	8.74	8.67	1.58	1.32
Ngwo-7	21.01	0.00	0.25	3.81	24.31	1.04	3.12
Ngwo-8	24.81	0.85	7.16	1.52	30.35	0.00	8.61
Ngwo-9	5.17	1.72	5.17	11.70	2.82	1.14	4.31
Ngwo-10	0.00	1.53	1.83	10.64	11.67	2.48	0.00
Abor-11	28.45	0.00	7.90	6.38	7.08	0.71	2.76
Abor-12	32.13	0.93	1.36	11.97	14.17	1.10	2.16
Abor-13	22.47	2.28	2.49	1.65	19.15	2.61	6.06
Abor-14	9.95	1.41	5.97	1.02	27.51	1.00	7.00

125

Table 4. Pollution Index (Pi) of Metals in soil samples

Sample Location	Pollution Index (Pi) of Metals						
	Cd	Cr	Cu	Ni	Pb	Co	Zn
Obioma-1	76.50	0.01	0.10	0.10	0.37	0.03	0.09
Obioma-2	65.62	0.01	0.09	0.02	0.07	0.01	0.03
Obioma-3	59.37	0.00	0.01	0.17	0.31	0.00	0.03
Nsude-4	57.75	0.02	0.00	0.11	0.15	0.00	0.05
Nsude-5	31.25	0.02	0.03	0.04	0.20	0.05	0.06
Nsude-6	14.00	0.00	0.02	0.17	0.17	0.03	0.02
Ngwo-7	14.00	0.00	0.00	0.06	0.43	0.01	0.05
Ngwo-8	9.00	0.00	0.06	0.01	0.29	0.00	0.08
Ngwo-9	3.12	0.02	0.08	0.18	0.04	0.01	0.06
Ngwo-10	0.00	0.02	0.03	0.18	0.20	0.04	0.00
Abor-11	11.75	0.00	0.13	0.11	0.12	0.01	0.04
Abor-12	20.00	0.01	0.02	0.20	0.23	0.01	0.03
Abor-13	14.00	0.03	0.04	0.02	0.31	0.04	0.10
Abor-14	4.62	0.01	0.07	0.01	0.34	0.01	0.08

126

Table 5. Average and range values of heavy metal content (mg/kg) of all sampled sites and related to background values from different sources.


127

Metal	Values of sites (this study)	Background values from different sources				
		Bowen (1979)	Shacklette/Boerngen (1984)	Kabata-Pendias (1995)	Krauskopf (1979)	Rudnick/Gao (2003)
Cd	Min. 0.00	0.01	—	—	—	—

	Max. 6.12 Ave. 2.18	2.00 0.35	—	0.05	0.15	0.08
Cr	Min. 0.00 Max. 5.12 Ave. 1.99	42.0 329.2 124.5	54.00	—	100.00	135.00
Cu	Min. 0.12 Max. 3.75 Ave. 1.44	14.2 68.9 40.0	25.00	100.00	50.00	27.00
Ni	Min. 0.87 Max. 11.87 Ave. 6.11	54.6 378.0 157.8	19.00	30.00	75.00	59.00
Pb	Min. 0.50 Max. 4.75 Ave. 2.59	33.4 163.3 80.9	19.00	20.00	12.50	11.00
Co	Min. 0.00 Max. 1.12 Ave. 0.43	- - -	—	50.00	22.00	20.00
Zn	Min. 0.00 Max. 7.25 Ave. 4.41	187.9 1087.0 476.7	—	300.00	70.00	72.00

129

130



131

132

133


134

Figure 2. Spatial distribution of the concentrations of some heavy metals (mg/kg): (a) Cadmium(Cd) (b) Chromium(Cr) (c) Nickel(Ni) (d) Lead(Pb)

Table 2 and figure 2 shows the concentrations (mg/kg) of metals determined from the soil samples while table 5 shows the range of values used for comparison. Cadmium (Cd) in unpolluted soils is normally in the concentration range of 0.01 – 0.35 mg/kg according to tolerable standards by [22]. However, cadmium in table 2 ranged from <0.37 to 6.12 mg/kg with mean values exceeding the tolerable limits of 2.00 mg/kg as given by [22], 0.05 mg/kg by [24], 0.15 mg/kg by [25] and 0.08 mg/kg as given by [26]. It implies that Cd has impacted negatively on the agricultural soil and the environment at large. High cadmium concentrations in the study area may be a direct result of both natural and anthropogenic sources. Similarly, chromium ranged from < 1.12 to 5.12 mg/kg with an average value of 1.99 mg/kg. At sampling sites 3, 6, 7 and 11, chromium was not detectable. Chromium values and the mean value are much lower than the referenced standard levels [22-26]. Copper varied from 0.12 to 3.75 mg/kg, with a mean value of 1.44 mg/kg. Similarly, copper was lower than the established tolerable limits. This means that Cr and Cu may not have had any significant effect on the environment. Nickel ranged from 0.87 – 11.87 mg/kg with an average concentration of 6.11 mg/kg while cobalt ranged from 0.25 to 1.22 mg/kg with 0.43 mg/kg as the average value. Cobalt was not detected in some locations (Table 2). Average concentrations of Cr (1.99 mg/kg) and Co (0.43 mg/kg) were very low when compared with established limits of [26], [23] and [22] in table 5. They therefore pose no risk to health and the environment. However, the concentrations of lead (Pb) is slightly below toxic level. Lead concentrations across the study area ranged from 0.50 to 4.75 mg/kg with the highest concentrations revealed at sites 7 and 14. At these two locations, auto mobile mechanic workshops are located close by and are believed to have directly contributed to the high concentrations at the locations. Zinc concentrations ranged from <1.87 to 7.25 mg/kg with an average value of 4.41 mg/kg. Concentration sequence of the examined metals in descending order is presented below: Ni > Zn > Pb > Cd > Cr > Cu > Co (Table 2). The average concentration value of nickel (Ni) of 6.11 mg/kg is considered high but below the 50 mg/kg tolerable limit set by [22]. The toxicity of excess Ni present in the soil could account for the poor growth of weeds at some of the locations. Nickel is carcinogenic and affects the reproductive health of organisms [28]. The high level in the study area may be due to aerial deposition, manure and phosphate fertilizer application.

The concentration of an element in the soil solution is believed to depend on the equilibrium between soil solution and the solid phase, with pH playing a decisive role [29]. While discussing the relatively high mobility of heavy metals with regard to pH, [30] considered that in acid soils (pH 4.2 – 6.6) the elements: Cd, Ni, and Zn are highly mobile, Cr is moderately mobile, and Cu and Pb practically immobile. Soil pH in this study ranged from 5.0 - 6.0 with an average value of 5.7 (Table 2). This is quite acidic, indicating a high potential of acidic leachate generation and confirming the lack of inherent alkalinity producing minerals. Unpalatable situations would in turn affect the larger predators and compel them to move to other places once they lose their food supply. Crops cannot naturally grow well and flourish in a polluted soil. The acidic level may have adverse effect on agricultural soil. Table 3 and figure 3 shows the Enrichment Factors (EF) in the sampled soils. Enrichment factor can fully reflect the impact which was brought to the environment by human activities [31]. It is also a convenient measure of geo-chemical trend for making comparisons between an area and overtime [32]. Based on (EF), the health risk level (HRL) is inferred. When the (EF) is less than 1, it indicates that there is no contamination, HRL is zero. When it is equal to one or two, it indicates that the pollution is minimal and HRL is 1. When (EF) lies in between 2 and 5, it indicates a low level and HRL is 2. When (EF) ranges from 5 - 20, HRL is 3. This shows an intermediate level of pollution. When (EF) is 20 – 40, HRL is 4 and it indicates a high level of pollution but when (EF) is greater than 40, HRL is 5,

184 it means that pollution is intense. Average values (Table 3) indicate that the enrichment trend
 185 of the metals is in this order: Cd > Pb > Ni > Zn > Cu > Cr > Co. Cobalt and Chromium, each
 186 has HRL of 2 and this indicates a low level pollution condition. Nickel, Zinc and Copper are
 187 at intermediate levels of pollution with a HRL of 3. Cadmium and Lead show respective
 188 HRL of 5 and 3. Their pollution levels are considered high.
 189



190

191 Figure 3. Spatial distribution of the enrichment factor of some heavy metals: (a) Cadmium(Cd)
 192 (b) Copper (Cu) (c) Nickel(Ni) (d) Cobalt (Co)

193

3.1 POLLUTION INDEX AND ENRICHMENT FACTOR

194 Pollution is any process whether natural or man-made which leads to harmful or
 195 objectionable increase in the amount of any factor in the environment [27]. Pollution Index
 196 (PI) and Enrichment Factor (EF) were employed to assess the pollution of individual metals
 197 in the top soils. The enrichment factor estimates presented in table 3 was computed using the
 198 method prescribed by [31] as shown in equation 1 below:

$$EF = \frac{C_n(\text{sample})/C_{ref}(\text{sample})}{B_n(\text{Background})/B_{ref}(\text{Background})} \dots \dots \dots (1)$$

199

200 Where C_n (Sample) is the concentration of the given element in the examined environment,
 201 and C_{ref} (Sample) is the concentration of the element in the reference sample. B_n
 202 (Background) is the concentration of the element in the background samples in the
 203 environment. B_{ref} (Background) is the concentration of the element in the reference
 204 background sample environment. Enrichment factor (EF) represents the level of the element
 205 in considered space [31]. Pollution Index (Pi) was also used in this study to assess the

contamination of individual metals. Similarly, the pollution index (PI) presented in table 4 and presented in figure 4 was calculated using [31] as shown in equation 2 below:

$$PI = C_i/S_i \dots \dots \dots (2)$$

Where P_i = Pollution Index, C_i = Concentration of heavy metal in the medium and S_i = relevant standard for this metal [26]. The tolerable limits set by various established standards of non-polluted soils from [22-26] as shown in table 5 were used as the assessment criteria.




Figure 4. Spatial distribution of the pollution index of some heavy metals: (a) Chromium (Cd) (b) Copper (Cu) (c) Lead (Pb) (d) Zinc (Zn)

The pollution index (P_i) was introduced as part of assessing pollution levels by considering the joint effects of all polluting metals in soils [33]. Pollution index was extensively used for soil pollution studies in China [34-36] and proved to be simple and effective because it related to a certain quality criterion. The Pollution Index data (Table 4) highlights the pollution status of the soils in the area. Pollution index (P_i) was also adopted here to assess the contamination of individual metals. The pollution index was calculated based on concentration of seven metals namely: Cd, Cr, Cu, Ni, Pb, Co and Zn which are considered significant in the pollution assessment of the area. The pollution index (P_i) was calculated by averaging the ratios of the concentration of the heavy metals to the hazard criteria (tolerable level) using the formula as shown $P_i = C_i/S_i$ [37], where P_i = Pollution index; C_i = Average concentration of heavy metal in the medium and S_i = relevant standard (tolerable level) for this metal [26]. Based on Pollution Index (P_i), soil contamination levels can be classified into unpolluted ($P_i < 1$); slightly polluted ($1 \leq P_i < 2$); moderately ($2 \leq P_i < 3$) and heavily polluted ($P_i \geq 3$). The pollution index of Cadmium in an unpolluted soil from literature should be less than one. Table 4 shows Cadmium pollution index ranged from 0.00 - 76.50 with an average value of 27.73 mg/kg. This implies a high pollution level according to the criteria of [26].

Lead varied from 0.07 mg/kg to 0.37 mg/kg without local enrichment at any of the localities. The average Lead value 0.14 mg/kg is below the toxic limit. All the remaining five metals analyzed were either below detection levels or are very low in pollution index criteria. They ranged from (0.00 – 0.43) mg/kg with averages Cr (0.01) mg/kg, Cu (0.03) mg/kg, Ni (0.09) mg/kg, Co (0.01) mg/kg and Zn (0.05) mg/kg. The pollution trend is in the order: Cd > Pb > Ni > Zn > Cu > Cr > Co. None of the average values exceeded one (Table 4). This, according to the classification of [26], implies no pollution by these metals. Their health risk level is zero and thus does not pose any health risk to humans though may have some implications to soil and associated plants.

4.0 SUMMARY AND CONCLUSION

Heavy metal analysis of fourteen top soil samples collected from old Awka-Enugu road were carried out using atomic absorption spectrophotometer (AAS). Result of the analysis revealed that cadmium concentration in the study area ranged from <0.37 to 6.12 mg/kg. High cadmium concentration in the study area may be associated with natural and anthropogenic sources. Similarly, chromium concentrations ranged from < 1.12 to 5.12 mg/kg with an average value of 1.99 mg/kg while copper concentrations varied from 0.12 to 3.75 mg/kg, with mean value of 1.44 mg/kg. Nickel concentrations across the study area ranged from 0.87 – 11.87 mg/kg while cobalt ranged from 0.25 to 1.22 mg/kg. Lead concentrations across the study area ranged from 0.50 to 4.75 mg/kg with the highest concentrations revealed at sites 7 and 14 where automobile mechanic workshops are located. Finally, zinc concentrations ranged from <1.87 to 7.25 mg/kg with an average value of 4.41 mg/kg.

In conclusion, cadmium concentration was generally high in the examined area compared to the typical range found in unpolluted agricultural soils. The study area has been highly contaminated by cadmium and lead but moderately polluted with respect to nickel, chromium and zinc. Copper and cobalt are deficient and pose no risk to the environment. Enrichment Factor (EF) for the elements in the different sites shows significant variability. It is suggested that the (EF) variability is due to anthropogenic activities in these areas. Anthropogenic sources include agricultural (fertilizers, animal manures, pesticides), carbides used for welding works as well as micro-electronics, sewage sludge and scrap disposals [38]. Traffic pollution may be a likely source for heavy metal contamination. Another source is bush burning which according to [39] releases the heavy metals contained in leaves and tissues in plants into soils.

5.0 REFERENCES

- [1] Hopke, P. K., Lamb, R.E and Natusch, F. S.,1980. Multi-elemental characterization of urban road way dust. *Environmental Science and Technology* 14;pp 164-172.
- [2] Beckwith, P. R., J.B.Ellis, and D.M. Revitt., 1986. Heavy metal and magnetic relationships for urban source sediments, *Physics of the Earth and Planetary Interiors*, 42 :67-75, 1986.
- [3] Xie, S., J. Dearing., J.Bloemendal., and J.Boyle.,1999. Association between the organic matter content and magnetic properties in street dust. *Liverpool, UK, The Science of the Total Environment*, 241: 205-214.
- [4] Lecoanet, H., Leveque, F and Ambrosi, J .P. (2003). Combination of magnetic

parameters: an efficient way to discriminate soil-contamination sources (south France)
Environmental Pollution 122;pp.229-234.

[5] Robertson, D .J, Taylor, K .G and Hoon, S .R. (2003) Geochemical and mineral magnetic
characterization of urban sediment particulates. Manchester, UK *Applied Geochemistry*
18: 269-282.

[6] Charlesworth, S., Everett, M., McCarthy, R., Ordonez, A., & De Miguel, E. (2003). A
comparative study of heavy metal concentration and distribution in deposited street
dusts in a large and a small urban area: Birmingham and Coventry, West Midlands, UK.
Environment International,29:563–573,2003; doi:10.1016/S0160-4120(03)00015-1.

[7] Virad, B., Pihan, F., Promeyrat, S., Pihan, J.C., 2004. Integrated assessment of heavy
metal (Pb, Zn, Cd) highway pollution: bioaccumulation in soil, Graminaceae and land
snails. *Chemosphere* 55, 1349– 1359.

[8] Robertson, D. and Taylor, K. J. , 2007. Temporal Variability of Metal Contamination in
Urban Road-deposited Sediment in Manchester, UK: Implications for Urban Pollution
Monitoring; Water, air and soil pollution, Volume 186, Issue 1, pp 209–
220;doi:10.1007/s11270-007-9478-x

[9] Li, X. D., Poon, C. S., & Liu, P. S. (2001). Heavy metal contamination of urban soils
and street dusts in Hong Kong. *Applied Geochemistry*,16,pp.1361–1368.
doi:10.1016/S0883-2927(01)00045-2.

[10] Culbard, E.B., Thornton I., Watt J., Wheatley M. and Moorcroft S., Thompson M.
(1988) Metal contamination in British urban dusts and soils; *J Environ Qual.*, 17, 226–
234

[11] Iwegbue, C.M.A., N.O. Isirimah., C. Igwe and E.S. William., 2006.Characteristic levels
of heavy metals in soil profiles of automobile mechanic waste dumps in Nigeria.
Environmentalist, 26: 131 – 137.

[12] Garcia,R and Millan,E.,1998. Assessment of Cd, Pb and Zn Contamination in roadside
Soils and grasses from Gipnzkoa (Spain). *Chemosphere* 37(8):1615 – 1625.

[13] Ward, N., Brocks, R. R., Roberts, E and Boswell, C. R.,1977. Heavy metal pollution
from automotive emissions and its effects on roadside soils and pasture species in
New Zealand. *Environ, Sci. Technol.*, 11:917 – 920.

[14] Turer, D. G and Maynard, B. J.,2003. Heavy metal contamination in highway Soils.
Comparisons of Corpus Christi. Texas Cincinnati, Ohio shows organic matter is key to
mobility. *Clean Technol Environ. Policy* 4(4):235 - 245.

[15] Nabuloa, G., Oryem-Origa, H and Diamond, M.,2006. Assessment of Pb, Cd and Zn
contamination of roadside Soil, surface films and vegetables in Kampala City,
Uganda. *Environ. Res* 102 (2006) 42 – 52.

- 327 [16] Nwajide, C. S and Reijers, J.A. ,1996. Selected chapters on Geology, App.D, Excursion
328 guide (1996) pp160 – 165.
329
- 330 [17] Cliff, D and P.M. Theodorakos.,1993. Rock Samples Preparations, USGS Open-File
331 Report 02-0223-A1.
332
- 333 [18] Taylor, C. D and Theodorakos, P. M., 2002. Rock Samples Preparations. Analytical
334 Methods for Chemical Analysis on Geologic Materials, Geological Survey Open- File
335 Report 0223, [Online] <http://pub.usgs.gov/of/2002/ofr>
336
- 337 [19] Crock., J. G., Lichte, F.E and Briggs, P.H., 1983. Determination of elements in
338 National Bureau of Standards geologic reference materials SRM 278 Obsidian and
339 SRM 688 Basalt by inductivity coupled plasma-atomic emission spectroscopy: Geo-
340 standards Newsletter, Vol. 7 (1983) 335 – 340.
341
- 342 [20] Briggs, P. H., 1979. Forty elements by inductivity Coupled-plasma atomic emission
343 spectrometry in Arbogast, B .E., (ed). Analytical methods manual for the Mineral
344 Resources Program U.S Geological Survey Open-File Report 96-525 (1996) 77-94.
345
- 346 [21] H.J. M Bowen, Environmental Chemistry of the Elements. Academic Press. New York.
347 NY.1979
348
- 349 [22] Shacklette, H. T and Boerngen, J.G.,1984. Element Concentrations in Soils and other
350 surface contaminous materials of the United States: USGS Professional paper 1270.
351 U.S Government printing office. Journal of Environ. Sci, 22 (3):113 – 119.
352
- 353 [23] Kabata-Pendias, A., 1995. Agricultural Problems related to Excessive Trace Metals
354 Content of Soil. In Salomons, W; Forstner, U and Made (eds) Heavy Metals (Problems
355 and Solutions). Springer-Verlag,Berlin; pp.3 – 18.
356
- 357 [24] Krauskopf, K. B.,1979. Introduction to Geochemistry, 2nd ed, McGraw Hill International
358 Series,Tokyo; 617pp.
359
- 360 [25] R. L. Rudnick and S. Gao, Composition of the continental crust, Treatise on
361 Geochemistry Vol. 3 (2003) 1 – 64.
362
- 363 [26] Ume, J. I., 2002. Environmental Pollution Control in Conceptual Analysis of process
364 systems. Published by New Generation Ventures Ltd, Nnamdi Azikiwe Stadium
365 Shopping Mall, Enugu-Nigeria;p.102.
366
- 367 [27] Finkelman, R. B and Zheng, B., 1999. Health Impacts of Domestic Coal use in China.
368 National Academy of Science proceedings, Vol. 96; p.3427 – 3431.
369
- 370 [28] Lindsay, W. L.,1979. Chemical equilibria in soils. John Wiley and Sons New York, NY.
371
- 372 [29] Fuller, W. H.,1977. Movement of selected metals, asbestos and cyanide in soil:
373 Application to waste disposal problem. EPA – 600/2-77-020.Solid and hazard waste
374 research division, U.S. Environmental protection agency, Cincinnati, OH.
375
- 376 [30] Liao, Guo-Li and Chao, W.C.,2006. Heavy metal Pollution and Control in

376 Lindsay. Transactions Nonferrous Metals Society of China 18 ; pp 207 – 211.
377
378 [31] Kumar, S. P and Edward, J. K.,2009. Assessment of metal concentration in the sediment
379 Cores of Manakudy estuary South West Coast of India; vol.38 (3); pp.235 – 348.
380
381 [32] Lee, J., Chon, H and Kim, K.,1988. Migration and Dispersion of trace Elements in the
382 rock-soil-plant system in areas underlain by black shales. Journal of Geochem. Explor.
383 vol. 61; pp.1 -78.
384 [33] Chem, T. B., Zheng, Y. M and Tain, O. Z.,2005. Assessment of heavy metal pollution in
385 the surface soils of urban parks in Beijing, China. Chemoshpere,605;pp.42 – 551.
386
387 [34] Fang, X., Tian, D. L and Xie, R. X.,2006. Soil physical and chemical properties of the
388 waste land in Xiangtan manganese mine. Atca Ecol. Sin 26 (5); pp.494 – 501.
389
390 [35] Zheng, G. Z., Yue, L. P., Li, Z. P and Chen, C.,2006. Assessment of heavy metals
391 pollution of agricultural soil in Guanzhung District. Journal Geograph. Sci. 16 (1)
392 ;pp.105 –113.
393
394 [36] Liao, Guo-Li and W. U. Chao.,2004. Application of Fuzzy mathematics approach on
395 Total Mine Environment. Journal of Environmental Science Trends, Vol.3;pp15 – 17.
396
397 [37] Mattigod, S. V and Page, A. L.,1983. Assessment of Metal Pollution in Soils in
398 Thornton, I (ed) Applied Environmental Geochemistry, Academic Press; pp.355 – 391.
399
400 [38] Ezech, H. N., Anike, O. L and Egboka, B. C. E., 2008. Evaluation of heavy metals
401 pollution of soils around the derelict Enyigba mines and their sources. International
402 Journal of Applied Sciences ISSN 0973 – 6077; vol. 3(1);pp. 1 – 10.
403