

Original Research Article

ASSESSMENT OF HEAVY METAL POLLUTION LEVEL OF AN AUTOMOBILE MECHANIC WORKSHOP IN NORTH CENTRAL REGION OF NIGERIA.

ABSTRACT

Following the reports in the literatures of the harmful components of spent motor oil, this study was geared towards determining the concentration of some heavy metals: Fe, Cd, Cu, Cr, Ni, Zn and Pb, from the soil of a mechanic workshop located in lorry (tipper) garage opposite AY filling station at Gwagwalada Area Council of the Federal Capital Territory Abuja and levels pollution of this site using pollution indices. The determined heavy metals were compared with the guideline stated by Nigerian regulatory agency, international standards and studies from other literatures. Seven spots were sampled at 0 – 15 cm depth at the mechanic workshop while a control sample was obtained from a SHESTCO environment in Sheda, Abuja. The concentrations of the heavy metals were determined using AAS after closed system digestion of the soil samples using 2M HNO₃. Other parameter of the soil characterized include: pH in water and KCl, electro-conductivity, carbonate content and total organic matter content. Statistical data analysis of the results was carried and the single indices of the heavy metals calculated. The order of the mean concentration of heavy metal content in the sampled soil was found to be Fe > Zn > Cu > Pb > Cr > Ni > Cd while the order of the mean concentration of the heavy metals in the control soil was found to be Fe > Cr > Zn > Pb > Cu > Ni > Cd. There was an indication that Ni has the same source in the sample soil with Fe and Zn with correlations at $r = 0.69$; $p = .043$ and $r = 0.85$; $p = .008$ respectively. The findings of this study would add to the environmental database of the soil of this mechanic workshop, which in due course, will assist the management and remediation strategies for heavy metal contaminated soil.

Keywords: Soil pollution, auto mechanic workshop; heavy metals; pollution indices, Gwagwalada

1. INTRODUCTION

The natural soil environment is clean and remains enjoyable for both plants and animals in the absence of pollution. Pollution is the introduction of harmful substances into a natural environment, usually by humans [1]. These harmful substances originates from agricultural chemicals such as pesticides, fertilizer, insecticides, weedicides, fumigant; industrial waste; radio active pollutants; biological agents; pollutants from automobiles ,urban waste, treated timber, lead paints, accidental spills and leakages of chemicals and oil, [2,3]. The pollutants could hamper human health, the quality of life or the natural functioning of the ecosystem by causing instability, disorder, harm or discomfort to the physical systems or living organisms therein [4]. Improper waste disposal and detrimental soil management methods have greatly reduced soil quality, thus resulting to soil pollution. Oil spillage caused by vandalized oil pipeline, indiscriminate disposal of oil at the mechanic workshops, filling stations, loading and pumping station, accidental spills during drilling from oil wells, spillage during refueling and lubrication of trucks and trains, etc; has been a concerned issued issue over the years [5] as these incident cause serious pollution of the soil environment. In developing cities of Nigeria, such as Gwagwalada in the Federal Capital territory, Abuja, there has been an increase in population with high demand for both new and fairly used cars. The maintenance of these cars has been back up by the springing up of several auto mechanic workshops to meet up with the repairs and servicing of these cars due to constant

usage. The servicing of these cars which involves mostly the change of the spent oil in the engine to new oil for effective and efficient use, has lead to the dumping of the spent motor oil indiscriminately on the soil [6].

This spent motor oil are disposed into gutters, water drain, open vacant plots and farmlands, a common practice by motor mechanics and generator mechanics [7]. Spent motor oil contains heavy metals among other chemical pollutants [4,5, 8-9] and include: Lead, Arsenic, Zinc, Barium, Chromium, Copper, Arsenic, Calcium, Aluminium and cadmium and result from wear and tear from engine parts when the oil burns inside the engine during use as an engine lubricant [7,10]. The heavy metal from spent oil have adverse physiological effects at relatively low concentration and bio-accumulate across the food chain in the ecosystem as they are absorbed and into the living tissues without been excreted as they non-biodegradable, thus become toxic: causing serious health problem to plants and animals (man inclusive) such as cardiovascular, kidney and liver problems; anemia, tremor and consequently, resulting in death [4-5 10-13]. Spent oil in the soil results to poor aeration, immobilization of soil nutrients and lowers the pH of the soil [7], which could be an unsatisfactory and detrimental condition to soil organisms; and could reduce growth and productivity of plants.

Therefore, there is a need to check the level of these heavy metals in mechanic workshop soil so as to provide useful environmental information and data for future monitoring to avoid alarming concentrations of these heavy metals in the soil environment.

2. MATERIAL AND METHODS

2.1 Sample preparation and analysis

Soil samples were collected from seven different spots of the mechanic workshop at the 0 -15 cm. The sampling was carried out with a trowel that was washed with soap and distilled water prior to sampling. The soil samples were stored in a black polyethylene bag and labelled accordingly before taken them to the laboratory. At the laboratory, the samples were air dried for 1 week and passed through a **of** 2mm sieve. The physicochemical properties of the soil were determined as follows: total Calcium trioxocarbonate (IV) according [14] method; wet Digestion of Soil Samples for metal analysis of- Fe, Cd, Cr, Cu, Ni, Zn and Pb, as the soil from each spot was digested in duplicates using 2M HNO₃ [15-17]; pH in water and KCl was carried out using the ph meter [18]; organic matter of the soil samples were determined based on Walkey- Black method according to the procedure of Estefan *et al* [14].

The data obtained were subjected to statistical analysis. One-way ANOVA analysis was use to test the significant difference of the mean of the heavy metals. Descriptive was to reveal the minimum, maximum, mean and standard deviation of the concentrations of the heavy metals obtained after AAS analysis. Correlation analysis was used to ascertain the **the** probable common source of the heavy metal pollutants in the contaminated soil [19,20].

2.2 Pollution indices

Some indicators were used to assess and interpret the contamination status of **the** each heavy metal in the contaminated soil. These indicators include contamination factor, degree of contamination, ecological risk factor, enrichment factor, and index of geo-accumulation etc.

Contamination factor:

Contamination factor is used to express the contamination of a given toxic substance [21].

Mathematically, it is expressed as

$$C_f^i = \frac{C_r^i}{C_R^i} \quad (1)$$

Where:

C_f^i = the contamination factor of a single metal;

C_r^i = the measured concentration of the metal in the sample;

C_R^i = the background concentration of the soil according to DPR [22]

Contamination factor is defined according to four categories [23]:

Contamination values < 1 = low contamination factor,

1 ≤ Contamination factor value ≤ 3 = moderate contamination,

3 < Contamination factor value ≤ 6 = considerable contamination) and

Contamination factor value > 6 = very high contaminated)

Degree of contamination:

The sum of the contamination factors of all the elements in the sample is referred to as the degree of contamination, which is mathematically expressed as:

$$C_D = \sum_{i=1}^n C_f^i \quad (2)$$

Where:

C_D = Degree of contamination

C_f^i = Contamination factor of a single element i

n = Count of the heavy metal

According to Hakanson, the degree of contamination in soil and sediments may be termed the sum of pollution [21]. Four categories has been defined for the degree of contamination as follows; (< 8 indicates low degree of contamination), (8-16 indicates moderate degree of contamination), (16-32 indicates considerable degree of contamination) and (>32 indicates very high degree of contamination) [23]

Potential Ecological risk factor

Hakanson [21] stated that initially, potential ecological risk factor was only applicable to water pollution control but have in recent times been effectively applied to determine the extent of pollution in soils and sediments. Therefore, this factor evaluates the potential harm of a given heavy metals in the studied soil. The categories of potential ecological risk factor and Index are as shown on (table 1)

The proposal by [21] as shown in equation (3) was followed in determining the potential ecological risk index of the heavy metals studied in the contaminated soil.

$$E_f^i = T_f^i \times C_f^i \quad (3)$$

Where:

E_f^i = the potential ecological risk factor of single metal;

T_f^i = the toxicity response factor of a given metal; and

C_f^i = Contamination factor of a single element i

The toxicity response factors of metals [21] are:

Cd = 30; Cr = 2; Cu = Pb= Ni= 5; Zn = 1

The Potential Ecological risk index was calculated based on the equation (4), which is a sum of the potential ecological risk of the single heavy metal in the sample from each spot. The format of calculating degree of contamination applies to potential risk index.

$$RI = \sum_{(i=1)}^n E_f^i \quad (4)$$

Where:

E_f^i = the potential ecological risk factor of single metal;

RI = the potential ecological risk index of many metals

n = Count of the heavy metal

Table 1: Categories of E_f^i and RI [20]

Ranges of Potential Ecological risk	Categories of Potential Ecological risk	Ranges of Potential risk index	Categories of potential risk index
< 40	Low	RI < 150	Low grade
$40 \leq E_f^i < 80$	Moderate	$150 \leq RI < 300$	Moderate
$80 \leq E_f^i < 160$	Higher	$300 \leq RI < 600$	Sever
$160 \leq E_f^i < 320$	High	$600 \leq RI$	Serious
$320 \leq E_f^i$	serious		

Index of Geo-accumulation

Index geo-accumulation is used to assess the effects of heavy metal contamination on agriculture and man [24]. Therefore, it is used to determine the extent of metal pollution as proposed by Muller [25].

Mathematically, it is stated as :

$$I_{geo} = \log_2 \frac{C_D^i}{1.5C_R^i} \quad (6)$$

Where:

Where:

C_r^i = the measured concentration of the metal in the sample;

C_R^i = the background concentration of the soil according to [22]

I_{geo} = Index of geo-accumulation

1.5 is the correction factor for compensating the background data as a result of lithogenic effects. The classification of geo-accumulation index is presented in (table 2) as reported by [26]

Table 2: Classification of geo-accumulation index

I_{geo} Value	Class	Soil Quality
≤ 0	0	Uncontaminated
0 – 1	1	From Uncontaminated to moderately contaminated
1 – 2	2	Moderately contaminated
2 – 3	3	From moderately contaminated to strongly contaminated
3 – 4	4	Strongly contaminated
4 – 5	5	From strongly contaminated to extremely contaminated
> 6	6	Extremely contaminated

3. Results and discussion

3.1 Physicochemical Properties of the soil

The results of some physicochemical properties of the soil from the mechanic workshop are as presented in (Table 1). The pH of the soil was found to be slightly alkaline as the pH of the contaminated soil sample determined in water ranged from 7.33 to 7.51 with a mean value of 7.41 ± 0.06 while the pH of the soil in KCl ranged from 7.26 to 7.58 with a mean value of 7.43 ± 0.011 . It was also found that there was no significant difference in all the values of pH obtained from the various soil samples, whether in water or in KCl. The pH values obtained were in line with those reported in the literatures, [27-28]. Though other authors [4, 29], reported pH that were slightly acidic.

The Electro-conductivity of the soil was found to range from 1151 to 1182 with a mean value of 1161.86 ± 10.84 . However, it was found to be much higher than the reported by Ekeocha and Anunuso[30].

The Carbonate content ranged from 1.48 to 1.86 with a mean concentration of 1.64 ± 0.13 %, which was lower than that reported by [27].

The mean value of the total Organic matter of the soil was found to be 3.20 ± 0.25 % as it ranged from 2.73 to 3.49.

Table 3. Physicochemical properties of the experimental soil (0-15 cm)

Sample size	pH in water	pH in KCl	Electro-conductivity	Carbonate content %	Total Organic Matter (%)
7	7.39	7.4	1161	1.65	3.1978
7	7.45	7.43	1157	1.59	3.25
7	7.40	7.53	1163	1.48	3.42
7	7.51	7.58	1182	1.66	3.20
7	7.33	7.26	1151	1.74	3.11
7	7.38	7.37	1168	1.86	2.73
7	7.42	7.45	1151	1.53	3.49
Min	7.33	7.26	1151	1.48	2.73
Max	7.51	7.58	1182	1.86	3.49
Mean $\pm$ SD	7.41 $\pm$ 0.06	7.43 $\pm$ 0.011	1161.86 $\pm$ 10.84	1.64 $\pm$ 0.13	3.20 $\pm$ 0.25

3.2 Heavy metal Concentrations

The AAS analysis results of the contaminated soil indicating the concentrations of the heavy metals are shown (table 2). All the heavy metals of interest were detected, though in some spots, the concentration of Cadmium was below the detection limit. More so, the descriptive analysis of the heavy metal concentration are presented (table 3). The heavy metal concentrations of the contaminated soil were compared with the allowable concentration of some countries is as shown in table 4.

Table 4: Heavy metal concentration of the soil samples

Sampling spots	Heavy metals (mg/kg)						
	Fe	Cd	Cu	Cr	Ni	Zn	Pb
1	490.11 $\pm$ 0.01	ND	59.10 $\pm$ 0.021	10.15 $\pm$ 0.01	4.18 $\pm$ 0.05	115.44 $\pm$ 1.48	60.57 $\pm$ 1.42
2	331.56 $\pm$ 3.42	0.011 $\pm$ 0.01	55.55 $\pm$ 1.91	9.44 $\pm$ 1.25	3.36 $\pm$ 0.94	121.62 $\pm$ 2.63	19.14 $\pm$ 1.55
3	387.24 $\pm$ 2.22	ND	48.71 $\pm$ 3.16	6.98 $\pm$ 2.66	3.07 $\pm$ 0.55	124.32 $\pm$ 0.13	28.92 $\pm$ 4.17
4	461.19 $\pm$ 1.81	ND	50.42 $\pm$ 2.28	8.51 $\pm$ 1.49	4.02 $\pm$ 0.03	119.75 $\pm$ 0.87	44.68 $\pm$ 3.82
5	373.17 $\pm$ 2.59	0.002 $\pm$ 0.12	47.33 $\pm$ 4.25	9.22 $\pm$ 0.79	2.22 $\pm$ 1.47	107.92 $\pm$ 6.56	79.55 $\pm$ 4.97
6	408.57 $\pm$ 0.43	ND	41.96 $\pm$ 3.44	10.05 $\pm$ 0.02	4.00 $\pm$ 0.11	122.38 $\pm$ 1.42	51.13 $\pm$ 5.30
7	485.62 $\pm$ 4.11	ND	55.74 $\pm$ 2.22	8.88 $\pm$ 1.83	3.92 $\pm$ 0.09	120.54 $\pm$ 3.81	65.78 $\pm$ 1.66
Control	11.14 $\pm$ 0.04	0.01 $\pm$ 0.01	0.91 $\pm$ 1.28	10.43 $\pm$ 4.01	0.78 $\pm$ 0.21	5.83 $\pm$ 2.98	3.99 $\pm$ 1.18

Value = mean $\pm$ SD, n = 2.

172 Table 5: some descriptive analysis of the contaminated soil

Sample size	Metals	Range	Min	Max	Mean±SD
7	Fe	158.55	331.56	490.11	419.64±60.74
7	Cd	0.01	ND	0.011	0.01±0.01
7	Cu	17.14	41.96	59.10	51.26±5.90
7	Cr	3.17	6.98	10.15	9.03±1.07
7	Ni	1.96	2.22	4.18	3.54±0.71
7	Zn	16.40	107.92	124.32	118.85±5.55
7	Pb	60.41	19.14	79.55	49.97±21.06

173 Iron Concentration

174 Iron was detected in all the spots of the mechanic workshop. The soil Fe concentration ranged from 331.56±3.42 to
 175 490.11±0.004 mg/kg. The mean concentrations of the Iron in the contaminated soil were found to be 419.64±60.74 mg/kg.
 176 There was an extreme significant difference between the mean concentration of Fe in the sample, 419.64±60.74 mg/kg,
 177 and the control soil, 11.14±0.04 mg/kg, at $p < 0.05$. The mean concentrations of Fe was found to be extremely higher than
 178 those of Cd, Cu, Cr, Ni, Zn and Pb ($p < 0.001$). The iron concentration in this study was found to be lower than the
 179 background value stated by DPR [22]. Some authors, [31-32] reported 748-70,606 ±10114.3 mg/kg and 11,776mg/kg of
 180 Fe concentration that were much higher.

183 Cadmium Concentration

184 The concentration of cadmium was found to be below detection limits for some of the spots analyzed as the results
 185 ranged from ND to 0.011±0.01mg/kg. The mean concentration of the cadmium in the contaminated soil was found to be
 186 0.01±0.01mg/kg. There was no significant different between the Cadmium concentration of the sample (0.01±0.01mg/kg)
 187 and that of the control soil (0.01±0.01 mg/kg). However, the mean concentration of Cd was found to be significantly lower
 188 than those of Cu ($p = 0.006$), Zn ($p < 0.001$) and Pb ($p = 0.007$) and Fe ($p < 0.001$). The cadmium concentration was found
 189 to be lower than those of: DPR background value and intervention values; Dutch target; UK allowable limits and other
 190 international standards. The cadmium concentration from this contaminated soil was lower than that reported in some
 191 literatures [31, 33-34]

193 Copper Concentration

194 The concentration of copper in the contaminated soil ranged from 41.96±3.44 to 59.095±0.02 mg/kg. The mean
 195 concentration of the copper in the contaminated soil was found to be 51.26±5.90 mg/kg. The Copper concentration of the
 196 sample, 51.26±5.90 mg/kg, was found to be significantly higher than that of the control soil, 0.91±1.28 mg/kg. Though in
 197 the sample from the mechanic workshop, the concentration of Cu was found to be statistically difference from the
 198 concentrations of Fe ($p = 0.000$), Cd ($p=0.006$); Cr ($p=0.036$); Ni ($p=0.012$) Zn ($p=0.000$) but did not differ significantly with
 199 that of Pb ($p = 1.000$). The Copper concentration was lower than the intervention value of DPR, EEC guideline, UK
 200 allowable limits Austria allowable limits and France but higher than the background value of DPR Dutch target value,
 201 Germany threshold values, Canadian criteria and Sweden allowable limits. The Cu concentration reported by [31,33-34]
 202 were all lower that the concenetratation found in this study..

204 Chromium Concentration

205 Chromium was present in all the samples of the contaminated soil analyzed, with a ranging concentration of 6.98±2.66 to
 206 10.15±0.006 mg/kg. The mean concentration of the chromium in the contaminated soil was found to be 9.03±1.07 mg/kg.
 207 There was no significant difference between the concentration of Chromium in the sample soil, 9.03±1.07 mg/kg and that
 208 of the control soil, 10.43±4.01 mg/kg. More so, the mean concentration of Cr differed significantly with the concentrations
 209 of Fe ($p < 0.001$); Cu ($p = 0.036$); Zn ($p < 0.001$) and Pb ($p = 0.046$). The concentration of Chromium was found to be
 210 lower than the allowable concentrations presented by most countries: France, Canada, Sweden, Austria, Germany, UK,
 211 Europe, Dutch and that of DRP. A Cr concentration of 12.9 mg/kg, which was higher was reported [32]

213 Nickel Concentration

214 The range of the Nickel concentration in the contaminated soil was 2.22±1.47 to 4.18±0.049 mg/kg, with a mean
 215 concentration of 3.54±0.71 mg/kg. The Nickel concentration in the sample soil, 3.54±0.71 mg/kg, did not differ significantly
 216 with that of the control soil 0.78±0.21 mg/kg at $p < 0.05$. However, in the contaminated soil, the concentration of Ni was
 217 significantly lower than the concentrations of those of Fe ($p = 0.001$); Cu ($p = 0.012$); Zn ($p = 0.001$) and Pb ($p = 0.016$).

The mean concentration of Nickel was found to be lower than the concentration limits stated by DPR, Austria, Germany, UK, France, Canada, Europe, Dutch and Sweden.

Zinc Concentration

The concentration of Zinc ranged from 107.92 ± 6.56 to 124.32 ± 0.127 mg/kg, though the mean concentration was found to be 118.85 ± 5.55 mg/kg. The Zinc concentration of the sample soil, 118.85 ± 5.55 mg/kg was significantly higher than that found in the control soil, 5.83 ± 2.98 mg/kg. An extreme significant difference was observed between the mean concentration of Zn and all the other heavy metals at $p < 0.001$. The concentration of Zinc in the contaminated soil was found to be **with in** the allowable concentration  some countries as shown on table 4 but exceeded that of Canada. The Zinc concentration reported by Abdullah **et al**  was higher than that in this study while Karim **et al** [32] reported a concentration of 123.03 mg/kg similar to 124.32 mg/kg in this study.

Lead Concentration

Lead was found present in all the samples obtained from the different spots of the mechanic workshop and was found to range from 19.14 ± 1.55 to 79.55 ± 4.97 mg/kg. The mean concentration of Pb, 49.97 ± 21.06 mg/kg, in the contaminated soil was found to be high 3.99 ± 1.18 mg/kg for Pb. The concentration of Pb in the sample soil, 49.97 ± 21.06 mg/kg, was found to differ significantly from that of the control soil, 3.99 ± 1.18 mg/kg, at $p < 0.05$. However, the mean concentration of Pb was found to be statistically different from the mean concentration of Fe ($p < 0.001$); Cd ($p = 0.007$); Cr ($p = 0.046$); Ni ($p = 0.016$) and Zn ($p < 0.001$). Though the concentration of Lead was lower than permissible limits stated by DPR, France, Austria, Germany, UK EEC and Dutch, it surpassed the allowable concentrations in Canada and Sweden. Karim **et al** [32] reported higher concentration of Pb.

The order of the mean concentration of heavy metal content in the sampled soil was found to be $Fe > Zn > Cu > Pb > Cr > Ni > Cd$ whereas, the concentration profiles of the metals in the soil of Iwegbue **et al** [41] follow the order $Fe > Pb > > Cr > Zn > Ni > Cu > Cd$. In both study, Iron had the highest concentration while Cd had the least concentration. . More so, the order of the mean concentration of the heavy metals in the control soil was found to be $Fe > Cr > Zn > Pb > Cu > Ni > Cd$. The concentration order of Fe, Pb, Ni and Cd in the contaminated site and control of this study were found to be the same.

Iwegbue **et al** [29] reported lower concentrations of heavy metals from their control site as follows: 0.93 mg/kg for Cr; 1.8 mg/kg for Cu; 4.8 mg/kg Zn; 2.6 mg/kg for Pb; 1893 mg/kg for Fe, 4.1 mg/kg for Ni as Cd found was below their detection limit. While Ilemobayo and Kolade [6] reported heavy metal concentrations (Fe, 611.24mg/kg; Cd 0.00mg/kg; Cu 124 mg/kg; Cr 115 mg/kg; Ni 43.57mg/kg; Zn 693 mg/kg Pb 218 mg/kg) which were higher than the concentration in this study. The control concentration of this study were lower than that reported by Ozulu **et al** [28], with Pb, Fe, Cu, Zn, and Cd concentration as 0.95 mg/kg, 999.88 mg/kg; 13.11 mg/kg; 23.11 mg/kg and 2.06 mg/kg respectively. Higher concentrations of Pb, Fe, Cu, Ni were reported from the control sites of Iwegbue **et al** [41]: 0.93 mg/kg for Cr, 1.8 mg kg⁻¹ for Cu, 4.8 mg kg⁻¹ for Zn, 2.6 mg kg⁻¹ for Pb, 1893 mg kg⁻¹ for Fe, and 4.1 mg kg⁻¹ for Ni, while Cd = BD: while Cr, Ni and Cd concentrations reported herein were higher than the concentration reported by Iwegbue **et al** [41].

282 Table 6: comparisons with some international guidelines

	Heavy metals (mg/kg)						
	Fe	Cd	Cu	Cr	Ni	Zn	Pb
Present study	419.64	0.01	51.26	9.03	3.54	118.85	49.97
Background value, DPR	4700	0.8	36	100	35	140	85
Intervention value DPR		17	190	380	210	720	530
Dutch target value		0.8	36	100	35	140	85
EEC guideline values (1993)		1-3	140-300			150 -300	50- 300
UK allowable limits		3	135	400	75		300
Germany allowable limits		1	40	60	50	150	70
Germany threshold values		0.4 -1.5	20 - 60	30 -100	15 -70	60-200	40-100
Austria allowable limits		1-2	60 -100	100	50 -70		100
Sweden allowable limits		0.4	40	60	30		40
Canadian Criteria (CCME)		0.5	30	20	20	60	25
France allowable limits		2	100	150	50		100

283 ECDGE [34], DPR [22], CCME [35], Iwegbue et al [29].

285 3.3 Data analysis

286 Correlation analysis of soil Heavy metal and other parameters of the contaminated soil are presented in (table 7). The correlation analysis is used to explore the inter-metal relationships of the heavy metals and ascertain the probable common sources of the heavy metal pollutants in the contaminated soil [19-20]. The linear relationship between two variables are ascertained on a scale of between -1 (negative or inverse relationship) to +1 (positive or sympathetic relationship). IBM SPSS Statistics 21 software package was used to analyze the correlation analysis between the heavy metal concentrations, pH in water and in KCl, Carbonate content, and Total organic matter content of the contaminated soil.

294 The mean concentration of Fe was found to be significantly correlated with Cd ($r = -.711$; $p = .037$) and Ni ($r = 0.69$; $p = .043$). More so, Zn and Ni were found to be significantly correlated at $r = 0.85$; $p = .008$. There is an indication that Ni has the same source in the sample soil with Fe and Zn. Other parameters that correlate significantly to each other include: pH in KCl and pH in water ($r = 0.84$; $p = .009$), Carbonate content and Cr ($r = 0.68$, $p = .048$), Total organic matter and Carbonate content ($r = - 0.96$; $p < 0.001$).

299 In addition, significant correlation were found between Fe, Cd, Ni, Zn,Cr, Carbonate, pH water, pHKCl and organic matter. However, Cu and Pb were found not to have any significant correlation with any of the other parameters at all.

302 Table 7: Correlation coefficient matrix for soil parameters

	Fe	Cd	Cu	Cr	Ni	Zn	Pb	pHH ₂ O	pH	Cond	Carbonate	O.M
									KCl	.		
Fe	1											
Cd	-.711*	1										
Cu	.373	.271	1									
Cr	.146	.183	.138	1								
Ni	.690*	-.265	.306	.332	1							
Zn	.600	-.435	.194	-.004	.848**	1						
Pb	.467	-.540	-.063	.363	-.144	-.274	1					
pHH ₂ O	.205	.186	.295	-.230	.531	.317	-.557	1				
pHKCl	.281	-.139	.188	-.607	.472	.551	-.602	.840**	1			

Conductivity	.225	-.282	-.288	-.142	.490	.430	-.347	.648	.647	1		
Carbonate	-.076	-.125	-.597	.676*	.045	-.171	.390	-.324	-.563	.230	1	
O. M	.188	.062	.649	-.610	-.097	.029	-.155	.286	.459	-.315	-.959**	1

*. Correlation is significant at the 0.05 level (1-tailed).

**. Correlation is significant at the 0.01 level (1-tailed).

O.M= Organic matter.

The calculated contamination factors, which represent the impact of the individual heavy metal element on the soil from the different spots analyzed are presented on (table 8). The highest contamination factor of Fe, 0.10, was from spot 1 and 7 while the mean contamination factor was 0.09. The maximum and mean values of the contamination factor of Cd were found to be 0.01 each, whereas, those of Cu were: 1.64 and 1.42 respectively. The maximum and mean contamination factor of Cr, Ni, Zn and Pb were: 0.1 and 0.09, 0.12 and 0.1, 0.89 and 0.85, 0.94 and 0.59 respectively. The categories of the contamination factor of the single heavy metals found to be low contamination for Fe, Cd, Cr, Ni, Zn and Pb while Copper was of moderate contamination. The moderate contamination of copper in the mechanic workshop could have originated from some mechanical parts of the cars that are plated with copper, a content of the gasket sealant and anti-seize compound, a component of bronze which is used in making crankshaft and camshaft bearings [36-37]. Stratson [37] also mentioned that other sources of copper could be from: Leaching from oil coolers, leaks from cooling systems, external contaminates and from excessive wear. Copper is reported to have originated from additives, such as anti-wear and corrosion inhibitors additives, that were added to engine oil used in the crankshaft of the car [38-40]. From (table 8), the degrees of contamination of the heavy metals in all the soils of the spots in the mechanic workshop were low. This could imply that the mechanic workshop

Table 8 . Values of contamination factor and degree of contamination of contaminated soil

Sampling Spots	Contamination factor							Degree of contamination
	Fe	Cd	Cu	Cr	Ni	Zn	Pb	C _d
1	0.10	0.00	1.64	0.10	0.12	0.89	0.71	3.56
2	0.07	0.01	1.54	0.09	0.10	0.82	0.23	2.86
3	0.08	0.00	1.35	0.07	0.09	0.87	0.34	2.80
4	0.10	0.00	1.40	0.09	0.11	0.86	0.53	3.09
5	0.08	0.00	1.31	0.09	0.06	0.77	0.94	3.25
6	0.09	0.00	1.17	0.10	0.11	0.87	0.60	2.94
7	0.10	0.00	1.55	0.09	0.11	0.86	0.77	3.48
Min	0.07	0	1.17	0.07	0.06	0.77	0.23	2.37
Max	0.1	0.01	1.64	0.1	0.12	0.89	0.94	3.80
Mean	0.09	0.01	1.42	0.09	0.1	0.85	0.59	3.15
C _f category	Low cont.	low cont.	Moderate cont.	low cont.	low cont.	low cont.	low cont.	

The values of the potential ecological risk factor of the different in the mechanical workshop under study, ranged from: 0.00 to 0.41 for Cd; 5.83 to 8.21 for Cu; 0.14 to 0.20 for Cr; 0.32 to 0.60 for Ni; 0.77 to 0.89 for Zn; and 1.13 to 4.68 for Pb. The mean values of the potential ecological risk factor of the heavy metals are: 0.10, 7.10, 0.18, 0.50, 0.84, and 2.93 for Cd, Cu, Cr, Ni, Zn and Pb respectively. According to the categories of the potential ecological risk factor by Hu [et al](#) (2017) present on (table 1), the potential ecological risk factor of each heavy metal on (table 1), all the heavy metals studied were found to have low potential ecological risk factor.

The potential index of the heavy metals in the soil from all the studied spot were found to be of low grade as shown in (table 9). Therefore, the heavy metals have not caused any harm to the mechanic workshop under study.

Sampling spot	E_f^i						RI
	Cd	Cu	Cr	Ni	Zn	Pb	
Spot 1	0	8.21	0.2	0.6	0.89	3.56	13.46
Spot 2	0.41	7.72	0.19	0.48	0.82	1.13	10.75
Sot 3	0	6.77	0.14	0.44	0.87	1.7	9.92
Spot 4	0	7	0.17	0.57	0.86	2.63	11.23
Spot 5	0.08	6.57	0.18	0.32	0.77	4.68	12.6
Spot 6	0	5.83	0.2	0.57	0.87	3.01	10.48
Spot 7	0	7.74	0.18	0.56	0.86	3.87	13.21
Min	0	5.83	0.14	0.32	0.77	1.13	9.92
Max	0.41	8.21	0.2	0.6	0.89	4.68	13.46
Mean	0.10	7.10	0.18	0.50	0.84	2.93	11.67
E_f^i	Low	Low	Low	Low	Low	Low	

The minimum, maximum and mean values of the I_{geo} of the metals in each spot analyzed are: for Fe, -4.41, -3.85 and -4.07; Cd, ND, -6.77 and -8.81; Cu, -0.36, 0.13, -0.08; Cr, -4.42, -3.88 and -4.05; Ni -4.56, -3.65 and -3.89; Zn, -0.96, -0.75 and -0.82; and Pb -2.74, -0.68 and -1.35 respectively. The minimum values of the all the I_{geo} of the metals were found to be negative except for Cadmium that the metal concentration was below detection limit. The maximum I_{geo} of the metals under consideration were as well negative though that of copper concentration was positive, 0.13. The mean I_{geo} values for the various metals in the spots were all negative. Following the classification of the I_{geo} on (table 2), there is an indication that the spots analyzed are uncontaminated as revealed on (table 10).

Geo- accumulation index of the heavy metals of the contaminated soil

[illegible]

352
353
354
355

Table 11.

Levels of the pollution indices studied							
Pollution indices	Spot 1	Spot 2	Spot 3	Spot 4	Spot 5	Spot 6	Spot 7
C_d	Low	Low	Low	Low	Low	low	low
RI	Low grade	Low grade	Low grade	Low grade	Low grade	Low grade	Low grade

C_d = degree of contamination
RI = Potential risk index

4. CONCLUSION

The heavy metals (Fe, Cd, Cu, Cr, Ni, Zn and Pb) determined in the soil from different spots of the mechanic workshop were compared with the heavy metals from the control soil samples and some international guideline. The order of the mean concentration of heavy metal content in the contaminated soil was found to be Fe > Zn > Cu > Pb > Cr > Ni > Cd while the order of the mean concentration of the heavy metals in the control soil was found to be Fe > Cr > Zn > Pb > Cu > Ni > Cd. The concentration order of Fe, Pb, Ni and Cd in the contaminated site and control of this study were found to be the same. The results showed that the concentrations of the heavy metal in the soil samples were below the intervention concentration stated by DPR. Soil pollution assessment using C_d , RI and I_{geo} indices confirmed the soil in the mechanic workshop is uncontaminated. Thus, this particular mechanic workshop can be considered uncontaminated.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Gammon K. Pollution Facts and Types Of Pollution. (2012). Assessed April 2016. Available: <http://www.livescience.com/22728-pollution-facts.html>
- Cornell Waste Management Institute (CWMI). Sources and impact of contaminants in soil. ND. Assessed September 2017: <http://cwmi.css.cornell.edu/sourcesandimpacts.pdf>
- Petruzzelli G, Gorini F, Pezzarossa B, Pedron F. The fate of pollutants in soil, CNR, Institute of the Ecosystem Studies (ISE), Pisa (Italy), 2010. Assessed on September 2017: http://dta.cnr.it/publications/other_publications/PIAS/PIAS-C1.pdf
- Adelakan BA and Adegunde KD "Heavy metal contamination of soil and underground water at auto mechanic villages in Ibadan, Nigeria" International Journal of Physical Sciences, 2011; 6(5):1045-1058
- Abdulsalam S, Adefila SS, Bugaje IM, Ibrahim S. Bioremediation of Soil Contaminated With Used Motor Oil in a Closed System. *Journal of Bioremediation and biodegradation*, 2012; 3(12): 172-179
- Ilemobayo O, Kolade I. Profile of heavy metals from Automobile workshop in Akure, Nigeria. *Journal of Environmental Science and Technology*, 2008; 1 (1): 19- 26, 2008.
- Achuba FI. and Peretiemo-Clarke BO. Effect of spent engine oil on soil catalase and dehydrogenase activities. *International Agrophysics* 2008 22: 1-4.
- Pam AA, Sha'Ato R, Offem JO. Evaluation of heavy metals in soils around auto mechanic workshop clusters in Gboko and Makurdi, Central Nigeria. *Journal of Environmental Chemistry and Ecotoxicology*, 2013; 5(11): 298-306,

9. Usman AA, Odoma, AN, Ozulu GU. Heavy Metals Concentration in Ground Water caused by Automobile Workshop Activities and its Health Implication on the Inhabitants of Makurdi Metropolis, Nigeria. *International Journal of Health and Medical Information*, 2013; 2(2): 36-43
10. Dike BU, Okoro BC, Nwakwasi, NN, Agbo KC. Remediation of Used Motor Engine Oil Contaminated Soil: A Soil Washing Treatment. *Journal of Civil & Environmental Engineering*, 2013; 3 (1): 129-131.
11. Idugboe SO, Tawari-Fufeyin P, Midonu AA. Soil pollution in two auto-mechanic villages in Benin City, Nigeria. *Journal Of Environmental Science, Toxicology And Food Technology* 2014; 8 (1): 09-14
12. Khan MN, Wasim AA, Sarwar A, Rasheed MF. Assessment of heavy metal toxicants in the roadside soil along the N-5, National Highway, Pakistan. *Environ Monit Assess*, 2011; 182:587–595.
DOI 10.1007/s10661-011-1899-8
13. Atlas RM, Cerniglia CE Bioremediation of petroleum pollutants: diversity and environmental aspects of hydrocarbon biodegradation. *Bioscience*, 1995; 45: 332-338.
14. Estefan G, Sommer R. and Ryan J. Methods Of Soil, Plant, And Water Analysis: a Manual for the West Asia And North Africa Region (3rd Edition). Beirut: ICARDA (International Center for Agricultural Research in the Dry Areas); 2013.
15. Joel OF. and Amajuoyi CA. Physicochemical Characteristics And Microbial Quality Of And Oil Polluted Site In Gokana, River State. *Journal of Applied Science and Environmental Management*, 2009 13 (13): 99 – 103.
16. Kabiru S, Yakubu R, Lukman A, Akintola T, Alegbemi M. Heavy Metal Content in Soil in Garki Area Council of Federal Capital Territory, Abuja, Nigeria. *Biochem Anal Biochem* 2015 4: 197. doi:10.4172/2161-1009.1000197
17. Odeyemi A., Onipe O. and Adebayo O. Bacteriological And Mineral Studies Of Road Side Soil Samples In Ado-Ekiti Metropolis, Nigeria. *Journal of Microbiology, Biotechnology and Food Sciences*: 2011 1 (3): 247-266.
18. Sikora F. and Kissel D. (2010) Soil pH: Application and Principle. Assessed January 2015. Available: http://www.clemson.edu/sera6/SoilpH_Sikora%20and%20Kissel_final%20Dec%2015.doc
19. Zamani AA. Yaftian MY. and Parizanganeh A. Multivariate statistical assessment of heavy metal pollution sources of groundwater around a lead and zinc plant. *Journal of Environmental Health Sciences & Engineering* 2012; 9:29. Doi:10.1186/1735-2746-9-29
20. Hu B, Zhou J, Liu L, Meng W, Wang Z. Assessment of Heavy Metal Pollution and Potential Ecological Risk in Soils of Tianjin Sewage Irrigation Region, North China. *J Environ Anal Toxicol*, 2017; 7: 425. Doi: 10.4172/2161-0525.1000425
21. Hakanson L. An ecological risk index for aquatic pollution control. a sediment ecological approach. *Water Research*, 1980; 14: 975-1001.
22. Department of Petroleum Resources (DPR). Environmental Guidelines and Standards For The Petroleum Industries in Nigeria (revised edition) Department of Petroleum Resources, Ministry of Petroleum and Mineral Resources, Abuja-Nigeria; 2002
23. Sam RA, Ofosu FG, Atiemo SM, Aboh IJK, Gyampo O, Ahiamadjie H, Adeti JP, Arthur JK. Heavy Metal Contamination Levels in Topsoil at Selected Auto Workshops in Accra. . *International Journal of Science and Technology* 2015; 4 (5): 222-229.
24. Boateng E, Dowuona GNN, Nude PM, Foli G, Gyekye P. and Jafaru HM. Geochemical assessment of the impact of mine tailings reclamation on the quality of soils at Anglogold concession, Obuasi, Ghana. *Res. J. Environ. Earth Sci.* 2012; 4: 466-474.
25. Müller, G. Index of geoaccumulation in sediments of the Rhine River. *Geojournal*, 1969; 2: 108-18.
26. Muller G "The Heavy Metal Pollution of the Sediments of Neckars and Its Tributary: A stocktaking," *Chem. Zeit.*, Vol. 105, 1981, pp. 157-164.

27. Agbaji EB, Abechi SE, and Emmanuel SA. Assessment of Heavy Metals Level of Soil in Kakuri Industrial Area of Kaduna, Nigeria. *Journal of Scientific Research & Reports*. 2015; 4(1): 68-78.
28. Ozulu GU, Usiobaifo OB, and Usman AA. Assessing the Impact of Waste Gasoline on the Physicochemical Properties of Soils at Selected Automobile Workshops in Obiaruku, Southern Nigeria. *Universal Journal of Environmental Research and Technology*, 2013; 3, (4): 427-435
29. Iwegbue CMA, Bassey FI, Tesi GO, Nwajei GE, and Tsafe AI. Assessment of Heavy Metal Contamination in Soils around Cassava Processing Mills in Sub-Urban Areas of Delta State, Southern Nigeria. *Nigerian Journal of Basic and Applied Science*, 2013; 21(2): 96-104
30. Ekeocha CI, and Anunuso CI. Comparative Analysis Of Index Of Geoaccumulation Of Heavy Metals In Some Selected Auto Mechanic Soils In Abuja, NIGERIA. *J. Chem. Soc. Nigeria*, 2016; 41 (2): 96-102.]
31. Nwachukwu MA, Feng H, Alinnor J. Assessment of heavy metal pollution in soil and their implications within and around mechanic villages. *Int. J. Environ. Sci. Tech.*, 2010; 7 (2), 347-358.
32. Karim Z, Qureshi BA, Mumtaz M. Geochemical baseline determination and pollution assessment of heavy metals in urban soils of Karachi, Pakistan. *Ecological Indicators*, 2015; 48: 358-364
33. Abdullah MZ, Louis VC, and Abas MT. Metal Pollution and Ecological Risk Assessment of Balok River Sediment, Pahang Malaysia. *American Journal of Environmental Engineering* 2015, 5(3A): 1-7
DOI:10.5923/c.ajee.201501.01
34. European Commission Director General Environment, (ECDGE). Heavy Metals and Organic Compounds from Wastes Used as Organic Fertilizers. Final Rep., July. WPA Consulting Engineers Inc. Ref. Nr. TEND/AML/2001/07/20, 2010; 73-74. http://ec.europa.eu/environment/waste/compost/pdf/hm_finalreport.pdf.
35. Canadian Council of Ministers of the Environment (CCME). Interim Canadian environmental quality criteria for contaminated sites. Report CCEM EPC-C534, Winnipeg, Manitoba. 1991.
36. Anonymous, Metro Tech Systms: Sources of elements found in oil analysis. 2011 Metro Tech Systems Ltd. Accesses from: <http://www.metrotechsystems.ca/TInfoElement.html>
37. Stratson, Advanced Lubrication solutions. High Copper Levels in Used Oil Analysis. Assessed September 2017. Available: <http://stratson.eu/high-copper-levels-used-oil-analysis/>
38. Zajac G, Szyszlak-Bargłowicz J, Słowik T, Kuranc A, Kamińska A. Designation of Chosen Heavy Metals in Used Engine Oils Using the XRF Method. *Pol. J. Environ. Stud*, 2015; 24 (5): 2277-2283.
39. Palkendo JA, Kovach J, Betts TA. Determination of Wear Metals in Used Motor Oil by Flame Atomic Absorption Spectroscopy. *J. Chem. Educ.*, 2014; 91, (4), 579.
40. Ayoola SO, Bassey BO, Alimba CG, Ajani EK. Textile effluent induced genotoxic effects and oxidative stress in *Clarias gariepinus*. *Pak J Biol Sci*. 2012;15 (17):804-812.
41. Iwegbue CMA, Nwajei GE, Ogala JE, Overah CL. Determination of trace metal concentrations in soil profiles of municipal waste dumps in Nigeria. *Environ Geochem Health*, 2010; 32:415–430. DOI 10.1007/s10653-010-9285-y