

Original Research Article

Enumeration of total heterotrophic and petroleum degrading bacteria counts in water and sediments from Diobu Creek, Port Harcourt, Nigeria.

Introduction: Bacteria play major role in biochemical cycling of nitrogen, carbon, sulphur, and oxygen in aquatic environments. Hydrocarbons are released naturally from oil seeps and incidental discharges which represents significant source of pollution. Estuarine systems are particularly susceptible to anthropogenic hydrocarbon contamination. Although oil contaminants are weathered by photo-oxidation and evaporation but complete degradation is dependent on the metabolic activities of the microbial population inherent to the area.

Aim: The aim of this work is to determine the total heterotrophic bacterial counts (THBCs) and petroleum degrading bacteria counts (PDBC) in water and sediment from Diobu creek.

Methodology: The THB and PDB in water and sediments from a polluted creek (Diobu Creek) were determined by serial dilution and plating on nutrient agar and petroleum degrading bacteria agar and the isolated bacteria were identified.

Results: The THBCs in water were from 6.3×10^3 CFU/mL and 6.33×10^3 CFU/mL, the highest THBCs were recorded in June (wet season). The THBCs in sediments was from 1.7×10^6 CFU/g to 1.85×10^6 CFU/g. The highest THBCs were recorded in the month of June. The PDBC in water were from 0.2×10^3 CFU/mL and 3.9×10^3 CFU/mL, whereas the PDBC in sediments ranged from 3.4×10^6 to 9.5×10^6 CFU/g, high counts were obtained if sampling were carried out in rain or after rain. The bacteria isolated were *Bacillus sp.*, *Pseudomonas sp.*, *Corynebacterium sp.*, *Acinetobacter sp.*, *Alkaligenes sp.*, *Escherichia coli*, *Micrococcus*, *Klebsiella sp.* and *Flavobacterium sp.*

Conclusions: The increased counts of PDBs in aquatic environments might be stimulated by the presence of pollutant hydrocarbons or chemicals discharged into creek which were degradable by bacteria. The activity of PDBs in detoxifying polluted environments is the most eco-friendly.

Keywords: Enumeration, Total heterotrophic, Petroleum degrading, polluted, Creek.

1. INTRODUCTION

The ubiquity of bacteria confers on them the ability of inhabiting any habitat on planet earth surface, having greater biomass than any other group of organisms. This is achieved by large surface area to volume ratio, metabolically versatile and obtaining energy by oxidizing carbon, parasitism, chemoautotrophy and photo-autotrophy [1]. Bacteria play major role in biochemical cycling of nitrogen, carbon, sulphur, and oxygen in aquatic environments [2, 3]. About 470 thousand and 8.3 million tons of petroleum hydrocarbons are released into aquatic environments each year globally [4]. Half of these are released naturally from oil seeps, incidental discharges which represents significant source of pollution [3]. From 2000 to 2013, 43 large marine oil spills and 167 medium sized marine oil spills were reported [5]. Estuarine systems are particularly susceptible to anthropogenic hydrocarbon contamination.

The concentrations of poly-aromatic hydrocarbons (PAHs) were shown to exceed 100 mg/Kg sediment in UK location at Melford Haven [6].

Although oil contaminants are weathered by photo-oxidation and evaporation, complete degradation depends on the metabolic activities of the microbial population inherent to the area [7]. Petroleum hydrocarbons are the most widespread contaminants within the marine environment. Pollution by hydrocarbons in marine environments may be caused by various natural seepages and/or anthropogenic activities (discharge during tanks and/or ships transportation and/or pipeline failures) coupled with chronic pollution (ships, harbours, oil terminals, fresh water run-off, rivers and sewage systems). Of particular concern is the accumulation of low molecular weight PAHs such as naphthalene, which had been found at concentration of 2.4 mg/Kg dry weight sediment in the Tyne estuary [6], and are acutely toxic to aquatic invertebrates at concentrations as low as 8 mg/L [8]. In addition, many high molecular weight PAHs such as chrysene, which was found at concentrations of about 6.94 mg/Kg dry weight sediment at Milford Haven [6], were classed as carcinogens and can cause chronic toxic effects in fish and invertebrates [9].

Different hydrocarbon degrading bacteria have been isolated from hydrocarbons polluted environments as hydrocarbon contaminants are released into the environment by human activities [10,11]. The presence of these hydrocarbons in an environment were known to stimulate the presence of hydrocarbons degrading bacteria in the affected environment. The total heterotrophic bacterial and petroleum degrading bacteria counts obtained from oil polluted Bodo Creek in Rivers State, Nigeria; Gokana, Rivers State and Ennore Creek, India, [1, 15, 16] were similar to the bacterial counts and genera isolated by other researchers [17, 13, 12].

The hydrocarbon utilizing bacteria isolated and identified from above studies were; *Bacillus*, *Nocardia*, *Staphylococci*, *Pseudomonas*, *Flavobacterium*, *Escherichia*, *Acinetobacter*, *Enterobacter*, and *Bacillus* sp. The gram positive bacteria (rod) *Bacillus* were the most predominant, followed by the gram negative *Pseudomonas*; these bacteria have also been isolated from hydrocarbons polluted environments by other investigators [17].

About 247 strains of bacteria were isolated by other workers from hydrocarbon polluted environments. The bacteria isolated were predominantly gram-negative bacteria with prevalence of 62.34% and gram-positive bacteria 35.63%. The gram negatives isolated were four genera namely, *Pseudomonas* sp., *Vibrio* sp., *Achromobacter* sp., and *Serratia* sp. with prevalence 39.68%, 15.78%, 3.64% and 3.24% respectively; whereas the gram-positive bacteria were two genera, *Bacillus* sp and *Micrococcus* sp with prevalence of 27.94% and 7.69% respectively [1, 17]

The bacteria isolated from the brackish polluted water of Bodo Creek were, *Bacillus* sp. *Alkaligenes*, *Enterobacter*, *Cetrobacter*, *Myroides*, *Instnibacillus*, *Pseudomonas*, and *Escherichia coli* [1].

Other workers also isolated *Bacillus* sp. *Proteus*, *Pseudomonas*, *Flavobacterium*, *Corynebacterium*, *Serratia*, *Micrococcus*, *Klebsiella*, *Enterobacter* and *Azotobacter* from polluted water and sediments [11,16]. In Ennore Creek, India, *Bacillus*, *Micrococcus*, *Pseudomonas*, *Vibrio*, *Acinetobacter*, *Serratia* were isolated from hydrocarbon polluted water and sediments [13]. The aims of this study is to determine the total heterotrophic bacterial counts (THBCs) and petroleum degrading bacterial counts (PDBC) in water and sediments from Diobu Creek in Port Harcourt.

2. MATERIALS AND METHODS

2.1 Study area

Diobu creek transects Port Harcourt metropolis and it originates from behind Mile four (4) in Rumueme Port Harcourt, Rivers State, Nigeria. It empties into Amadi Creek in the old

Government Reserved Area (GRA). Numerous activities were carried out by the creek and the most prominent are the mechanic workshops and car engine wash, coupled with municipal wastes discharging into the creek. The creek is constantly exposed to petroleum hydrocarbons and other pollutants. For the purpose of this study, four sampling stations were established along the creek which were at least 200 meters apart, while station one was marked as control station upstream was about 1Km from station 2, with relatively little human activities and pollution. The coordinates for the stations were established using Gram 76GPS.

2.2 Collection of Samples

Samples of water and sediments were collected monthly during low tide at the established sampling stations. This was possible by using the tidal data published by the Nigerian Navy Hydrographic School as a guide. Samples were collected from November, 2016 to October, 2017 from each of the sampling stations.

2.3 Collection of Water and Sediments for Bacteriological Analysis

The water samples for bacteriological examinations were collected once monthly aseptically to avoid contamination. Detailed quality assurance and quality control procedures were followed for sample collection, holding and analysis (APHA, 1976). The water samples were collected at a depth of about 15 – 25 cm in opposite direction of water current into commercial sterile universal bottles.

About 20g of sediment were collected at low tide from the oxidized thin layer (1 – 5cm) surfaces of exposed mud flats along transects across intertidal zone into a wide mouth sterile glass containers sterilized by autoclaving at 121°C-121°C for 15 minutes. The water and sediment samples were immediately taken to the laboratory for bacteriological examinations.

2.4 Preparation of Media

Nutrient agar was prepared according to the manufacturer's instruction and stored in the refrigerator for total heterotrophic bacterial counts.

2.5 Media for the isolation of petroleum degrading bacteria: The isolation of petroleum degrading bacteria were carried out using engine oil/diesel 3:1 ratio 5mL, ammonium chloride 0.5g/L, dipotassium hydrogen phosphate 0.5g/L, disodium hydrogen phosphate 2.5g/L, agar 15g [18]. The media was sterilized by autoclaving at 121°C for 15minutes and dispensed into disposable petri-dishes, allowed to solidify and stored in the refrigerator for subsequent uses.

2.6 Isolation

The method of isolation used was the ten-fold dilution technique. Decimal dilution of the samples were made by adding 1mL (water) or 1g (sediment) of the sample to 9.0 mL sterile normal saline to give an initial dilution of 1:10. Subsequent serial dilutions were made by adding 1mL of the last dilution to 9.0mL of fresh sterile saline. Lastly, 0.1mL of appropriate dilution were plated out in duplicate on agar medium and evenly spread with a sterile glass rod spreader [19].

The plates for total heterotrophic counts were incubated at 30 - 35°C-35 °C for 18-24 hours on nutrient agar and the plates for petroleum degrading bacteria incubated at ambient temperature.

2.7 Identification of isolated bacteria

Series of tests including Gram's stain, chemical and biochemical tests were used for the identification of isolated bacteria such as catalase, coagulase, indole, citrate, methyl red, hydrogen sulphide production, Vourges Prausker, oxidase, and carbohydrate fermentation tests etc [18].

2.8 Statistical analysis

Statistical analysis was carried out using GraphPad Prism 5.07.

3. RESULTS

3.1 Total heterotrophic bacterial counts in water (10^3 CFU) and sediments (10^6 CFU)

The THBCs from water was high in March, April, and May which were 6.33×10^3 , 5.38×10^3 and 5.96×10^3 CFU/mL respectively. The highest count was in the month of June, while the lowest counts were recorded in November. High THBCs from sediments obtained were in the months of November, 1.76×10^6 CFU/g, January, 1.60×10^6 CFU/g, February, 1.70×10^6 CFU/g, June, 1.84×10^6 CFU/g and September, 1.65×10^6 CFU/g respectively. The highest THBC was obtained in July. The total heterotrophic bacteria counts in sediments were high compared to results obtained in water, as shown in Figures 1 and 2.




Figure 1: THBCs in water (monthly)




Figure 2: THBCs in sediments (monthly)

3.2 Petroleum degrading bacterial counts in water and sediments

The monthly petroleum degrading bacterial counts (PDBC) in water were between 0.2×10^3 CFU/mL and 3.9×10^3 CFU/mL, the counts were high in December, April and February as shown in figure 3. In sediment samples PDBC range from 3.4×10^6 CFU/g to 9.5×10^6 CFU/g, the highest PDBC were recorded in February, whereas the lowest was in March as shown in figures 4.




Figure 3: Monthly PDBC in water

Formatted: Font: Not Italic




Figure 4: Monthly PDBs in sediments

Table 1: Identification of isolated bacteria.

S/No	Colour	Surface	Edge	Translucency	Texture	Gram Rxn	Size	Shape	Motility	Methyl Red	Voges Proskaver	Oxidase	H ₂ S Production	Indole	Coagulase	Catalase	Citrate	Urease	Starch Hydrolysis	Glucose	Lactose	Sucrose	Melactose	Galactose	Manitol	Arabinose	Oxidative	Fementative	Bacteria
1	M	P	C	O	D	+	M	P	+	+	+	+	+	+	+	+	+	+	+	A	A	A	A	A	+	+	+	+	Basillus sp.
2	IG	IS	IM	IO	Mt	+	M	P	+	+	+	+	+	+	+	+	+	+	+	A	A	A	A	A	+	+	+	+	Pseudomonas sp.
3	IG	IS	IM	IO	D	+	M	P	+	+	+	+	+	+	+	+	+	+	+	A	A	A	A	A	+	+	+	+	Alcaligenes Sp.
4	W	IS	IM	IO	D	+	M	P	+	+	+	+	+	+	+	+	+	+	+	A	A	A	A	A	+	+	+	+	Corynebcterium Sp.
5	K	IS	IM	IO	Mt	+	M	P	+	+	+	+	+	+	+	+	+	+	+	A	A	A	A	A	+	+	+	+	Flavobacterium Sp.
6	K	IS	IM	IO	Mt	+	M	P	+	+	+	+	+	+	+	+	+	+	+	A	A	A	A	A	+	+	+	+	Chromobacteria Sp.
7	IG	IS	IM	IO	Mt	+	M	P	+	+	+	+	+	+	+	+	+	+	+	A	A	A	A	A	+	+	+	+	Escherichia coli
8	IO	IS	IM	IO	Mt	+	IS	IO	+	+	+	+	+	+	+	+	+	+	+	A	A	A	A	A	+	+	+	+	Micrococcus Sp.
9	IG	IS	IM	IO	Md	+	IS	IO	+	+	+	+	+	+	+	+	+	+	+	A	A	A	A	A	+	+	+	+	Vibro Sp.
10	IG	IS	IM	IO	Md	+	IS	IO	+	+	+	+	+	+	+	+	+	+	+	A	A	A	A	A	+	+	+	+	Klebsiella Sp.
11	IG	IS	IM	IO	Mt	+	IS	IO	+	+	+	+	+	+	+	+	+	+	+	A	A	A	A	A	+	+	+	+	Acenetobacter Spp

	colour	surface	edge	Transparency	Texture	Gram-Run	Size	Shape	Motility	Methyl Red	Voges-Proskauer	Oxidase	H ₂ S-Production	Indole	Coagulase	Catalase	Citrate	Urease	Starch-Hydrolysis	Glucose	Lactose	Sucrose	Mannitose	Galectose	Manitol	Arabinose	Oxidative	Fementative	
1	M	R	E	C	D	+	M	R	+	-	+	-	-	-	-	+	+	-	+	A	A	A	A	A	±	-	+	+	<i>Bacillus-spp.</i>
2	G	S	E	C	Mt	-	M	R	+	-	-	+	-	-	-	+	+	-	-	A	A	A	A	A	±	±	+	-	<i>Pseudomonas-spp.</i>
3	G	S	Sr	O	D	-	C	R	+	-	-	+	-	-	-	+	-	-	-	A	-	-	-	A	-	A	+	-	<i>Alcaligenes-Spp.</i>
4	W	S	E	O	D	+	E	R	-	-	-	-	-	-	-	+	-	+	-	A	A	A	-	A	A	A	-	-	<i>Corynebacterium-Spp.</i>
5	K	S	E	C	Mt	-	St	R	-	-	-	+	-	+	-	+	-	-	-	-	-	A	-	-	A	-	+	-	<i>Flavobacterium-Spp.</i>
6	K	D	E	O	Mt	-	St	R	+	+	-	-	+	-	-	-	+	+	-	A	-	-	A	-	-	A	+	+	<i>Chromobacteria-Spp.</i>
7	G	S	E	C	Mt	-	M	R	+	-	-	-	-	+	-	+	-	-	-	A	A	A	A	A	A	A	+	A	<i>Escherichia-coli</i>
8	O	S	E	C	Mt	+	S	C	-	-	-	-	-	-	-	+	-	-	-	A	-	-	A	A	-	-	+	-	<i>Micrococus-Spp.</i>
9	G	S	E	O	Md	-	S	C	+	-	+	+	-	-	+	+	-	-	-	A	-	-	-	-	-	-	+	-	<i>Vibro-Spp.</i>
10	G	S	E	O	Md	-	S	R	-	-	+	-	-	-	-	+	+	+	-	A	A	A	A	A	A	A	+	+	<i>Klebsiella-Spp.</i>
11	G	S	E	C	Mt	-	St	R	+	-	-	-	+	-	-	+	+	-	-	-	-	A	-	-	A	-	+	-	<i>Acinetobacter-Spp</i>

173

174 Key:

175 Cultural Characteristics

176 M = Milky, G = Green, Cr = Creamy, W = White, S = Smooth, Or = Orange, V = Violet,
177 Yellow, Dl = Dull, O = Opaque, C = Clear, E = Entire, U = Undertake, Sr = Serrated, Mt,
178 Moist, D = Dry, Md = Muccoid, R = Rough, N = Not detected M = Moderate, R = Rod, St =
179 Short, CB = coco bacillary, C = Cocci, Cr = Curved, S = Small, L = Large

180

181 ~~M = Milky, G = Green, Cr = Creamy, W = White, S = Smooth, Or = Orange, V = Violet,~~
182 ~~Y = Yellow, Dl = Dull, O = Opaque, C = Clear, E = Entire, U = Undertake, Sr = Serrated~~
183 ~~Mt = Moist, D = Dry, Md = Muccoid, R = Rough, N = Not determined,~~
184 ~~M = Moderate, R = Rod, St = Short, CB = coco bacillary, C = Cocci, Cr = Curved, S~~
185 ~~= Small, L = Large,~~

186

187

Comment [NZ1]: For this "N" my suggestion is
If you replace it by "Nd" so it will be better
otherwise it is correct.

Comment [NZ2]: One "M" is given above for
milky, so you can change it to "Mo" or other chage
form etc.

Comment [NZ3]: Cr is given once for Creamy
you can change it to Cd or other change form etc.

Comment [NZ4]: One S is given already for
Smooth you can change it to Sm or other etc.

Formatted: English (United States)

4. DISCUSSION

-Bacteria isolated from water and sediments from Diobu creek were: *Bacillus* sp, *Pseudomonas* sp, *Corynebacterium* sp, *Acentobacter* sp, *Vibrio* sp, *Alcaligenes* sp, *Escherchhia coli*, *Micrococcus*, *Chromobacterium* sp, *Klesiella* sp, *Flavobacterium* sp. The bacterium with the highest percentage occurrence was *Bacillus* sp. The bacteria isolates from this work were similar to the isolates obtained by other workers. [17,1,13]. Similarly the current work was in line of the previous study having also more counts in wet seasons [20], the difference was attributed to increase water content of the soil in wet season. The results of similar works showed that THBCs and PDBC were more in wet season compared to dry season and the bacterium with the highest prevalence from oil polluted water and sediments was *Bacillus* sp. [1]. In a comparable work in River Nun at Amasoma, Bayelsa State, Nigeria, THBCs 1.5 to 8.67×10^5 CFU/g were obtained in sediments [21]. In Uppanar Estuary (habour) Cuddalore coast [22] THBCs 8×10^2 CFU/mL were obtained in April (summer) and 4.56×10^2 CFU/mL in December (monsoon-wet season) which showed that THBCs were season driven or influenced and sediment had higher counts of THB and PDB compared to the values obtained in water. The interaction of bacteria with benthic organisms, the chemoattractant of bacteria to nutrients at the base or bottom of the creek (on sediment) and the gentle ebbing of water which deposits most bacteria on the sediments may be responsible for the difference in counts observed in sediments and water in this study. Using chi-square $p < 0.05$, statistical analysis showed significant deference in the counts of THB and PDB in water and sediment.

Comment [NZ5]: Enter a dot “.”

Comment [NZ6]: This sign < will be used for less than not >.

In a research conducted in Awash River mouth in Ethiopia, higher counts were observed in the (Monsoon) wet season when compared with (summer) dry season [23]. In the above research, the highest THBCs in water was 2.6×10^4 CFU/mL and lowest was 9.8×10^3 CFU/mL whereas, THBCs in sediment was between 3.0×10^6 to 1.13×10^7 CFU/g. They also noted that the populations of bacteria were actually influenced by the physio-chemical parameters of the creek and THBCs were more in sediment. In an investigation of THB counts and human pathogens in Cuddalore fishing harbour after Tsunami, counts in range of 1.0×10^6 to 5.0×10^6 CFU/mL were recorded in coastal water and 5×10^4 to 1.0×10^6 CFU/mL in estuaries [24]. The results of other workers were similar to what was obtained from this study. Other researchers also had analogous results in their studies [25, 26, 27]. The THBCs were in the range of 3.6×10^3 to 1.47×10^3 CFU/mL, and THBCs 2.5×10^3 to 5.1×10^3 CFU/g were obtained in water and sediment of Persian Gulf [28]. In Padma River, Bangladesh, THBCs in sediments were 2.1×10^3 CFU/g in dry season (summer) and 3.46×10^6 CFU/g in wet season [23, 13]. In the Niger Delta Region where this work was carried out, rainfall is experienced through the year which might be why there was no significant difference in THB and PDB counts between the seasons. The most prevalent bacteria in environments highly polluted with hydrocarbons are those stimulated by their presence. The pollutant hydrocarbons and other chemicals in the creek may stimulate the boost of PDBs that were capable of degrading or mineralizing these pollutants. It is always very difficult to remove PAHs from environment due to their high hydrophobicity that increases with increasing molecular weight and this may result in high toxicity and long persistence in the environment [29]. Variety of microbes capable of degrading certain PAHs stimulates significant interest in studying microorganisms in contaminated creeks as a means of bioremediation [30]. PAHs consist of two or more than two fused benzene rings which are arranged in linear, angular or clustered forms. PAHs were present in the environment due to natural and mostly anthropogenic activities as the case in Diobu Creek, this was also noted by other researchers [6,13]. There are over 175 bacteria genera that have been known to utilizing different crude oil components as source of carbon and energy, thus petroleum hydrocarbon contaminants in aquatic environments were commonly biodegraded by insitu microbial communities [32]. PAHs are potential threat to the environment due to their toxic,

mutagenic, and carcinogenic properties [33]. Moreover some PAHs have been listed as great concern by the US Environmental Protection Agency [29, 34]. The counts of total heterotrophic bacteria and petroleum degrading bacteria are crucial to ascertain their presence and activities. The mineralization of hydrocarbons by microorganisms is the only eco-friendly means of detoxifying the environment from these pollutants.

4. CONCLUSION

The THBCs and PDBCs isolated from sediments were higher compared to the results obtained in water. The counts of THB and PDB were also increased when sampling were carried out in rain or after rainfall irrespective of the season. The increased counts of PDB in this aquatic environment were possibly stimulated by the presence of the pollutant hydrocarbons or chemicals discharged into creek which they mineralize. The activities of PDBs in a polluted environment is the most efficient and eco-friendly means of riding off pollutants from (detoxifying) the environment.

COMPETING INTERESTS

All author declared there was no conflict of interest in the course of the research and during write-up.

CONSENT (WHERE EVER APPLICABLE)

This was not applicable in this research.

ETHICAL APPROVAL (WHERE EVER APPLICABLE)

This was not applicable in this research.

REFERENCES

[1] Ichor, T., Okerentugba and Okpokwasili G., Molecular characterization of aerobic heterotrophic bacteria isolated from petroleum polluted brackish waters of Bodo Creeks, Rivers State Nigeria. Open Journal of Ecology, 2014: 4: 715 – 722.2.

[2] Kirchman, D.L, Microbial Ecology of the Oceans. 2nd Edition, 2008, John Wiley and Sons Inc., USA. <http://dx.doi.org/10.1002/9780470281840>.

[3] Alexopoulos, A., Plessas, S. and Beitzoglout, E., Water Microbial Ecology. An Overview. EOLSS, 2013, Orestiada, Greece.

[4] National Research Council (NRC), Oil in the Sea III, Inputs, Fates, and Effects National Academy, 2003, Press, Washington, D.C.

[5] ITOPE: Oil Tanker Spill Statistics: 2005. London: The International Tanker Owners Pollution Federation Ltd; 2006.

Comment [NZ7]: One space

Comment [NZ8]: Set it according to the journal

- 285 [6] Woodhead, R.J., Law, R.J., Matthiessen, P., Polycyclic aromatic hydrocarbons in surface
286 sediments around England and Wales, and their possible biological significance. Marine
287 Pollution Bulletin, 1999; 9: 773-790.
- 288 [7] Harayama, S., Kishira, H., Kasai, Y., and Shutsubo, K., Petroleum biodegradation in
289 marine environments. Journal of Molecular Microbiology Biotechnology, 1999; 1: 63-70.
- 290 [8] Sanborn, H.R., Malins, D.C., Toxicity and metabolism of naphthalene a study with marine
291 level invertebrates. Experimental Biology and Medicine, 1977; 154: 151-155.
- 292 [9] Barron, M.G., Carls, M.G., Heintz, R., Rice, S.D., Evaluation of fish early life stage toxicity
293 models of chronic embryonic exposures to complex polycyclic aromatic hydrocarbon
294 mixtures. Toxicology Science, 2003; 78: 60 - 67.
- 295 [10] Essien, J.P., Benson, N.U. and Antai, P.S. Application of Correlation Analysis in
296 Assessment of Relationships between Mineral Hydrocarbon Levels and Hydrocarbon
297 Bacteria Count in Tropical Mangrove Estuary Sediments. Scientific Research and Essays,
298 2008; 3; 94-101.
- 299 [11] McGenity, T.J., Folwell, B.D., Mckew, R.A. and Sanni, G.O., Marine Crude Oil
300 Biodegradation: A Central Role for Interspecies Interactions. Aquatic Biosystems, 2008; 8;
301 10. <http://dx.doi.org/10.1186/2046-9063-8-10>
- 302 [12] Chikere, B.C and Ekwuabu, B.C. Culture dependent characterization of hydrocarbon
303 utilizing bacteria in selected crude oil impacted sites in Bodo, Ogoniland, Nigeria. African
304 Journal of Environmental Science and Technology, 2014; 8(6): 401- 406.
- 305 [13] Subathra M.K., Immanuel G, Suresh A.H. Isolation and identification of hydrocarbon
306 degrading bacteria from Ennore creek. Bioinformation, 2013; 9(3): 150-
307 157.<http://dx.doi.org/10.6026/97320630009150>
- 308 [14] Edlund A, Jansson J.K. Changes in active bacterial communities before and after
309 dredging of a highly polluted Baltic Sea sediments. Applied Environmental Microbiology,
310 2006; 72: 680-6807.
- 311 [15] Chikere, C.B., Okpokwasili G.C., Chikere B.O., Monitoring of microbial hydrocarbon
312 remediation in the soil. Biotechnology, 2011; 1: 117-138.
- 313 [16] Kasai Y, Kishira H, Harayama S, Bacteria belonging to the genus Cycloclasticus play a
314 primary role in the degradation of aromatic hydrocarbons released in a marine environment.
315 Applied Environmental Microbiology, 2002; 68:5625-5633.
- 316 [17] Kostka J.E.^{1*}, Om-Prakash¹, O., Will A., Overholt, W.A.¹, Stefan J., Green, S.J.,
317 1,3Gina-Freyer, G., 1,2, Andy Canion¹, A., Jonathan Delgardio, J., 1 Nikita Norton¹, N.,
318 Terry C., Hazen, T.C. and Markus Huettel, M. Hydrocarbon degrading bacteria and the
319 bacterial community response in Gulf of Mexico beach sand impacted by Deep water
320 horizon oil spill. Applied and Environmental, 2011; 77(22): 7962 – 7974.
- 321 [18] IPS, Ecological post impact study of Ebubu-Ochani oil spillage. Institute of Pollution
322 Study Rivers State University of Science and Technology, Port Harcourt, Rivers State,
323 Nigeria. 1990; RSUST/IPS/TR/90/02.23 pp.

Comment [NZ9]: One space

Comment [NZ10]: Replace ; by : and One space
also with the page numbers accordingly start page
and last page

Comment [NZ11]: Replace , by .

Comment [NZ12]: Enter .Dots

Comment [NZ13]: Enter Dot

Comment [NZ14]: Enter dot

Comment [NZ15]: Enter .Dot befor and after B,
C and O.and reomove comma from the last O onl dot

Comment [NZ16]: Remove the number and
asterisk (*) from the names

Comment [NZ17]: Correct it according to th
journal

324 [19] Harrigan and McCane, Laboratory Methods in Microbiology, 1966; Academic press,
325 London and NewYork.

326 | [20] Eze, V.C. and Okpokwasili, G.C., Microbial and heavy metal characteristics of Niger
327 Delta River receiving industrial effluents. Tropical Journal of Biomedical and Allied Sciences
328 Research, 2008; 3(1): 238-249.

329 | [21] Kigigha, L.T., Seiyaboh, E.I., Obua, V.J. and Izah, S.C., Contamination of River Nun at
330 Amassoma, Bayelsa State, Nigeria Due to Microbial Diversity in Sediments. Environmental
331 Toxicology Studies Journal, 2018; 2(1): 2

332 | [22] Devanathan, K., Srinivasan, M. and Balakrishnan, S., Studies on total heterotrophic
333 population density from Uppanar Estuary (habour) Cuddalore coast. Advances in Biological
334 Research, 2010; 4(2): 139-145.

335 | [23] Lakew, W., Seenvasa, V., Prabhadevi, L., Natarajan, P. and Khilare, Y. Heterotrophic
336 bacterial population in water and sediment and fish tissues collected from Koka Reservoir
337 and Awash river Ethiopia. International Journal of Agriculture, 2015; 5(11): 1-5.

338 | [24] Mahalaksmi M., Srinivasan, M., Murugan, M., Balakrishnan, K. and Devanathan, K.,
339 Isolation and identification of total heterotrophic bacteria and human pathogens in water from
340 Cuddalore fishing harbour after Tsunami. Asian Journal of Biological Science, 2011; 4(2):
341 148-156.

342 | [25] Es, F.B. and Meyer-Reil, L.A., Biomass and metabolic activity of heterotrophic bacteria.
343 Advanced Microbial Ecology, 1982; 6:111-120

344 [26] Azam, F. France, T. Gray, J.F. MeyerReil, L.A. and Thingstad, T. The ecological role of
345 water column microbes in the sea. Marine Ecology Prog. Ser., 1983; 10: 257-263.

346 | [27] Akiko, T., Mikihide, D. and Yasuwa, F., Changes in microbial communities, including
347 uncultured and culturable bacteria, with mid-ocean blast water exchange during voyage
348 from Japan to Australia. PLOS One, 2014; 9(5)

349 | [28] Vahid, N.K., Sina, O., Iraj, N., Afshin, O., Hossein, A., Amir, V. Reghayeh, M.,
350 Mozghan, K., Fatemeh, G. and Ferzgeh, K. Indicator bacteria community in sea water and
351 coastal sediments, the Persian gulf as a case. Journal of environmental Science and
352 Engineering, 2017; 15(1):1-2.

353 | [29] Chauhan, A., Rahman, F., John, G., Oakeshott and Jain, R.K., Bacterial metabolism of
354 polycyclic aromatic hydrocarbons: strategies for bioremediation. Review of Indian Journal of
355 Microbiology; 2008; 48:95-113. <http://dx.doi.org/10.1007/s12088-008-0010-9>

356 | [30] Daane, L.L., Harjono, I., Zylstra, G.J. and Haggblom, M.M., Isolation and
357 characterization of polycyclic aromatic hydrocarbon degrading bacteria associated with the
358 rhizosphere of saltmarsh plants. Applied and Environmental Microbiology, 2001; 67(6):
359 2863-2691. <http://dx.doi.org/10.1128/AEM.67.6.2683-2691.2001>

360 | [31] Coitinho, J.B., Costa, D.M.A., Guimara, S.L. and Miranda de Góes, A., Nagem
361 RAP Expression, purification and preliminary crystallographic studies of NahF, a salicyl
362 aldehyde dehydrogenase from *Pseudomonas putida* G7 involved in naphthalene
363 degradation. Acta Crystallographic Section F: Structural Biology and Crystallization
364 Communications, 2012; 68:93- 97. <http://dx.doi.org/10.1107/S174430911105038x>

- 365 [32] Prince, R.C., Gramain, A., McGenity, T.J., Prokaryotic hydrocarbon degraders. In:
366 | Timmis, K.N. (Ed.), Handbook of Hydrocarbon and Lipid Microbiology, 2010, pp1671-e1692.
367 | Springer-Verlag, Berlin
- 368 | [33] Hemalatha S.,¹ and Veeramanikandan P.² Characterization of Aromatic Hydrocarbon
369 | Degrading Bacteria from Petroleum Contaminated Sites. Journal of Environmental
370 | Protection, 2011; 2: 243-254. <http://dx.doi.org/10.4236/jep.2011.23028>
- 371 | [34] Coral, G.¹ and Karagoz, S.² Isolation and characterization of phenanthrene-degrading
372 | bacteria from a petroleum refinery soil. Annals of Microbiology, 2005; 55(4): 255-259.

Comment [NZ18]: Set this reference according to the journal