

Bacteriological Quality and Antibigram of Isolates Obtained from Creek Town River, Odukpani L.G.A., Cross River State, Nigeria

ABSTR ACT

The bacteriological quality and antibiogram of isolates obtained from Creek Town River were investigated to ^{determine} ascertain the quality of the river and ^{water} sensitivity patterns of the bacterial isolates. The bacteriological assessment was studied using pour plate and membrane filtration techniques. The total heterotrophic bacterial and total coliform counts were shown to be highest in the CTR3 (8.0×10^6 cfu/ml) and CTR2 (3.0×10^5 cfu/ml). The total *E. coli* counts ranged from 1.5×10^2 cfu/ml to 5.8×10^2 cfu/ml. There was significant difference ($P=.05$) in total heterotrophic bacterial count and also between *E. coli* and coliform counts. Bacterial counts were higher than the acceptable limit of the WHO standards. The bacteria isolated and characterized included eleven (11) bacterial genera: *Escherichia coli*, *Klebsiella* spp., *Pseudomonas* spp., *Bacillus* spp., *Enterobacter faecalis*, *Streptococcus* spp., *Salmonella* spp., *Staphylococcus aureus*, *Proteus* spp., *Shigella* spp. and *Citrobacter* spp. Out of these organisms, ^{Eut.} *Enterobacter faecalis*, *S. aureus*, *Klebsiella* spp., *E. coli* and *Proteus* spp. are the most prevalent (100%) in the samples. The antibiogram results indicate that there is significant statistical variation ($P=.05$) in the sensitivity profile of the test organisms to antibiotics. However, *S. aureus* had the highest resistance followed by *Streptococcus* spp. *E. coli* and *Proteus* spp. ^{which} were the most sensitive among the organisms. Result also revealed that chloramphenicol was the most effective antibiotic with 100% sensitivity, and there was significant difference ($P=.05$) in the effectiveness of each drug against test organisms. These findings indicate that the water from Creek Town River is not suitable for direct human consumption and it poses a serious threat to the health of the consumers. Water treatment is therefore recommended before it can be used for domestic purposes.

Keywords: Bacteriological quality, Total heterotrophic bacteria, Total *E. coli*, Total Coliform, Creek Town River, Antibiotics, Antibigram,

1.0. INTRODUCTION

Water is a vital resource and supports life of all living organisms. Though, it occupies about 70% of the earth's surface, yet a greater percentage of the world's population, most especially in developing countries, live without access to safe water [1, 2, 3]. This is due to lack of infrastructure for the treatment of water and its eventual distribution for the populace. The multifarious uses of water for drinking, bathing, washing and cooking are well known. Water meant for human consumption should be free from pollution, safe and acceptable. The bacterial quality of water sources should not exceed the maximum limits specified in the water quality guidelines [4, 5, 6].

However, the bacterial quality of river water sources in rural communities in the Cross River State, Nigeria has been reported to be poor, unsafe and not acceptable for human consumption [7, 8]. Several bacterial enteropathogens, namely *Campylobacter jejuni*, *Salmonella*, *Shigella*, *Plesiomonas*, *Aeromonas*, *Vibrio cholera* and *Escherichia coli* were reported to have been isolated from the river water sources [9]. These enteric bacterial pathogens are variously incriminated in cases of diarrhoea, which accounts for a substantial degree of morbidity and mortality in different age groups worldwide [10].

In Nigeria, lack of efficient water supply facilities has led to the prospecting of river water by individuals especially in rural areas for the provision of drinking water. Because of the easy contamination of river water and reports on alarming prevalence of various disease-causing microbes in this water source, there seems to be increased search of underground water by rural dwellers [11, 9].

Isolation of pathogens from water sources connotes a serious public health risk for consumers. To further compound this problem, enteric bacterial pathogens have been widely reported to demonstrate resistance to several antibiotics [12, 2]. For example, in 1984, 82% of *Campylobacter* strains from Lagos, Nigeria, were sensitive to erythromycin, and 10 years later only 20.8% were sensitive [13]. In Thailand, ciprofloxacin resistance among *Campylobacter* species increased from 0% before 1991 to 84% in 1995 [12]. Strains of *S. typhi* with multiple resistances to chloramphenicol, ampicillin and trimethoprim have led to several outbreaks [14]. In the United States, several rivers were reported to be reservoirs of antibiotic resistant bacteria [2]. Antibiotic resistance may occur spontaneously, by selective pressure or because of antibiotic abuse by humans or over use in animal husbandry [9]. Although antibiotic resistance is common, antibiotics are still indicated in the management of diarrhoea. Antibiotics shorten the duration of diarrhoea, decrease stool output and may mitigate complications [15].

However, in spite of the poor water quality in rural regions of Cross River State, data on antibiotic susceptibilities of bacterial isolates from river water sources are not sufficient. Such data, if available, will be useful in the empiric management of patients with diarrhoea in the regions because antibiograms vary with time and geographical region [3, 16].

Since most of the water obtained from these sources is seldom treated because of the lack of awareness and erroneous perception that they are generally safe to drink, this study is therefore, aimed at evaluating bacteriological attributes of water samples of Creek Town River in Odukpani L.G.A, Cross River State, Nigeria.

2.0. MATERIAL AND METHODS

2.1. Study Area

The study site is a rural community in the Village known as Creek Town, located in Odukpani L.G.A., Cross River State. The occupation of the rural dwellers of this community is predominantly fishing and farming. The town is located at an elevation of 99 meters above sea level and its population amounts to 171,672. Its coordinates are $4^{\circ}48'60''\text{N}$ and $8^{\circ}16'60''\text{E}$ in DMS (degrees minutes seconds) or 4.9833 and 8.28333 (in decimal degrees). It was observed during sample collection that faeces are discharged into the Creek Town River by the rural dwellers. The river is used for bathing and washing of clothes, and sometimes drunk by children during recreational activities. The river is surrounded by plant vegetation and decaying organic matter.

? to be
really a
rural
community?

2.2. Sample collection

Water samples were collected from Creek Town River (CTR), Odukpani Local Government Area. For the purpose of clarity and reference, the selected samples were coded CTR1, CTR2, CTR3, CTR4, CTR5 and CTR6. A total of six (6) samples (3 samples in the morning and 3 samples in the evening) were randomly and aseptically collected at different points of the river. The water samples were collected in 1000ml sterile plastic bottles and transported on ice to the Microbiology Laboratory, Cross River University of Technology, Calabar for microbiological analysis. The samples were analyzed within 6 hours after collection.

?

??

2.3. Microbiological analysis samples

2.3.1. Enumeration of Total Heterotrophic Bacterial (THB) Counts

Total heterotrophic bacterial count was determined using serial dilution and pour plate method as described by Cheesbrough [17]. Ten (10) fold serial dilutions of the samples were carried out and 1ml from 10^{-4} - 10^{-6} dilutions was dispensed into sterile Petri dishes containing molten Nutrient agar in triplicates. The plates were swirled to spread the inoculum evenly within the agar medium. They were incubated for 24 hours at ambient temperature. Thereafter, the isolates are counted and recorded in colony forming units.

What is the ambient temp °C?

inoculum / inocula

in plates

are already
in plates!

Safe to write
inoculum

??

2.3.2. Screening of Total *E. Coli* and Total Coliform Counts of the Isolates

One (1) milliliter of the dilutions was also cultured on McConkey agar (MCA) for screening of Total *E. coli* count (bright pink colonies). The plates were incubated at 37°C for 24 hours before enumeration. Membrane Filtration (MF) Technique was employed in the evaluation of total coliform count. Sterilized

?

forceps were used to transfer sterilized membrane filter (47mm diameter, with 0.45µm pore size: Merck Millipore, USA) onto the porous plate of the membrane filtration unit with the grid side up and a sterile meshed funnel placed over the receptacle and locked in place. One hundred (100ml) mills of the sample water were added to the membrane filtration unit using the funnel measure. After filtration, the filtrates were discarded and the funnel unlocked and removed. In each case, sterile forceps were then used to transfer the membrane filter and then rolled over the surface of a Petri dish containing Eosin Methylene Blue agar (EMB) carefully making sure that air bubbles were not trapped between the filters and the medium. The mean total coliforms (blue-green colonies) were counted after incubation for 24hrs at 37°C. Aseptic standards were maintained throughout the experiment.

2.3.3. Isolation and Identification of Bacterial Isolates

The colonies were isolated and characterized using standard bacteriological procedures. For the isolation of *Salmonella* and *Shigella* species, 1ml water sample and 10⁻¹ serial dilution were inoculated in 9ml selenite-F-broth and incubated for 18-24h at 37°C for enrichment. The enriched samples were plated on *Salmonella-Shigella* Agar (Oxoid) and incubated for 48h at 37°C. Small colourless colonies were subcultured on nutrient agar slants and identified using methods described by Cowan and Steel [18]. For the identification of *S. aureus*, isolates were subcultured on Gelatin Mannitol Salt Agar (GMSA; Oxoid Limited, Basingstoke, UK) and incubated at 37°C for 24 hours after which growth was observed for colonies of white, surrounded by yellow zone and tested coagulase positive.

Other enterobacteria were identified by culturing samples on MacConkey agar (Oxoid) and characteristic colonies sub-cultured on nutrient agar slants for further biochemical tests. Non-enterobacterial species were identified by culturing on nutrient agar and subsequent biochemical tests carried out following standard bacteriological procedures [18].

2.4. Determination of antibiogram of isolates

Pure isolates were cultured for antibiotic susceptibility assessment using Kirby-Bauer disc diffusion method described by Jorgenson and Ferraro [19]. Pure bacterial inoculum (0.1ml), which was standardized according to 0.5 McFarland turbidity standard (10⁴ cells equivalent) was spread on Mueller-Hinton agar plates, and antibiotic discs (Antibiotic Susceptibility Discs, Oxoid Ltd., England), were aseptically placed using sterile forceps and plates were incubated for 24hrs at 37°C. Zones of inhibition were measured and compared with standard values. Antibiotics in the panel included Ciproflox (10µg), Norfloxacin (10µg), Gentamycin (10µg), Amoxil (20µg), Streptomycin (10µg), Rafamycin (20µg), Erythromycin (30µg), Chloramphenicol (30µg), Ampiclox (20µg) and Levofloxacin (20µg) with concentrations as recommended by the National Committee for Clinical Laboratory Standards [20].

2.5. Statistical analysis

Data generated from the experiment were subjected to Analysis of Variance (ANOVA) at 0.05 α- level of significance using SPSS to determine if there is significant statistical difference in bacteriological counts,

sensitivity pattern of the test organisms to antibiotics and the effectiveness of each antibiotic against bacterial isolates [21].

3.0. RESULT AND DISCUSSION

3.1 Enumeration of total heterotrophic bacterial (THB), total *E.coli* and Coliform counts

The results of total heterotrophic bacterial (THB), total *E. coli* and Coliform counts revealed a high level of bacterial load associated with the samples (Table 1). Creek Town River 3 (CTR3) had the highest THB count of 8.0×10^6 cfu/ml and sample CTR1 had the least total heterotrophic bacterial counts (4.0×10^4 cfu/ml). The results of the mean total *E. coli* count showed that 5.8×10^2 cfu/ml (being the highest) count was obtained from CTR3 while the least count (1.5×10^2 cfu/ml) was obtained from CTR6. Total coliform count obtained from the different samples ranged from 2.0×10^3 cfu/ml to 3.0×10^5 cfu/ml. According to the result, CTR2 had the highest number of coliform count (3.0×10^5 cfu/ml). This was closely followed by count from CTR4, which had 2.8×10^4 cfu/ml while CTR1 had the lowest count of 1.0×10^4 cfu/ml.

The presence of high THB, total *E. coli* and Coliform counts could be traced to anthropogenic activities, as the River is used for bathing, washing of clothes and other recreational activities indulged by children.

The high total *E. coli* and coliform counts is indicative of faecal contamination. The WHO standard for heterotrophic bacteria in potable water states that the total heterotrophic bacteria count should not be more than 100cfu/ml [22]. WHO [4] reported that the greatest risk to humans from water sanitary point of view is from faecal contamination of water supplies and that the sanitary quality of water is based on the presence and density of faecal coliform or *E. coli*. The levels of faecal coliforms in the river water could be associated with defaecation into the river by inhabitants of densely populated settlements in and around the region. Washing of faecal material deposited within adjoining land into the river by rain has its own contribution to the level of contamination of the river [23]. Ewa *et al.* [24] reported that Industrial and residential wastes were largely responsible for the temporal and spatial variation in water quality of the Calabar River.

None of the sampling points of the water sources complied with WHO standard for coliform in water and this could be supported by evidence advanced by Omoigberale *et al.* [25], who worked on seasonal variation in the bacteriological quality of Ebutte River in Ehor community, Edo State, Nigeria. According to WHO standard, every water sample that has coliform must be analyzed for faecal coliforms (*E. coli*) [26] with a view to ascertaining contamination with human or animal waste and possibly pathogenic bacteria. Results of related studies conducted in the Limpopo and Mpumalanga Provinces of South Africa showed that water sources used by the people were contaminated with potential enteric bacterial pathogens [27]. However, some of these residents do not have alternative water sources; hence, they continue using the water regardless of the risks involved. Furthermore, the level of hygiene in these rural populations is generally low [28]. Water is regarded as clean and safe for consumption when it complies with the World

Health Organization guidelines for drinking-water [29]. These guidelines are used around the world, and when drinking-water does not comply with such guidelines, it is regarded as not suitable for human use.

In this study, the results obtained indicated that the bacterial quality of the water was poor and not suitable for human consumption. This finding agrees with reports on the physiochemical and bacteriological analysis of water used for drinking and swimming purposes in Abeokuta, Nigeria by Shittu *et al.* [30]. The result is in tandem with that obtained from various water sources around the Venda region, South Africa [9]. It also corroborates the findings of Akubuenyi *et al.* [31], which reported high total heterotrophic bacterial counts, *E. coli* and coliform counts from most major sources of water for domestic uses in Calabar metropolis. The poor quality of water may cause a wide range of diseases, including cholera, typhoid fever, hepatitis, dysentery, giardiasis, and other gastrointestinal infections in rural communities. This corroborates the findings of Stanley *et al.* [32] that the water from the New Calabar River has poor microbiological quality, according to quality guidelines for drinking water

3.2. Isolation and identification of bacterial isolates

Bacterial isolates obtained from the water at various sampling points in the study and their prevalence is shown in Table 2. They include: *Enterobacter faecalis* (100%), *Escherichia coli* (100%), *Klebsiella* spp (100%), *Streptococcus* spp (66.64%), *Shigella* spp. (100%), *Salmonella* spp. (49.98%), *Staphylococcus aureus* (100%), *Pseudomonas* spp. (66.64%), *Proteus* spp (100%), *Bacillus* spp (66.64%) and *Citrobacter* spp (33.32%). The result revealed that *Enterobacter faecalis*, *Escherichia coli*, *Staphylococcus aureus*, *Klebsiella* spp and *Proteus* spp had the highest prevalence (100% each) while *Citrobacter* spp had the least prevalence (33.32%).

In a related study, the most commonly-isolated organisms from water sources included *E. coli* (22.7%), *Enterobacter aerogenes* (2.5%), *Salmonella* spp. (13.3%), *Shigella* spp. (19.3%), *Proteus* spp. (18.5%), *Klebsiella* spp. (19.3%), and *P. aeruginosa* (4.2%) [16]. *Shigella* has been identified as one of the most common pathogens responsible for the outbreaks of diarrhoea in Italy [33]. Obi *et al.* [9] reported the presence of *E. coli*, *V. cholerae*, *A. hydrophila*, *Shigella*, *Plesiomonas*, and *Campylobacter* spp. as the most common isolates in their study. In a related study, Akubuenyi *et al.* [31] recommended adequate treatment of water collected from major water sources in Calabar metropolis before consumption in order to epidemic of water related diseases. *E. coli* and *Salmonella* represents important water-borne pathogens. Enterohaemorrhagic *E. coli* have emerged as a serious gastrointestinal pathogen in many countries, albeit from consumption of contaminated meat [34]. Most of these organisms are known causative agents of outbreaks of diarrhoea in areas where there is lack of potable water. *Salmonellae* are frequently found in streams that receive sewages and industrial wastes and it is associated with salmonellosis [4]. The consumption of the river water poses potential health risk for humans.

H it = *Salmonella* 6

3.3. Antibigram of isolates

The result of the antibiotic sensitivity testing of bacterial isolates is shown in Table 3. It revealed that *E. coli* and *Proteus* spp. had the highest percentage sensitivity of 90% each. This was closely followed by *Bacillus* spp., *Enterobacter* spp. and *Citrobacter* spp. (80%) respectively. *Klebsiella* spp. and *Shigella* spp. had 70% sensitivity each while *Salmonella* spp. and *Pseudomonas* spp. had 50% each. *Streptococcus* spp. had 40% sensitivity, and the least percentage sensitivity was obtained from *Staphylococcus aureus* with 30% sensitivity.

The result of percentage sensitivity, in terms of the effectiveness of each antibiotic against all the bacterial isolates, revealed that chloramphenicol was most effective with 100% sensitivity. Streptomycin was the next antibiotic with percentage sensitivity (91%). Ciprofloxacin had 82% sensitivity while gentamycin and ampiclox had 73% each. Erythromycin had 64% and amoxil had 57%. Norfloxacin had 55% while rafamycin had the least percentage sensitivity (45%).

The antibiotic susceptibility testing is vital in the treatment and management of bacterial diseases. Unfortunately, organisms have been shown to develop resistance against commonly-used antibiotics [35]. Multiple drug resistance defined as resistance to three or more antibiotics was exhibited by *Staphylococcus aureus* and *Streptococcus* sp in this study. These findings are similar to those obtained by Obi *et al.* [9] who showed that bacterial isolates demonstrated multiple drug resistance to antibiotics. In the present study, it was found that susceptibility to amoxil, lovofoxacin, ampiclox is low. These antibiotics have been reported to be potent against a wide range of pathogens [9]. In a study in India, there was no evidence of resistance against this antibiotic, particularly among *Shigella* isolates from stool samples [36]. The high resistance exhibited by *S. aureus* could be due to the overuse of these antibiotics in the hospital environment and in the general population, leading to adaptive resistance. In a study of the isolates from river samples in the Limpopo province, the resistance level varied from 8% to 15% depending on the organisms tested, with lower resistance among *Shigella* sp. (8%) [37]. Profiles of resistance to about 20 antibiotics at a time were noted by Wasfy *et al.* [36] in Egypt where more than 25% of bacteria were resistant to three or more antibiotics. In a study in Nigeria, Oluyeye *et al.* [16] found that over 10% of bacteria were resistant to four or more antibiotics, and antibiotic resistance was the highest in members of the genera; *Enterobacter*, *Pseudomonas*, and *Proteus*. The findings of the present study suggests that the water samples collected from Creek Town river were heavily contaminated with potential bacterial pathogens that have evolved resistance to one antibiotic or the other. This poses a serious public-health threat, as children and other dwellers used the water daily. Hence, the river is unfit for drinking purposes.

The effectiveness of each antibiotic showed that all enteric bacterial isolates were markedly sensitive to chloramphenicol (100%), while streptomycin and ciproflox showed 91% and 82%, respectively. These

drugs may therefore be of value for enteric infections requiring antibiotic therapy. These reported susceptibilities are in harmony with reports of other investigators [35, 38]. It should be noted that susceptibility of bacteria to antibiotics is not static and resistance may be due to antibiotic abuse, antibiotic overuse or may be chromosomally or plasmid mediated [39]. Antibiotic usage must therefore be carefully regulated and monitored.

3.4. Statistical analysis

Analysis of variance (ANOVA) on total heterotrophic bacterial (THB) count showed that there was significant difference ($P=.05$) among the six samples. ANOVA on total coliform and faecal coliform counts were significant with $P=.05$ level of significance. There was significant difference ($P=.05$) in total heterotrophic bacterial count and also between *E. coli* and coliform counts. There was significant difference ($P=.05$) in sensitivity patterns of organisms to the different antibiotics. ANOVA also revealed that there is significant statistical difference ($P=.05$) in the effectiveness of each drug against test organisms.

Table 1: Mean Total Heterotrophic bacterial (THB), Total *E. coli* and Total Coliform counts

S/N	Sample code	THBC (CFU/ml)	<i>E. coli</i> count (CFU/ml)	Coliform count (CFU/ml)
1	CTR1	4.0×10^4	4.3×10^2	1.0×10^4
2	CTR2	2.5×10^6	4.0×10^2	3.0×10^5
3	CTR3	8.0×10^6	5.8×10^2	2.5×10^4
4	CTR4	1.7×10^6	4.6×10^2	2.8×10^4
5	CTR5	5.5×10^5	3.0×10^2	6.1×10^3
6	CTR6	3.1×10^6	1.5×10^3	2.0×10^3

CTR = Creek Town river; CFU/ml = Colony forming unit per milliliter

Table 2: Prevalence of bacterial isolates obtained from Greek Town water samples

Microorganisms isolated	CRT1	CTR2	CTR3	CTR4	CTR5	CTR6	Prevalence (%)
<i>Enterobacter faecalis</i>	+	+	+	+	+	+	100.00

<i>Escherichia coli</i>	+	+	+	+	+	+	100.00	
<i>Klebsiella</i> spp.	+	+	+	+	+	+	100.00	
<i>Streptococcus</i> spp	+	-	+	+	-	+	66.64	73
<i>Shigella</i> spp.	-	+	+	+	+	-	66.64	
<i>Salmonella</i> spp.	+	-	-	+	+	-	49.98	750
<i>Staphylococcus aureus</i>	+	+	+	+	+	+	100.00	
<i>Pseudomonas</i> spp	+	-	+	+	+	-	66.64	7
<i>Proteus</i> spp.	+	+	+	+	+	+	100.00	
<i>Bacillus</i> spp.	+	-	+	+	-	+	66.64	76.64
<i>Citrobacter</i> spp.	-	+	+	-	-	-	33.32	73.33

CTR = Creek Town river; sp. = Species; + = Present; - = Not present; % = Percentage

Table 3: Antibigram of bacterial species isolated from Creek Town River

Organisms	Antibiotics and their concentrations (mcg) and zones of inhibition (mm)										% Sensitivity
	CPX (10)	NB (10)	CN (10)	AML (20)	S (10)	RD (20)	E (30)	CH (30)	APX (20)	LEV (20)	
<i>Salmonella</i> spp.	18 (S)	13 (I)	21 (S)	20 (S)	11 (I)	13 (I)	12 (I)	19 (S)	18 (S)	0 (R)	50
<i>Enterobacter faecalis</i>	22 (S)	12 (I)	23 (S)	11 (I)	19 (S)	18 (S)	21 (S)	22 (S)	23 (S)	20 (S)	80
<i>Bacillus</i> spp.	21 (S)	23 (S)	21 (S)	9 (R)	19 (S)	12 (I)	21 (S)	20 (S)	20 (S)	19 (S)	80
<i>Pseudomonas aeruginosa</i>	18 (S)	17 (S)	11 (I)	0 (R)	19 (S)	8 (R)	13 (I)	22 (S)	19 (S)	12 (I)	50
<i>E. coli</i>	19 (S)	14 (I)	20 (S)	22 (S)	23 (S)	19 (S)	21 (S)	24 (S)	20 (S)	18 (S)	90
<i>Staphylococcus aureus</i>	10 (R)	9 (S)	13 (I)	10 (R)	19 (S)	12 (I)	11 (I)	20 (S)	0 (R)	7 (R)	30
<i>Klebsiella</i> spp.	22 (S)	21 (S)	19 (S)	20 (S)	18 (S)	11 (I)	13 (I)	24 (S)	17 (S)	0 (R)	70
<i>Streptococcus</i> spp.	11 (I)	0 (R)	13 (I)	12 (I)	18 (S)	23 (S)	19 (S)	22 (S)	8 (R)	9 (R)	40
<i>Shigella</i> spp.	23 (S)	20 (S)	21 (S)	8 (R)	19 (S)	12 (I)	22 (S)	21 (S)	19 (S)	13 (I)	70
<i>Proteus</i> spp.	19 (S)	18 (S)	18 (S)	12 (I)	20 (S)	22 (S)	16 (S)	17 (S)	19 (S)	18 (S)	90
<i>Citrobacter</i> spp.	20 (S)	11 (I)	20 (S)	21 (S)	23 (S)	21 (S)	19 (S)	20 (S)	11 (I)	17 (S)	80
% Sensitivity	82	55	73	57	90.5 91	45	64	100	73	45	

CPX = Ciproflox; NB = Norfloxacin; CN = Gentamycin; AML = Amoxil; S = Streptomycin; RD = Rafamycin; E = Erythromycin; CH = Chloramphenicol; APX = Ampiclox; LEV = Levofloxacin; R = Resistant; S = Sensitive; I = Intermediate; spp. = Species; Conc. = Concentration; *Sal.* = *Salmonella*; mcg = Micrograms; mm = Minimeter

4.0. CONCLUSION

Creek Town River contains total heterotrophic bacterial, total *E. coli* and total Coliform counts that exceeds WHO acceptable limits. The high faecal coliform count showed that the river was contaminated with human excrement. Some of the bacterial isolates identified from the water sample are known pathogenic bacteria. Appropriate water treatments or safe potable water sources should be provided in the area to improve the welfare of the riverine dwellers. There is need to educate the villagers on how to handle and locally treat water for domestic use. The government should evolve sanitation programmes and propagate these through environmental education throughout the communities in the river catchments to prevent pollution of water bodies and consequent transmission of water-related diseases. Control of human activities which could prevent faeces and refuse from entering water body is the key to avoiding bacterial contamination of the river water.

REFERENCE

1. Eja ME. Water pollution and sanitation for developing countries. Seasprint (Nig) Co., Calabar, Nigeria. 2001; pp. 2-4.
2. Ash RJ, Mauck B, Morgan M. Antibiotic resistance of Gram negative bacteria in rivers, United States of America. *Emerging Infectious Disease*. 2002; 8:7-12.
3. Itah AY, Akpan CE. "Potability of Drinking Water in an Oil Impacted Community in Southern Nigeria", *Journal of Applied Sciences and Environmental Management*. 2005;9: 135-141.
4. World Health Organization. Guidelines for Drinking Water Quality 4th ed. World Health Organization Geneva, Switzerland. 2011: 28.
5. Department of Water Affairs and Forestry (DWAF). South African Water Quality guideline for Domestic Use (2nd edn.) Pretoria .1996.
6. Water Research Commission (WRC) Quality of Domestic Water Supplies. Col.1-Assessment Guide. Water Research Commission Report No TT 101/98. Pretoria, South Africa. 1998
7. Eja ME, Ogri O R, Arikpo G E. (2003). Bioconcentration of heavy metals in surface sediments from the great Kwa River Estuary, Calabar, South Eastern Nigeria. *Journal of Nigeria Environmental Society*. 2003; 2: 247-256.
8. Ikpeme E, Mfongeh J, Enyi-Idoh K, Eja ME, Etim L. (2011). Antibiotic susceptibility profiles of enteric bacterial isolates from dumpsites utisols and water sources in a rural community in Cross River State, Southern Nigeria. *Nature and Science*. 2011;9(5): 46-50.
9. Obi CL, Potgieter N, Bessong PO, Matsaung G. Assessment of the microbial quality of river water sources in rural Venda communities in South Africa. *Water in South Africa*. 2002; 28 (3) 287-291.
10. El-sheikh SM, El-Assouli SM. Prevalence of viral, bacterial and parasitic enteropathogens among young children with acute diarrhoea in Jeddah, Saudi Arabia. *Journal of Health Population and Nutrition*. 2001; 19 (1): 25-30.
11. White DG, Piddock LJ, Maurer JJ, Zhao S, Ricci V, Thayer SG. Characterization of fluoroquinolone resistance among veterinary isolates of avian *Escherichia coli*. *Journal Antimicrobial Agents Chemotherapy*. 2000;44 (10): 2897-2899.

12. Hoge CW, Gambel JM, Srijan A, Pitarangsi C, Echeverria P. Trends in antibiotic resistance among diarrhoeal pathogens isolated in Thailand over 15 years. *Clinical Infectious Disease*. 1998;26:341-345.
13. Coker AO, and Adefeso AO. The changing patterns of *Campylobacter jejuni/coli* in Lagos, Nigeria after ten years. *Eastern Africa Medical Journal*. 1994;71:437-440.
14. Rowe B, Ward LR, Threlfau EJ. Multidrug resistant *Salmonella typhi*: a worldwide epidemic. *Clinical Infectious Disease*. 1997; 24 (1): 5106-5109.
15. Black RE. Persistent diarrhea in children in developing countries. *Pediatrics and Infectious Diseases Journal*. 1993;12:751-761.
16. Oluyeye JO, Dada AC, Odeyemi AT. Incidence of multiple antibiotic resistant Gram-negative bacteria isolated from surface and underground water sources in south western region of Nigeria. *Water Science and Technology*. 2009;59:1929-36.
17. Cheesbrough M. *District Laboratory Practices in tropical countries*, Part II, Cambridge University Press, United Kingdom .2000; pp. 182-187.
18. Cowan ST, Steel K J. *Manual for Identification of Medical Bacteria*. Cambridge University Press, UK. 1985; pp. 11115.
19. Jorgensen JH, Ferraro MJ. Antimicrobial susceptibility testing: General Principles and Contemporary Practices. *Clinical Infectious Diseases*. 1998; 6: 973-980.
- ? | 20. NCCLS. Performance standards for antimicrobial susceptibility testing: 12th Information supplement. 22(1): M2-A7 and M7-A5. 2001.
21. Miller JC, Miller JN. *Statistics for Analytical Chemistry*. Ellis Howood Limited, Chichester. 1986.
22. World Health Organization. Guidelines for drinking water quality. Third Edition, WHO press, Geneva, Switzerland. 2006; 398.
23. Abu GO, Egenonu C. The current pollution status of the new Calabar river in the Niger Delta region of Southern Nigeria: A survey of antibiogram profiles of its bacterial isolates. *African Journal of Environmental Science and Technology*. 2008;2(6):134-141
24. Ewa EE, Iwara AI, Adeyemi JA, Njar GN. Assessment of water quality of the Calabar River using Multivariate Statistical Techniques. *Journal of Applied Sciences Research*. 2013; 9(5): 3354-3363
25. Omoigberale NO, Isibor JO, Izegaebe JI, Iyamu MI. Seasonal variation in the bacteriological quality of Ebutteriver in Ehor community, Edo State, Nigeria. *American Journal of Research Communication*. 2013;1(7): 59-69.
26. European Union (EU), Council directive on the quality of water intended for human consumption. *Journal of European Communities*. 1998;330: 32-54.
27. Obi CL, Momba MN, Samie A, Igumbor JO, Green E, Musie E. Microbiological, physico-chemical and management parameters impinging on the efficiency of small water treatment plants in the Limpopo and Mpumalanga provinces of South Africa. *Water in South Africa*. 2007;33:229-38.
28. Potgieter N, Obi CL, Bessong PO, Igumbor EO, Samie A, Nengobela R. Bacterial contamination of Vhuswa—a local weaning food and stored drinking-water in impoverished households in the Venda region of South Africa. *Journal of Health and Population Nutrition*. 2005;23:150-155.

29. Al-Khatib I, Salah S. Bacteriological and chemical quality of swimming pools water developing countries: a case study in West Bank of Palestine. *International Journal of Environmental Health Resource*. 2013; 13: 17-22.
30. Shittu OB, Olaitan JO, Amusa, T S. Physiochemical and bacteriological analysis of water used for drinking and swimming purposes in Abeokuta, Nigeria. *African Journal of Biomedical Resource*. 2008 ;11: 285-290.
31. Akubuenyi FC, Uttah EC, Enyi-Idoh KH. Microbiological and Physicochemical assessment of major sources of water for domestic uses in Calabar metropolis, Cross River State, Nigeria. *Transnational Journal of Science and Technology*. 2013; 3(2): 31-44
32. Stanley HO, Immanuel OM, Nwagboso A. Water quality assessment of the New Calabbar River. *Journal of Applied Life Sciences Internationl*. 2017;15(2): 1-5
33. Blasi MF, Carere M, Pompa MG, Rizzuto E, Funari E. Water-related diseases outbreaks reported in Italy. *Journal of Water Health*. 2008 ;6:423-32.
34. Pall E, Niculae M, Kiss T, Sandru CD, Spinu M. Human impact on the microbiological water quality of the rivers, *Journal of Medical Microbiology*. 2013;62: 1635–1640.
35. Wasfy MO, Oyofa BA, David JC, Ismail TF, El-Gendy AM, Mohran ZS. Isolation and antibiotic susceptibility of Salmonella, Shigella, and Campylobacter from acute enteric infections in Egypt. *Journal of Health and Population Nutrition*. 2000;18:33-8.
36. Mota MI, Gadea MP, González S, González G, Pardo L, Sirok A. Bacterial pathogens associated with bloody diarrhea in Uruguayan children. *Review Argent Microbiology*. 2010. 42:114-7.
37. Samie A, Guerrant RL, Barrett L, Bessong PO, Igumbor EO, Obi CL. Prevalence of intestinal parasitic and bacterial pathogens in diarrhoeal and non-diarrhoeal human stools from Vhembe district, South Africa. *Journal of Health and Population Nutrition*. 2009;27:739-45.
38. Obi CL, Bessong PO, Momba MW, Potgieter N, Samie A, Igumber EO. Profiles of antibiotics susceptibilities of bacterial isolates and physico-chemical quality of water supply in rural Venda communities, South Africa. *South African Water Journal*, 2004;30(4): 515-519.
39. Eja ME, Out JU, Alobi NO, Uno UA, Obi-Abang M. An evaluation of the phytochemical and antimicrobial profiles of *Vernonia amygdalina* and bark of *Mangifera indica*. *New York Science Journal*. 2016; 9(5): 12-23.

2017 (1)
2016 (1)