

# Original Research Article

## Relations of some hydrochemicals with hydrogen sulphide levels in sediments of Lake Burullus, Egypt.

### Abstract

Lake Burullus, one of the northern deltaic lakes in Egypt, is an important economic, recreational and Fish breeding reservoir. The study used nine georeferenced stations to assess hydrogen sulphide (H<sub>2</sub>S) levels, its relationship with selected hydrochemical parameters, and the implication on this lake's biota.

The study reveals that areas mostly affected by drainage water with high load of organic matter, aid to the production of H<sub>2</sub>S into sediments and dispersion to water. The present results indicate that H<sub>2</sub>S in lake sediment increase with increasing water temperature, biological oxygen demand (BOD) and load of organic matter (OM) in water. On the other hand areas with high oxygen levels and clear water aid in reducing H<sub>2</sub>S levels in sediments as proved from correlation analysis. The distribution maps of H<sub>2</sub>S, OM and BOD reveal positive correlation of variables of organic matter and H<sub>2</sub>S. The amount of different wastes, particularly when in large quantities, increase the level of H<sub>2</sub>S, and therefore affected biota so it is highly recommended to treat wastewater to conserve the biodiversity of this lake.

**Keywords:** Lake Burullus, Pollution, Hydrochemicals, Hydrogen Sulphide

## Introduction

The drainage water carries huge amounts of deposits to the Lakes. These deposits are distributed by currents, flows and water movements along these Lakes. They are deposited on the bottom sediments of the Lakes (Saeed and Shaker, 2008). These sediments are carried annually into the lake through the drain water, sea water and wind. The lake's bottom along the northern shores extending from the lake-sea connection westwards is mainly clayey-sand; silty sand with some patches formed molluscan shells. The eastern and western regions of the lake are silty clay. The southern shore sediments which receive directly the drain discharges which formed from clay and silt with small areas covered with molluscan shells (Med. Wet. Coast Project, 2005).

Abdo (2005) explains that the total organic matter in sediments plays an important role in the accumulation and release of pollutants in lagoon water, and it is a source of nutrient for the living fauna in the lagoon.

Hydrogen sulphide concentration in water and sediment in aquatic system is considered good indicator of oxygen levels in the water and sediment as regards assessing the lake's water suitability for supporting biota (Golterman, 1975). Naturally, hydrogen sulfide occurs in the process of decomposing organic substances containing sulfur used by bacteria in anaerobic conditions (Wongsin, 2015). Also, Berner (1984) stated that surface sediments, which contain large amounts of the freshly deposited planktic organic compounds, are very important in the production of  $H_2S$  by sulfate reducing bacteria.

$H_2S$  is an extremely potent metabolic poison, lethal at low concentrations (<1 ppm) to most vertebrates alike (Evans, 1967, Smith *et al.*, 1976; Oscid and Smith 1974 a, b).

The toxicity of hydrogen sulphide for some fauna (*Tilapia gallilae*; Nauplii larvae of *Artemia salina* (*Ocenebra erinacea*) and *Idotea baltica* have been recorded by Tayel and Shriadah (1991). The aim of this research is to

study the interrelationship between selected hydrochemical variables and H<sub>2</sub>S level in the sediments of Lake Burullus, Egypt.

## 2. Material and Methods


### 2.1. Study area

Lake Burullus is located in Kafr El-Sheikh Governorate (30° 22' - 31° 35'N; 30° 33' - 31° 08'E) with an area of about 460 km<sup>2</sup>. It is situated on the eastern side of Rosetta branch of the River Nile. It occupies a central position along the Mediterranean coast of the Nile Delta. It is the second largest natural lake in Egypt after Lake Manzala (Maswada, 2004).

Lake Burullus is connected to the Mediterranean Sea through El-Burullus outlet (Boughaz El-Burullus) which is about 250 m wide and 5 m deep. The depth of the Lake varies between 40 cm in its middle sector and near the shores and 200 cm near the outlet to the sea (Zahran and Willis, 2009).

The lake receives (4 milliard m<sup>3</sup>/y<sup>-1</sup>) drainage water from several drains which were considered the main source of pollution in the lake (El-Bayomi, 1999). Drainage water is discharged into the lake through a group of pumping stations at the end tail of the drains except Gharbia drain which discharges its water freely without pumping (EMI, 2012). The estuarine water of Rosetta mouth of the River Nile is mixed with the lake water through Brimbil canal. Sea water may also flow into the lake at Burullus outlet (Al-Sayes *et al.*, 2007).

Surficial sediment samples were collected from nine stations covering Lake Burullus; (Figure 1). The description of these locations is as shown in Table (1).



**Fig.1. Sampling stations at Lake Burullus, Egypt**

**Table (1): Latitudes and longitudes of the sampling stations at Lake Burullus**

St. NO	Station name	Latitude N	Longitude E
1	El-Burullus (east)	31° 33' 29.9''	31° 04' 25.3''
2	inf. of drain 7	31° 27' 56.1''	30° 56' 17.5''
3	El-Zankah	31° 27' 53.3''	30° 47' 10.0''
4	Mastarouh	31° 29' 09.0''	30° 45' 24.4''
5	Abo-Amer	31° 26' 07.0''	30° 42' 23.3''
6	El-Tawelah	31° 23' 43.8''	30° 43' 52.8''
7	inf. of drain 8 & 9 (Shakhlobah)	31° 24' 46.9''	30° 45' 54.9''
8	inf. of drain 11 (El-Hoksa)	31° 23' 15.5''	30° 36' 15.3''
9	inf. of Brimbal Canal	31° 24' 06.0''	30° 35' 00.4''

St= station      inf= infront

## 2.2. Analytical methods



Nine geo-referenced water samples were collected within Lake Burullus. In the field, water temperature and dissolved oxygen (DO) were measured using

the DO meter (Lutron YK-22 DO meter). pH is measured using pH-meter (Model Lutron YK-2001, pH meter). EC was determined using EC-meter (Thermo, Orion 150 A+ advanced conductivity). The biological oxygen demand (BOD) determination was carried out using the conventional Winkler method (APHA, 1998). Organic matter (OM) is determined by Permanganate oxidation method (FAO, 1975).

Hydrogen sulfide is a colorless, flammable and toxic gases smell like rotten eggs, even at low concentrations (Tuntoolavest and Tuntoolavest, 2004).


Oxidation of hydrogen sulphide in natural waters either produces or consumes hydrogen ions, depending on products and other conditions (Tayel and Shriadah 1991).

Thus



In absence of biological activity, sulphide can be slowly oxidized to sulphur which then combines with remaining sulphide to form polysulphide.

Estimation of hydrogen sulphide in sediment samples occurred as follow: 0.1 –0.8gm wet acidified samples with nearly 5ml Conc  $\text{H}_2\text{SO}_4$  in closed system, (Figure 2). The involved hydrogen sulphide gas was displaced with oxygen free nitrogen gas into zinc acetate traps. The recovery of sulphide in this manner is 99% efficient. Sulphide collected in the traps was measured calorimetrically using methylene blue method (Youssef, 1999). Results are expressed as mg/gm.



**Fig. 2. The apparatus used for estimation of hydrogen sulphide**

### 2.3. Statistical analysis:

The statistical analysis for the data were carried out to determine the correlation coefficient (r) using the formula

$$r = \frac{n(\sum xy) - (\sum x)(\sum y)}{\sqrt{[n\sum x^2 - (\sum x)^2][n\sum y^2 - (\sum y)^2]}}$$

Where X the concentration of H<sub>2</sub>S and Y is the corresponding concentration of variant and n is the number of data.

### 2.4. Geo-statistical Analysis

Inverse distance weight (IDW) is a deterministic interpolation procedure that estimates values at prediction points (V) using the following equation

$$V = \frac{\sum_{i=1}^n v_i \left(\frac{1}{d_i^p}\right)}{\sum_{i=1}^n (1/d_i^p)}$$

Where d is the distance between prediction and measurement points,  $V_i$  is the measured parameter value, and p is a power parameter (Isaaks *et al.* 1989). The main factor affecting the accuracy of inverse distance interpolator is the value of the power parameter p, as well the size of the neighborhood and the number of neighbors are also relevant to the accuracy of the results (Burrough and McDonnell, 1998).

### 3. Results and discussion

Results of hydrogen sulphide concentrations in sediments as well as concentrations of some related parameters in the water as organic matter, dissolved oxygen, biological oxygen demand and hydrogen ion concentration were shown in Table (2) and the spatial distribution maps of depth, pH, EC, BOD, DO, OM and  $H_2S$  within water and sediment of Lake Burullus are as shown in Figure 3 (A-G).

$H_2S$  concentration in sediments ranged between 4.3 at Abu-Amer and 7.7 at El-Tawelah and Brinbal canal with mean value of 6.72 mg/g. The highest value was recorded at Brinbal, and may attributed to the nature of sediment characteristics of clay and high content of organic matter that aid in the release of  $H_2S$  in sediments. Radwan and Lotfy (2002) estimated that sediments of Lake Burullus have a complex nature. More specifically, the sediments change from coarse particles-sand, usually abundant in the northern coast and at the coast of islets, whereas it's muddy in the southern parts of lakes.

Organic matter also takes the same distribution of  $H_2S$  as its high percent content was found at Brinbal canal may attributed to agricultural wastes from different agricultural areas. High discharge of drained water in the southern part



of the lake led to the consumption of DO due to oxidation of such OM. This is agreed with observations of El-Ghobashy (1990) in Lake Manzala.

In Lake Burullus, the highest concentrations of organic matter and organic carbon were distributed at the western, southern and eastern parts of the lake; this agrees with Masoud (2011) and El-Alfy (2015).

The southern parts are described as having clayey sediments or fine particles which contain high amount of organic carbon not as sandy soils which are very poor with organic matters at the northern parts of the lakes (Palma *et al.* 2012).

The site at Brinbal canal is distinguished by high density of vegetation especially hydrophytes i.e. *Eichhornia crassipes* and other vegetative plants. So it's an important reason for high concentration of organic matter (OM) in these areas as may attributed to sinking and decaying of dead plants on the bottom (Nafea, 2005).

These results are in agreement with Moussa *et al.* (1994) and Khalil *et al.* (2007) for Lake Edku where the content of OM in sediment was controlled by the amount of clay and silt in addition to the plant detritus from nearby vegetative areas.

Electrical conductivity (EC) fluctuated between 3.9 at Brinbal canal (source of fresh water from Rosetta Branch /River Nile) to 30.9 ms/cm at east of El-Burulls (this sites is near from El-Boughaz area) so it may highly affected by the sea water intrusion.

Hydrogen sulphide was produced in the anoxic part of the sediment, with reduction of sulphate. It's noticeable that, the reduction of sulphate in sediments reaches a percent of nearly 13% of total organic matter in acidic conditions and to 50% in marine sediment (Kühl and Jorgesen, 1992).

pH is very significant parameter in the metabolic and physiological processes that is important in growth of aquatic organisms (Lawson, 2011).

Values of pH changed within different sites. As it was acidic especially in the



outlets of drains, and may attributed to release of different nutrients like ammonia that responsible for acidification and decreasing of pH. This is in agreement with Koerkamp *et al.* (1998) and Ibrahiem *et al.* (2012). Also Abbas *et al.* (2001) and Sayed (2003) stated that low pH values are attributed to liberation of H<sub>2</sub>S during the decompositions of OM. The highest value of pH was recorded in site 5, may attributed to high density of hydrophytes as increase of pH value is accompanied by a flourishing photosynthesizing organisms (El-Sonbati *et al.* 2009).

The excess of OM produced during photosynthesis process in the euphotic zone eventually sinks down through the water to the sediments where respiration processes dominate. The depth of the Lake does not exceed 1.5 meter, thus, a significant difference often exists between the oxygen rich euphotic zone and underlying oxygen-poor aphotic zone. The presence or absence of oxygen has significant effect on the oxidation-reduction chemistry, also attributed to the anaerobic bacteria where the biological oxygen demand is an empirical test used to determine the relative oxygen requirements needed for the biochemical decomposition and oxidation of OM and inorganic material. The highest concentrations of BOD were recorded in stations close to the point of discharges as pronounced at station 7 (drains 8&9), where huge amount of OM originated from drains led to more consumption to DO by the bacterial activities which leads to oxygen depletion and rise in H<sub>2</sub>S level in the sediment.

Sedimentary production of hydrogen sulphide can increase the oxygen demand rate of sediment leading to a reduction in dissolved oxygen in the overlying water as shown in our investigation at stations 5 and 9. Utilizing combined oxygen as sulphate, purification then occurs resulting from decomposition of OM to hydrogen sulphide as end product (Klein, 1962). From the results obtained in Table 2, its clear that, the whole water body of the lake is well oxygenated during the time of sampling with a minimum of 5.1mg<sup>-1</sup> at

station 7 (in front of drain 8,9) a maximum of  $10.6 \text{ mg}^{-1}$  at station 5 in the middle sector.

From the statistical analysis (Table 3), it is obvious that highly inverse significant proportion was observed between hydrogen sulphide and dissolved oxygen ( $r = -0.67$ ). Meanwhile, there was a positive significant correlation with organic matter ( $r = 0.74$ ). On the other hand, the relation was insignificant between hydrogen sulphide, BOD, ( $r = 0.32$ ), pH ( $-0.24$ ) and with water temperature ( $r = 0.47$ ). The distribution maps of depth, pH, EC, BOD, DO, OM and  $\text{H}_2\text{S}$  within water and sediment of Lake Burullus as shown in figure (3) proved the relation between the presences of different parameters within the  $\text{H}_2\text{S}$ , which highly attributed to drainage waters from different waste drains. El-Amier *et al.* (2016) and El-Alfy *et al.* (2017) used geostatistical and deterministic methods for creating spatial distribution maps of different pollutants in Lake Burullus.

## Conclusion

It's concluded that areas besides drainage water as waste drains recorded high levels of  $\text{H}_2\text{S}$ . There are strong relations between drained water containing low concentrations of dissolved oxygen, high concentration of BOD and high levels of OM in sediments with the levels of  $\text{H}_2\text{S}$ . Areas with low pH values or characterized by acidic nature may be indication for high levels of  $\text{H}_2\text{S}$  in sediments of lake. So it's highly recommended to reduce organic load to the lake by using different methods of remediation aid in reducing of  $\text{H}_2\text{S}$  sources in sediments to keep the aquatic life. Also removal of invasive aquatic plants from Lake Burullus' water could aid to solve such problems.

242 Table (2): Hydrogen sulphide concentration (mg/g) in sediment and  
 243 concentration of selected hydrochemicals in water of Lake Burullus.

St.	H <sub>2</sub> S mg/gm	OM %	DO mg/l	BOD mg/l	pH	T°C	EC	Depth
1	6.9	1.8	8.5	3.8	8.67	22.7	30.8	70
2	7.2	3.1	5.8	6.5	8.78	23.0	9.11	60
3	7.2	2.4	9.1	5.7	8.55	23.0	10.1	90
4	6.8	2.9	8.1	11.4	8.78	22.8	9.29	110
5	4.3	1.6	10.6	7.3	8.83	22.1	9	120
6	7.7	3.6	5.9	13.5	8.0	25.0	8.61	100
7	6.8	2.9	5.1	18.3	7.86	24.0	4.52	70
8	5.9	2.1	7.4	10.6	6.88	25.0	4.1	80
9	7.7	4.4	6.0	21.4	6.37	25.7	3.9	90
σn	0.997	0.842	1.728	5.596	0.8518	1.1935	7.74	18.72
σn.1	1.058	0.893	1.833	5.936	0.9035	1.2658	8.21	19.61
X	6.722	2.7525	7.38	10.944	8.08	23.700	9.94	87.77

244 H<sub>2</sub>S: hydrogen sulphide; OM: organic matter; DO: dissolved oxygen; BOD: biological  
 245 oxygen demand ; pH: hydrogen ion concentration; T°C: temperature and EC: electrical  
 246 conductivity.


248 Table (3): Pearson moment correlation matrix between some hydrochemical parameters and  
 249 H<sub>2</sub>S in sediments.

Variables	H <sub>2</sub> S	OM	DO	BOD	pH	T°C
H <sub>2</sub> S	1					
OM	0.743*	1				
DO	-0.675*	-0.741*	1			
BOD	0.329	0.756*	-0.648	1		
pH	-0.246	-0.524	0.471	-0.734*	1	
T°C	0.48	.692*	-0.666	0.754*	-0.917**	1


250 \*. Correlation is significant at the 0.05 level (2-tailed).

251 \*\*. Correlation is significant at the 0.01 level (2-tailed).

252  
 253 H<sub>2</sub>S: hydrogen sulphide; OM: organic matter; DO: dissolved oxygen; BOD: biological  
 254 oxygen demand ; pH: hydrogen ion concentration and T°C: temperature




A)






B)

C)



D)








Fig. 3.(A-G) Spatial distribution of depth, pH, EC, BOD, DO, OM and H<sub>2</sub>S within water and sediment of Lake Burullus.

#### 4. References

- Abbass, M., Shakweer, L.M. and Youssef, D. (2001). Hydrochemical characters of lake Edku. *Bulletin National Institute of Oceanography and Fisheries*, Egypt, 27: 65-93.
- Abdo, M. H. (2005): Assessment of some heavy metals, major cations and organic matter in the recent sediments of Bardawil lagoon, Egypt. *Egyptian Journal of Aquatic Research*, 31(1): 214-228.
- Al-Sayes. A.; Radwan, A. and Shakweer, L. (2007). Impact of drainage water inflow on the environmental conditions and fishery of Lake Burullus. *NIOF. Egyptian Journal of aquatic research*, 33(1): 312-351.
- APHA (1998). Standard methods for the examination of water and waste water, 19<sup>th</sup> Edition. American Public Health Association, American Water Work Association, Water Pollution Control Federation, Washington, D.C.



273 Berner, R. A. (1984) Sedimentary pyrite formation: an update. *Geochim*  
 274 *Cosmoshim Acta* 48:605–615

275 Burrough, P. A. and McDonnell, R. A. (1998). Principles of Geographical  
 276 Information Systems. Spatial Information Systems and Geostatistics.  
 277 Oxford University Press, New York, 333 p.

278 El-Alfy, M. A. (2015). Comarative Ecological studies on the northern Deltaic  
 279 Lakes, Egypt. Ph. D. Thesis. Mansoura University, Egypt.

280 El-Alfy, M. A.; Abd El-Azim, H. and El-Amier, Y. A. (2017). Assessment of  
 281 heavy metal contamination in surface water of Burullus Lagoon, Egypt.  
 282 *Journal of Scientific Agriculture*, 1: 233-243

283 El-Amier, Y. A.; Abd El-Azim, H. and El-Alfy, M. A. (2016). Spatial  
 284 Assessment of Water and Sediment Quality in Burullus Lake Using GIS  
 285 Technique. *Journal of Geography, Environment and Earth Science*  
 286 *International*, 6(1): 1-16.

287 El-Bayomi, G. M. (1999). Lake Burullus: A geomorphological Study. PhD.  
 288 Thesis, Faculty of Arts. Helwan University, 328pp.

289 El-Ghobashy, A. E. (1990). Biological studies on the western region of lake  
 290 Manzalah. Ph.D. Thesis. Alexandria University, Faculty of Science,  
 291 279pp.

292 El-Sonbati, M. A.; Al-Asmar, A. M.; Ahmed, M. A.; Zyada, M. and Ibrahim,  
 293 M. (2009). Assessment of some pollutants produced from industrial  
 294 activities at Damietta governorate- Egypt. *Mansoura Journal of Biology*,  
 295 36:184-185.

296 **EMI, 2012.** Egyptian ministry of irrigation, organization of mechanic and  
 297 electricity, the amount of agricultural drainage water which entered the  
 298 lake Burullus during 2012. Central administration of the central Delta  
 299 stations, Kafr El-Sheikh, Egypt.

300 Evans, C. (1967). The toxicity of hydrogen sulphide and other sulphides. *QII*  
 301 *Exp. Physiology.*, 52: 231-248.

302 FAO (1975). Permanaganate value of organic matter in natural waters. *Fisheries*  
303 *Technical Paper*, 137: 169-171.

304 Golterman, H. L. (1975). Physiological immunological approach to the  
305 physiology of Lake Ecosystem. Elsevier scientific publishing Co.  
306 Amesterdam-Oxford- New York.

307 Ibrahiem, M. S.; El-Sonbati, M. A. and El-Alfy, M. A., 2012. Detection of  
308 industrial pollution at the northeastern part of Manzala lagoon-Egypt.  
309 *Journal of Environmental Sciences*, 41(2): 211-235.

310 Isaaks, E.H. and Srivastava RM, et al. (1989). Applied geostatistics, volume 2.  
311 Oxford University Press New York.

312 Khalil, M. Kh., Radwan, A.M. and El-Moselhy Kh.M. (2007). Distribution of  
313 phosphorus fractions and some of heavy metals in surface sediments of  
314 Burullus lagoon and adjacent Mediterranean Sea. *Egyptian Journal of*  
315 *Aquatic Research*. 33 (1): 277-289.

316 Klein, L. (1962). River pollution II, Causes and effects, butter worths, London,  
317 G.B.

318 Koerkamp, P. W. G. G.; Metz, J. H. M.; Uenk, G. H.; Phillips, V. R.; Holden,  
319 M. R.; Sneath, R. W.; Short, J. L.; White, R. P. P.; Hartung, J.; Seedorf,  
320 J.; Schröder, M.; Linkert, K. H.; Pedersen, S.; Takai, H.; Johnsen, J. O.  
321 and Wathes, C. M. (1998): Concentrations and Emissions of Ammonia  
322 in Livestock Buildings in Northern Europe. *Journal of Agricultural*  
323 *Engineering Research*, 70: 79-95.

324 Kühl, M and Jorgensen B.B. (1992). Microsensor measurements of sulphate  
325 reduction and sulphide oxidation in compset microbial communities of  
326 aerobic bioflms, Appl. Environ. *Microbiology*, 58: 1164-1174.

327 Lawson, E.O. (2011). Physico-Chemical Parameters and Heavy Metal Contents  
328 of Water from the Mangrove Swamps of Lagos Lagoon, Lagos, Nigeria.  
329 *Advances in Biological Research*, 5 (1): 08-21.

330 Masoud, M. S.; Fahmy, M. A.; Ali, A. E. and Mohamed, E. A., 2011. Heavy  
 331 metal speciation and their accumulation in sediments of Lake Burullus,  
 332 Egypt. *African Journal of Environmental Science and Technology*, 5(4):  
 333 280-298.

334 Maswada, H. F., 2004. Ecological studies on Lake Borollus protected area,  
 335 Egypt. M. Sc. Thesis, Fac. Agric., Tanta Univ., Tanta, Egypt.

336 Med. Wet. Coast Poject (2005): sediment properties of Lake Burullus, Ministry  
 337 of Environmental Affairs. Puplication of National Biodiversity, 13-  
 338 578pp.

339 Moussa, A.A., El-Sabrouti, M.A., El-Rayis, O.A. and Khalil, M.Kh. (1994).  
 340 Phosphorus in sediments of lake Edku, Egypt. The influence of chemical  
 341 and grain size parameters. *Chemistry and Ecology*, 9: 31-40.

342 Nafea, E. (2005): A study on the ecology and sustainable development of the  
 343 northern delta lakes, Egypt. Ph.D. Thesis, Faculty of Science.  
 344 Mans. Univ.p.33.

345 Oseid, D. and Smith, L. (1974a). Factor influencing acute toxicity estimates of  
 346 hydrogen sulphide to fresh water invertebrates. *Water Reearch*, 8: 739-  
 347 746.

348 Oseid, D. and Smith, L. (1974b). Chronic toxicity of hydrogen sulphide to  
 349 Gammarus pseudolins. Naeus trans. Am. Fish. *philosophy Soc.*, 103:  
 350 819-822.

351 Palma, C.; Oliveira, A.; Filali, A.; Valenca, M. and Mhammdi, N., 2012.  
 352 Geochemical characteristics of water and sediment in summer period of  
 353 the Loukkos and Sebou estuaries (NW Morocco): preliminary study.  
 354 *Bulletin de l'Institut Scientifique*, Rabat, section Sciences de la Terre, 34:  
 355 69-77.

356 Radwan, A. R. and Lotfy, M. H., 2002. On the pollution of Burullus Lake water  
 357 and sediments by heavy metals. *Egypt. J. Aquat. Biol. & Fish.*, 6(4): 147-  
 358 164.

- Saeed, S. M. and Shaker, I. M., 2008. Assessment of heavy metals pollution in water and Sediments and their effect on *Oreochromis Niloticus* in the northern delta lakes, Egypt. 8th International Symposium on Tilapia in Aquaculture, 475:490.
- Sayed, E. (2003). Studies of some chemical and physical change of water and sediment of lake Idku. M.Sc. thesis chemistry Dep. Alex. Uni. 250p.
- Smith, L.; Oseid, D. and Olson, (1976). Acute and chronic toxicity of hydrogen sulphide to fathead minnow *pimephales promelas*. *Environ. Sci. Technol.*, 10: 565-68.
- Tayel, F. and Shriadah M. (1991). Effect of pollution by hydrogen sulphide on certain aquatic organisms. *Bulletin High Institutes of public health*, XXI(4): 801-813.
- Tuntoolavest, M. S. and Tuntoolavest, M. R., "(2004).Chemical analysis of water and wastewater," Chulalongkorn University Printing House, Bangkok, Thailand,.
- Wongsin, T.; Boonprab, K.; Okamoto, Y. and Salaenoi, J. (2015). Hydrogen Sulfide Distribution in Sediments Collected From Cockle Farm at Bandon Bay, Thailand. International Conference on Plant, Marine and Environmental Sciences (PMES-2015) Jan. 1-2, Kuala Lumpur (Malaysia).
- Youssef, D. H. (1999). Behaviour of some heavy metals in sulphidic Aquatic conditions. Doctor of philosophy in science, Faculty of Science, Alexandria, University.
- Zahran, M. A. and Willis, A. J. (2009). The Vegetation of Egypt. 2nd ed. Springer. Netherlands.