

Toxic Effect of African Black Soap (*Sabulun Salo*) on THE Histology of Albino Rat Stomach

ABSTRACT

Background: Sabulun salo (African traditional black soap) is widely used among various local communities in the northern Nigeria as medicinal herb. This experimental research was carried out to study the effect of oral administration of the soap on the stomach of Albino Rats. The administration lasted for the period of two weeks

Objectives: The objectives are to determine the effect of African black soap (Sabulun Salo) on the histology of stomach, hematological parameters and food and water consumption.

Methodology: 20 Albino rats used for this study were randomly divided into four groups. The first, second and third group (AC, BC, and FC) were orally administered with 2000mg/kg, 3000mg/kg and 4000mg/kg doses of sabulun salo respectively. The control group received normal saline, the animals were sacrificed under chloroform anaesthesia. The stomach was excised and processed for paraffin embedding. Blood samples were collected for haematological analyses.

Result: Incomplete metaplasia of the stomach epithelia were seen in the albino rats fed with 2000 mg/kg and 3000 mg/kg of the extract while complete metaplasia were observed among those on 4000 mg/kg. There were significant increase in white blood cell, hemoglobin and packed cell volume in the entire groups except the FC group when compared to control. There was decrease in food and water consumption across the groups

Conclusion: The results of this research have shown that the sabulun salo (African traditional black soap) is toxic to the albino rats exposed.

Keywords: sabulun salo, histological, albino rat, toxic effect, stomach

Abbreviations: UDUTH (Usmanu Danfodiyo University Teaching Hospital)

27

28 **Introduction:**

29 African traditional black soap is the type of soap that is widely used among various communities
30 within and outside the African continents [1].The name varies among various communities in
31 Africa. It is widely referred to as sabulun salo, ose dudu among the Yoruba and in Igbo as
32 Nchankota [2].This legend black soap was introduced in Ghana from Nigeria many years ago by
33 Yoruba traders doing business in Ghana [2]. Traditionally, African black soap which has in
34 combination of water, roasted plaintain skin or cocoa pod, palm oil, palm kernel oil or Shea
35 butter when put together, is collectively referred to as black soap [1]. Traditional medicine as
36 described as the combination of knowledge, practice and belief incorporating plant, animal and
37 mineral based medicine whether explicate or not, used in diagnosis, preventing or eliminating a
38 physical, mental or social diseases and which may rely exclusively on past experience from
39 generation to generation either verbally or in writing [3;4]. African Black Soap has numerous
40 benefit and importance [1]. The African Black Soap has the ability of improving or eliminating
41 uneven skin tone, razor bumps caused by ingrown hairs and skin rashes. It is an excellent for
42 cleaning up oily skin, acne due to its antiseptic properties. It is also used to prevent skin from the
43 rashes, ring worm measles, eczema and body odor. It is used in the treatment of many infectious
44 diseases caused by micro-organism. Black soap is highly used for African spiritual purification
45 [5].However, Sabulun Salo is special form of African Black Soap mainly used in the northern
46 part of Nigeria. This type is devoid of artificial caustic soda, perfume, color and any other
47 additives used in other types of African Black Soaps [6]. It is produced from the ash of straw of
48 millet and oil. The oil could be any of palm oil, groundnut oil, palm kernel oil, Shea butter.
49 These types of African Black Soap are highly medicinal as it is used orally for the treatment of
50 various forms of gastrointestinal tract infection. It is generally used in the northern part of

Nigeria for, cleaning and washing of stomach of toxic substances. The chemical constituents of sabulun salo are potassium hydroxide, sodium hydroxide, fatty acid, glycerol and other trace elements which includes selenium, copper, lead and cadmium [7]..

Materials and Methods:

Test Substance: The sabulun salo used in this study was bought from the Shehu Shagari central market in Sokoto, Nigeria.

Animal husbandry: Albino rats (About 3 weeks old) of either sex, bred in the animal house of Faculty of Pharmaceutical Sciences of Ahmadu Bello University Zaria, Nigeria, were used for the research. The animal were transferred into the animal house of the Faculty of Pharmaceutical Sciences of Usmanu Danfodiyo University Sokoto, Nigeria. They were housed in a meshed bottom cage and left for about 6 weeks for maturity and acclimatization under continuous 12h light/darkness and fed with pelletized feed from FA feeds, Sokoto. The ethical standard of handling the laboratory animals was completely adhered to during the study.

Experimentation: The total number of twenty albino rats was used for the study. The concentrations (2000, 3000 and 4000) mg/kg of weight of the sabulun salo solution were prepared. The experiment commenced on 4th June, 2015, six weeks after the purchase of the albino rats and was allowed to stay for about 14 days. At the end of the 14th day, the rats were sacrificed and the blood and the stomach organ were removed, processed, histologically and stained in the histology laboratory.

Hematological examination: The albino rats sedated with chloroform in the laboratory. Total death was prevented to allow for continuous flow of blood for proper blood collection.. The blood was then collected from the cardiac puncture using syringe and needle. And it was

discarded into plane and EDTA bottle.. The blood parameters (PCV, Hbg, WBC, Platelets and Differential Leucocytes) were determined using sysmex auto-analyzer.

Results and Discussion

. The albino rats were divided into experimental group of fifteen animals and the control group of five animals. The experimental group was divided group AC, AB and FC. And the groups were administered 2000, 3000 and 4000mg /kg of sabulun salo respectively. From fig. 1, there are decreases in both food and water consumptions as the concentrations of the sabulun salo increase along the groups when compared to control. The haematological values obtained for animals fed with different doses of sabulun salo are presented in Table 1. The results indicated that WBC, HBG and PCV increase significantly ($p < 0.05$) in the entire groups except those animals in group FC (on the highest dose) when compared with control. The platelets increase significantly ($p < 0.05$) in the entire groups when compared with control. The lymphocytes increase significantly ($p < 0.05$) in the group BC, decrease significantly ($p < 0.05$) in the group FC and the results also indicated that there are no significant differences in the group AC when compared to control.

The Monocytes decrease significantly ($p < 0.05$) in the group BC and FC when compared to control and the group AC showed no significant difference when compared to control.

The granulocytes showed no significant difference in the entire groups except those in group FC (on the highest dose) which increase significantly ($p < 0.05$) when compared to control.

93 The reduction of the water and food consumption of the exposed albino rats may be due to the
94 irritation of the gastrointestinal tract of the albino rats caused by the sabulun salo solution
95 ingestion. The reduction of water and food consumption in the exposed group also support the
96 reported findings [8] that sub lethal concentration of soap reduced the food consumption and the
97 weight of catfish. It was observed that chronic exposure of soap also reduced food consumption
98 and retard the growth of the sea bass larvae exposed significantly [9]. However, the result did not
99 observe any significant reduction in the growth, food and water consumption in albino rats
100 exposed to the soap and detergents [10]. Poor physiological conditions of the exposed rats can
101 lead to the loss of appetite [11]. The fact that stomach organ of the exposed rats showed a
102 marked histopathological changes confirm that the soap has an adverse effect on the albino rats.
103 The histopathological changes were as a result of cytotoxic agents(11) in the soap solution which
104 caused multi organs injuries.

105 This result supports research findings that ingestion of soap solution can cause multi organ
106 damage leading to nervous system disorders, respiratory illness, cancer and even death [12; 13].
107 Furthermore , the result is in line with reported pathological changes in animal and plant exposed
108 to soap. However, this study did not find cancer changes which has been suspected by some
109 researchers [16], probably due to the short duration of the study.

110 Sodium silicate and potassium hydroxide, a major components of soap has been fingered in
111 gastrointestinal irritation leading to nausea, severe digestive tract burns and weight

112 loss in an exposed animal [18]. Reduction in the blood parameters (WBC, Platelets, PCV, HBG
113 and Differential) of the exposed albino rats compared to the control rat confirm the earlier report
114 of blood parameters reduction in rabbit exposed to the soap [18]. The reduction in the blood

parameters was as a result of destruction of the cells of the albino rat by the traces of amounts of heavy metals found in the soap [19]. It was reported that blood of an animal may be adversely affected by soap used as anti bacteria because they do so by killing cells. The increase in the WBC of the exposed albino rats was as a result of immune response of the albino rat to the toxic components of the soap solution [21].

Tables and Figures:


Table 1: Hematological parameters of the experimental groups

Parameters	Groups			
	Control	Group AC	Group BC	Group FC
WBC ($\times 10^3/\mu\text{l}$)	4.10 $\pm$ 0.00 ^a	4.80 $\pm$ 0.10 ^b	4.74 $\pm$ 0.11 ^{bc}	2.78 $\pm$ 0.24 ^d
HBG (g/dl)	11.30 $\pm$ 0.00 ^b	11.82 $\pm$ 0.26 ^c	11.42 $\pm$ 0.34 ^{ab}	11.24 $\pm$ 0.09 ^{bd}
PCV (%)	45.00 $\pm$ 0.00 ^c	49.60 $\pm$ 1.82 ^b	45.40 $\pm$ 0.55 ^{ac}	44.82 $\pm$ 0.61 ^{cd}
Platelets ($10^3/\mu\text{l}$)	545 $\pm$ 0.00 ^a	693.2 $\pm$ 28.86 ^b	798.6 $\pm$ 116.95 ^c	611.80 $\pm$ 47.97 ^{ab}
Lymphocyte (%)	88.10 $\pm$ 0.00 ^d	87.48 $\pm$ 1.21 ^d	89.12 $\pm$ 0.91 ^{cd}	45.76 $\pm$ 1.84 ^b
Monocytes (%)	4.60 $\pm$ 0.00 ^e	4.86 $\pm$ 0.37 ^e	4.04 $\pm$ 0.21 ^b	3.00 $\pm$ 0.40 ^c
Granulocytes (%)	7.30 $\pm$ 0.00 ^a	7.46 $\pm$ 1.15 ^a	6.80 $\pm$ 0.78 ^a	51.24 $\pm$ 2.01 ^b


Values are presented as mean $\pm$ SD. Values with different superscripts per row are significantly different ($p < 0.05$). WBC=white blood cell, HBG=haemoglobin level, PCV=packed cell volume.

129

130 Figure1: Effect of sabulun salo on food and water consumption



131



132

133 Fig. 2: Control group of stomach histology showing the short foveolae (long arrow) and the
 134 tightly packed glands. Purplish zymogenic cells predominate at the base, and pinkish parietal
 135 cells predominate in the upper part of the glands (short arrow) H& E X 40.

136

137 Fig. 3: AC group stomach histology showing an incomplete intestinal metaplasia. The epithelium
 138 consists of a mixture of intestinal-type goblet cells (long arrow) and columnar mucus-secreting
 139 cells, morphologically resembling those of the normal gastric epithelium (short arrow).

140




Fig. 4: BC group stomach histology showing the epithelium consists of a mixture of intestinal-type goblet cells (short arrow) and columnar mucus-secreting cells, morphologically resembling those of the normal gastric epithelium (long arrow)

Fig. 5: FC group stomach histology Showing a complete intestinal metaplasia with fully developed goblet cells (short arrow) and the contour of the mucosa changes with the development of villi (long arrow).H&E X40

Conclusion

The results of this research have shown that the sabulun salo (African traditional black soap) is toxic to the albino rats exposed. It is also presumed that all other animals including man will suffer similar fate if exposed to the sabulun salo solution. Finally, it is hoped that the result from this investigation will enlighten people on the health risk posed by sabulun salo consumption.

References

- (1) Aliyu, G.M., Malik, A.I. and Kashari, O. (2012). Analysis of the antibacterial activity of African Black soap on some selected pathogens. *ARPJN Journal of Science and Technology*. 2: 2225-7217.

(2) Beauty out of African (2011). Available at shearadiance.typepad.com/shearadiance/. Accessed on 7th march, 2015

(3) Sofowora, A. (1982): Medical plants and Traditional Medicine in Africa, 1st edition. John Wiley and Sons Limited. 68-69.

(4) David, O. (2005). Brief history of black soap and ingredients (Plantain skin). Available in <http://www.oakfordavid.com>. and accessed on 27th August, 2015

(5) Karen, A. (2001). Black Soap Story: Traditional Soap making in Ghana <http://www.yendordrums.com>. 14th August 2014.

(6) Taiwo O.E. and Osinowo F.A. (2001), Evaluation of various agro waste for traditional black soap production. *Science Direct journal*. **79**: 95-97.

(7) Simmon, W.H and Appleton, H.A (2007). "The Handbook of soap manufacture". The Oil and Colour Trades Journal. London Scott, Greenwood and Son office 8 Broadway, Ludgate Hill, E.C. Available at <http://www.gutenberg.org>. Accessed on 3rd February, 2015.

(8) Esenowo, I.K. and O.A Ugwumba, (2010). Growth response of cat fish (*Clarias gariepinus*) exposed to water soluble fraction of detergent and diesel oil. *Environmental Research Journal*. **4**:298-301.

(9) Rejeki, S., Desrina, D. and Mulgana, R.A. (2006). Chronic effect of detergent surfactant (linear alkylbenzene sulfonate) on the growth and survival rate of sea bass (*Lateolabrax niloticus*) larvae. Agriculture study program fisheries Department, Faculty of fisheries and marine Science, Diponegoro university, Semarang, Indonesia. 1-18.

(10) Faremi, A.Y. and Oloyede, O.B., (2010). Biochemical Assessment of the effect of soap and industrial Effluent on some enzymes in the stomach of albino rats. *Research Journal of Environmental Toxicology*, **4**:127-133.

- 182 (11) Ogunbileje, J.O. and Akinosun, M.O. (2011). Biochemical and Hematological profile in
183 Nigeria cement factoring workers. *Research Journal of Environmental Toxicology*. **4**: 123-157.
- 184 (12) Ayandiran, T.A., Fawole, O.O., Adewoye, S.O. and Ogundiran, M.A. (2009). Bio-
185 concentration of metal in the body muscle and gut of *Claris gariepinus* exposed to sub lethal
186 concentration of soap and detergent effluent. *Journal of Cell and Animal Biology*. **3**: 113-118..
- 187 (13) Gholami, M., Fatemi, S.M.R., Falahi, M., Esmaili, A. and Mashinchiyan, A. (2010). Effect
188 of heavy metal (copper and cadmium) and detergent on white fish fry *Rutilus frisii kutum*.
189 *Research Journal of Environmental Toxicology*. **4**: 231-236.
- 190 (14) Akanji, M.A., Olagoke, O.A. and Olayode, O.B. (1993). Effect of chronic consumption of
191 metal bisulphate on the integrity of rat liver cellular system. *Toxicity*, **81**:173-179.
- 192 (15) Uaboi-Egbenni, P.O., Okolie, P.N., Adejuyitan, O.E., Sobande, A.O. and Akinyemi, O.,
193 (2009). Effect of industrial effluents on the growth and anatomical structures of *Abelmoschus*
194 *esculeutus*. *African Journal of Biotechnology*. **8**:3251-3260.
- 195 (16) Kassem., N., (2010). Allergens and toxin in laundry detergent: synthetic ingredient that are
196 harmful to people and the environment. [http://organizedwisdom.com/allergens-and-toxins-in-](http://organizedwisdom.com/allergens-and-toxins-in-laundry-detergents-synthetic-ingredient-that-are-harmful-to-people)
197 [laundry-detergents-synthetic-ingredient-that-are-harmful-to-people](http://organizedwisdom.com/allergens-and-toxins-in-laundry-detergents-synthetic-ingredient-that-are-harmful-to-people)
- 198 (18) Warne, M.J. and schifko, A.D. (1999). Toxicity of loundary detergent components to a
199 fresh watter cladoceran and their contribution to detergent toxicity. *Ecotoxicology and*
200 *Environmental Safety*. **44**:196-206.
- 201 (19) Wadaan, M.A.. and Mubarak, M. (2009). Blood chemistry changes as an evidence of the
202 toxic effect of anionic surfactant dodecyl sulphate. *Asian Journal of Scientific Research*. **2**:113-
203 118.
- 204 (20) Abulude, F.O., Ogunkoya, M.O., Ogunleye, R.F., Emidun, O. and Abulude, A.I. (2007).
205 Assessment of the content of Pb, Cd, Ni and Cr in soap and detergents from Akure, Nigeria.
206 *Research Journal of Environmental Toxicology*. **1**:102-104.
- 207 (21) Riaz, S.,A. Ahmed and S. Hasnain., (2009). Antibacterial activities of soap against daily
208 encountered bacteria. *African Journal of Biotechnology*. **8**:1431-1436.