

**ABO BLOOD GROUPS AND MALARIA PREVALENCE IN A
REFERRAL HOSPITAL IN NIGERIA**

ABSTRACT

A cross-sectional randomized study was carried out to evaluate the relationship between malaria prevalence and ABO blood groups in a referral hospital in Rivers State, Nigeria. Venous blood samples were collected from all consenting study participants, films (thick and thin) were made from each blood sample, stained with 10% Giemsa stains and viewed microscopically with oil immersion objective to detect *Plasmodium* species using standard parasitological techniques. Agglutination technique using monoclonal Antisera A, B and D was used to determine the ABO blood groups. Data generated were statistically analyzed using Chi-square test and significance level was set at $P < 0.05$. An overall malaria prevalence of 43.1% was observed among study participants. Blood groups A and O had higher malaria prevalence of 61.1% and 42.6% respectively while blood groups B and AB had lower prevalence of 28.0% and 0% respectively ($P < 0.05$). Malaria parasite density levels > 1000 parasites/ μL in relation to ABO blood groups showed a higher prevalence of 46.7% and 22.7% in blood groups A and B respectively while lower prevalence of 0% and 15.7% occurred in blood groups AB and O respectively ($P < 0.05$). The findings of this study showed that individuals with blood group O were susceptible to contracting uncomplicated malaria but had a higher resistance to developing severe malaria compared to non – O blood groups (A and B). More detailed molecular researches are needed to fully establish the relationship between malaria parasitaemia and ABO blood groups, therefore malaria interventions and control strategies should be directed equally among individuals irrespective of their blood groups.

Keywords: Malaria, ABO, Prevalence, Rivers State

1. INTRODUCTION

Malaria is a parasitic disease caused by *Plasmodium* species, it is transmitted mainly via the bites of an infected female *Anopheles* mosquitoes during a blood meal and this disease has plagued man for centuries [1]. Malaria is a serious public health concern in sub-Saharan Africa, causing high morbidity and

mortality rates [2,3]. Some individuals in malaria-endemic zones are less susceptible to malaria infection despite the prevailing high morbidity and mortality rates compared to others who experience frequent malaria bouts. Some factors such as G-6-P-Dehydrogenase activity levels, sickle cell trait, host immune response and ABO blood groups have been linked to individuals with less or increased malaria susceptibility [4]. ABO blood groups are genetic traits with different frequencies possessed by humans and are transferred from one generation to the next but some factors such as sexual relationship, race, ethnic group, socio-economic status and alleles affect the frequencies of the ABO blood groups among populations [5]. In developing new malaria control strategies and scaling up existing ones, understanding the relationship between ABO blood groups and malaria parasitaemia could aid in effectively combating malaria especially in endemic regions of sub-Saharan Africa. This study was conducted to evaluate the relationship between ABO blood groups and malaria prevalence among randomly selected consenting study participants in University of Port Harcourt Teaching Hospital (UPTH), Rivers State, Nigeria.

2. MATERIALS AND METHODS

2.1 Study Area and Population

A cross-sectional study was carried out in UPTH (Latitude 4°53'58"N and Longitude 6°55'43"E) in Obio – Akpor L.G.A., Rivers State, Nigeria using 1000 randomly selected consenting study participant using computer generated random sequence by Statistical Analysis System (SAS) software and minimum sample size determination was done using Leslie – Kish formula [6]. The study population was made up of males and females as well as individuals of different age groups. Inclusion criteria for this study was that study participants attended the selected healthcare centres, had suspected body temperatures $\geq 38^{\circ}\text{C}$ for less than 10 days, were examined by a physician and gave their oral or written consent to be part of the research. Exclusion criteria included individuals undergoing malarial treatment (or took antimalarial drugs within two weeks before the research), diagnosed with mental illness, measles, chickenpox, infected wounds, pneumonia and those with suspected body temperatures $\geq 38^{\circ}\text{C}$ but did not give their consent to be part of the study. Ethical clearance was obtained from the Rivers State Ministry of Health and UPTH Ethics Committee before the commencement of the study.

2.2 Data Collection and Analysis

The consent (oral or written) of all study participants (consent was obtained from guardians or parents of participants below 18 years) was obtained. Five

74 milliliter (5ml) of venous blood was collected from each study participant for
 75 *Plasmodium* species detection using Giemsa microscopy technique and ABO
 76 blood grouping by agglutination technique [7]. For *Plasmodium* species
 77 detection, blood films (thick and thin) were made, air dried, stained with 10%
 78 Giemsa stains for 10 minutes and then observed microscopically using oil
 79 immersion objective [7]. ABO blood grouping was done with monoclonal
 80 antisera A, B and D (Agappe Diagnostics Ltd, India) and each blood group was
 81 determined by agglutination technique [7]. The degree of *Plasmodium* species
 82 density was graded as uncomplicated (1 – 1000 parasites/ μ L) and severe (>1000
 83 parasites/ μ L) [8]. Data obtained in this study were compared with Chi square
 84 (χ^2) test (McNemar chi-square for paired and Pearson independence chi-square
 85 for unpaired comparisons) using statistical package for social sciences (SPSS)
 86 version 17. The data were presented in frequencies, percentages and means
 87 while a p-value less than 0.05 was considered significant.

88 3. RESULTS

89 An overall prevalence of 43.1% was recorded in the study and *Plasmodium*
 90 *falciparum* was the only malaria parasite species identified (Table 1). Among
 91 study subjects examined, individuals with blood group O had the highest
 92 population while those with blood group AB had the least population (Tables 1
 93 and 2). Individuals with ABO blood groups B and AB had lower malaria
 94 prevalence of 28.0% and 0% respectively while those with blood groups A and
 95 O had higher malaria prevalence of 61.1% and 42.6% respectively ($P < 0.05$)
 96 (Table 1). Malaria parasite density levels >1000 parasites/ μ L in relation to ABO
 97 blood groups showed an overall prevalence of 19.0% with higher parasitaemia
 98 of 46.7% and 22.7% in individuals with blood groups A and B respectively
 99 compared to 0% and 15.7% in individuals with blood groups AB and O
 100 respectively ($P < 0.05$) (Table 2).

101 Table 1: Malaria prevalence in relation to ABO blood groups

ABO BLOOD GROUPS	NUMBER EXAMINED	NUMBER INFECTED (%)
A	90	55 (61.1)
B	75	21 (28.0)
AB	1	0 (0)
O	834	355 (42.6)
TOTAL	1000	431 (43.1)

102 $\chi^2 = 18.970$, $Df = 2$, $P - value = 0.000$ ($P < 0.05$)

103 Table 2: Malaria parasite density levels >1000 parasites/ μ L in relation to ABO
 104 blood groups

ABO BLOOD GROUPS	NUMBER EXAMINED	NUMBER INFECTED (%)
------------------	-----------------	---------------------

A	90	42 (46.7)
B	75	17 (22.7)
AB	1	0 (0)
O	834	131 (15.7)
TOTAL	1000	190 (19.0)

$\chi^2 = 51.253$, $Df = 2$, $P - value = 0.000$ ($P < 0.05$)

4. DISCUSSION

Data obtained in this study showed that study subjects with blood groups AB and O had the least and highest population respectively. This observation is comparable to some other similar researches conducted in Nigeria and Ethiopia [4,9,10]. Malaria prevalence was highest in blood groups A (61.1%) and O (42.6%) but lower in blood groups AB (0%) and B (28.0%). This finding is comparable to similar studies that reported higher malaria parasitaemia among individuals with blood groups A and O when compared to those with blood groups B and AB [11,12,13]. Malaria parasite density levels >1000 parasites/ μ L showed significant higher prevalence in blood groups A and B compared to blood groups AB and O. This finding agrees with the reports of similar researches [10,14,15,16,17]. *Plasmodium falciparum* infected erythrocytes bind with some serum molecules whose levels are linked with the ABO locus and these are used as biomarkers for inflammatory processes and damaged vascular endothelial cells [18,19]. *Plasmodium falciparum* – infected erythrocytes form stronger and larger rosette (clustering of cells) with uninfected erythrocytes in non – O blood groups (A and B) but weaker and smaller rosettes in blood group O [20]. Also, blood groups A and B antigens are good rosetting receptors on the surface of *Plasmodium falciparum* – uninfected erythrocytes [20] and these antigens are held by PfEMP1 which is expressed on the surface of *P. falciparum* infected erythrocytes [21]. Therefore, the high severe malaria observed among individuals with blood groups A and B could be attributed to the presence of more cellular factors (e.g. complement receptor 1 and glycoproteins) in their erythrocytes which aid invasion by *P. falciparum* and rosetting.

5. CONCLUSION

More detailed molecular researches are needed to completely establish the relationship between malaria parasitaemia and ABO blood groups. Therefore, malaria therapeutic interventions and control measures should be directed equally at all individuals irrespective of their blood groups in malaria endemic regions.

COMPETING INTEREST

Authors have declared that no competing interests exists.

REFERENCES

1. Joy D, Feng X, Furuya T, Chotivanich K, Krettli A, Wang A, White N, Suh E, Beerli P. Early origin and recent expansion of *Plasmodium falciparum*. Science. 2003; 300(5617): 318 – 321.
2. Oshikoya KA. Malaria treatment in Lagos private clinics/hospitals: physicians' compliance with the World Health Organization's recommendations. Nigerian Medical Practitioners, 2006; 49(5): 102 – 110.
3. World Health Organization. World Malaria Report. Pp 1 – 195; 2012. Available at: www.who.int/malaria
4. Otajevwo FD. Prevalence of malaria parasitemia and its association with ABO blood grouping among students of Igbinedion University Okada, Nigeria. British Journal of Medicine and Medical Research. 2013; 3(4): 1164 – 1177.
5. Chakraborty S. Heterozygosity and fixation index for ABO gene in Barak Valley populations. Notulae Scientia Biologicae. 2011; 3(1): 7 – 11.
6. Araoye MO. *Research Methodology with statistics for the Health and Social Sciences*, Ilorin, Nigeria: Nathadex Publishers; 2014; P. 118 – 119.
7. Cheesbrough M. District laboratory manual for tropical countries. 2nd Edition update Volume 1. Bulterworth – Heinemann Ltd. Oxford ox28DP. Pp 249; 2005
8. Nwagha U, Ugwu V, Nwagha T, Anyaehie B. Asymptomatic *Plasmodium* parasitaemia in pregnant Nigerian women: Almost a decade roll back malaria. Transactions of the Royal Society of Tropical Medicine and Hygiene. 2009; 103(1): 16 – 20.
9. Ilozumba P, Uzozie C. Prevalence of malaria parasitemia and its association with ABO blood group in Odaokpu Area of Onitsha South Local Government Area, Anambra State, Nigeria. Nigerian Annals of Natural Sciences. 2009; 8(2): 1 – 8.
10. Zerihun T, Degarege A, Erko B. Association of ABO blood group and *Plasmodium falciparum* in Bafeno Area, Southern Ethiopia. Asian Pacific Journal of Tropical Biomedicine. 2011; 1(4): 289 – 294.

- 174 11. Onanuga A, Lamikanra A. Association of ABO blood group and
175 *Plasmodium falciparum* malaria among children in the Federal Capital
176 Territory, Nigeria. African Journal of Biomedical Research. 2016; 19(1):
177 1 – 6.
- 178 12. Sigei J. *Plasmodium falciparum* infection among individuals with
179 different ABO and Rhesus blood groups attending Kipkelion and Fort-
180 Ternan Hospitals in Kericho County, Kenya. A thesis submitted in
181 partial fulfillment of the requirement for the award of the degree of
182 Master of Science (Applied Parasitology) in the School of Pure and
183 Applied Sciences of Kenyatta University; 2015
- 184 13. Muntaka S, Opoku – Okrah C. The prevalence of malaria parasitemia
185 and predisposition of ABO blood groups to *Plasmodium falciparum*
186 malaria among donors at the Ghanaian hospital. Assumption University
187 Journal of Technology. 2013; 16(4): 255 – 260.
- 188 14. Amala S, Nwibani C. Malaria among pregnant women and children and
189 the protective roles of ABO blood groups and HB – genotype.
190 International Journal of Current Research in Life Sciences. 2015; 4(10):
191 418 – 422.
- 192 15. Degarege A, Medhin G, Animut A, Legess M, Erko B. Association of
193 ABO blood group and *P. falciparum* malaria related outcomes: a cross –
194 sectional study in Ethiopia. Acta Tropica. 2012; 123(3): 164 – 9.
- 195 16. Tadesse H, Tadesse K. Assessing the association of severe malaria
196 infection and ABO blood groups in North – Western Ethiopia. Journal of
197 Vector Borne Diseases. 2013; 50: 292 – 296.
- 198 17. Akhigbe R, Ige S, Adegunlola G, Adewumi M, Azeez M. Malaria,
199 haemoglobin genotype and ABO blood groups in Ogbomoso, Nigeria.
200 International Journal of Tropical Medicine. 2011; 6(4): 73 – 76.
- 201 18. Barbalic M, Dupuis J, Dehghan A, Bis J, Hoogeveen R, Schnabel R.
202 Large scale genomic studies reveals central role of ABO in SP – selecting
203 and SICAM – 1 Levels. Pubmed. 2010; 19: 1863 – 1872.
- 204 19. Igbineghu C, Odaibo G, Olaleye D, Odaibo A. Malaria infection and
205 ABO blood grouping in Iwo community, Southwestern Nigeria. Research
206 Journal of Medical Sciences. 2012; 6(5): 247 – 250.
- 207 20. Barragan A, Kremsner P, Wahlgren M. Blood group A antigen is a co-
208 receptor in *Plasmodium falciparum* rosetting. Infection and Immunity.
209 2000; 68: 2971 – 2975.
- 210 21. Chen Q, Heddini A, Barragan, A. The semi conserved head structure of
211 *Plasmodium falciparum* erythrocyte membrane protein 1 mediates

212 binding to multiple independent host receptors. Journal of Experimental
213 Medicine. 2000; 192: 1 – 10.