

Acute intussusceptions ileo-ileal in adult

****A case report**

Abstract

We relate here the case of a 47 year old patient with no particular history, admitted in the emergency department for an occlusive syndrome which had developed gradually. The onset of symptoms was marked by a moderate abdominal pain with bilious vomiting. The symptomatic treatment had no effect; the pain became intense and diffused to the entire abdomen accompanied by uncontrollable vomiting and the passage of gas and stool passage were stopped. Ultrasonography of abdomen showed target signs in cross section and sandwich sign in longitudinal section which are characteristic of intussusceptions. The abdominal computed tomography (CT) allows diagnostic certainty of discovering the possible etiology. It showed the presence of an intestinal occlusion. The laparotomy revealed an ileo-ileal intussusception caused by an ileal tumor. We performed a segmental small bowel resection with anastomosis. Histological study confirmed the benign nature of the tumor evoking an aspect in favor of an inflammatory pseudotumor of the small intestine.

Keywords: Acute intussusception, ileo-ileal, adult

Introduction

Intussusception or invagination of the bowel is defined as the telescoping of one portion of the bowel into an immediately adjacent portion of the bowel. Intussusception is more common in the pediatric population than in adults. The intussusception in adults is rare accounting for 5% of all cases of intussusceptions and almost 1%-5% of bowel obstruction [2,8,9]. It is an epiphenomenon revealing in 80% of cases a particular tumor organic lesion[1]. In pediatric population, the diagnosis and management are different from those of adult populations

29

30 **The observation**

31 A 47 year old patient with no particular history was admitted in emergency for an intestinal
32 obstruction which had developed gradually. He presented an abdominal colic without severe
33 pain and presented with bilious vomiting. The onset of these symptoms was marked by
34 intestinal obstruction. The pain became intense in spite of taking symptomatic treatment and
35 diffused to the entire abdomen accompanied by uncontrollable vomiting. The gas and stool
36 passage were stopped. On physical examination, the abdomen was slightly distended with
37 tenderness in the left flank. Laboratory tests were normal. The abdominal X-ray showed the
38 image of many bright arches with air-fluid levels projecting the left flank (Figure 1).

39



40

41 **Figure 1.**The abdominal X-ray: bright arches with air-fluid levels.

42

43 Moreover, ultrasonography of abdomen showed target signs in cross section and sandwich
44 sign in longitudinal section which are characteristic of intussusception (Figure 2)

45



Figure 2.Ultrasonographic image in transverse section “target” signs.

The diagnosis is confirmed by the abdominal computed tomography scan showing ileo-ileal intussusception (Figure 3).




Figure 3.Abdominal computed tomography in adult intussusception.

The laparotomy also revealed an ileo-ileal intussusception (Figure 4) with a dilated proximal small intestine.

The intussusception was due to an ileal homogeneous well circumscribed solid mass with exophytic growth into intestinal lumen (Figures 5,6). The mass was measuring $5 \times 5 \times 4.5$ cm

60 in the location mentioned above. It was reduced and a segmental small bowel resection was
61 performed.

62 Histological study confirmed the benign nature of the tumor and revealed proliferation of
63 spindle-shaped cells with infiltration of plasma cells and lymphocytes evoking an aspect in
64 favor of an inflammatory pseudotumor of the small intestine. Immunohistochemistry was not
65 undergone carried out.



66

67 **Figure 4:** Intraoperative findings: a solid, well-defined mass as lead point of intussusceptum.



68

69 **Figure 5.** The surgical specimen after resection of the small bowel.



70

71

72

73

74

75 **Figure 6.** Specimen showed a firm, circumscribed endoluminal tumor.

76 Discussion

77 The acute intussusception is a rare cause of abdominal pain and represents 1-5% of intestinal
78 obstruction in adults. It is most often in the small bowel (48% -70%). Unlike the in children
79 where it is often idiopathic, in adults it is often secondary to an organic lesion in nearly 85%
80 of cases [5]. In 90% of adult cases, predisposing lesions can be found, but in the paediatric
81 population, organic lesions are found in only 10% of the cases [1], whereas in 58% of cases
82 of large bowel intussusceptions, a malignant aetiology has is to be expected [2,10,13]. Some
83 studies showed that approximately 30% of all small bowel intussusceptions are caused by
84 malignancy, whereas the remainder is caused by benign lesions (60%) or are idiopathic
85 (10%) [9,11,16].

86 The classic pediatric presentation of acute intussusception (a triad of cramping abdominal
87 pain, bloody diarrhea and a palpable tender mass) is rare in adults [2]. The diagnosis is often
88 difficult as the symptomatology evolves spontaneous resolve by pushing at least at the
89 beginning and is usually manifested as chronic abdominal pain [10,16]. Nausea, vomiting,
90 abdominal fullness sensation, diarrhea, constipation occur usually. Bowel obstruction outset
91 can also be observed. As for the small bowel tumor diagnosis is difficult outside the
92 complications of intussusception or bowel obstruction. More rarely, gastrointestinal bleeding
93 form or Melena can dominate in case of tumor ulceration.

94 Plain abdominal films are typically the first diagnostic tool, since in most cases the
95 obstructive symptoms dominate and the clinical picture demonstrates signs of intestinal
96 obstruction and may provide information regarding the site of obstruction.

97 Ultrasonography is a useful tool for intussusception diagnosis, both in children and in adults,
98 though variable appreciation, depending on the operator [5,16,17]. The classic appearance of
99 an intussuscepted bowel in a transverse plane is called the 'target sign' and in the longitudinal
100 appearance it is usually viewed as multiple parallel lines, which is termed as the 'sandwich
101 appearance [9,12,16,17].

102 Computed tomography for adult Abdominal is the reference imaging technique. It allows
103 conducting indisputably diagnostic certainty and discovering the possible etiology. It shows
104 the presence of an intestinal occlusion, the topography and the morphological characteristics
105 of any causal lesion [14,15]. The computed tomography sensitivity varies between 58 and
106 100%. This test is currently considered as the most sensitive radiologic method to confirm

intussusception and distinguishes the presence or absence of a lead point [4,6,9,14,15]. Adult intussusception secondary to inflammatory tumor can be demonstrated by MRI [15]. But the laparoscopy has also been used successfully in selected cases [7]. Among the adults 70 to 90% of cases of intussusception require definite treatment, of which surgical resection is, most often, the treatment of choice [2].

The term “inflammatory pseudotumor” has been used for any macroscopic or microscopic tumor [1]. Different terms have been used: Vanek’s tumour, Inflammatory myofibroblastic tumor (IMFT), inflammatory fibroid polyps, plasma cell pseudotumour, inflammatory myofibro histiocytic proliferation, and omental mesenteric myxoidhamartoma [3,12,13]. It was first described as polypoid fibroma by Konjetzny in 1920, then by Vanek in 1949. It was so called Vanek’s Tumour. Finally it was named as inflammatory fibroid polyps in 1953 by Helwig and Rainer, indicating that its nature was probably inflammatory [13]. The etiology is still unknown. Authors think that development of this tumor occurs after trauma surgery or infection, such as Epstein-Barr virus and human herpesvirus, related with reactive cytokine production. Histologically, it is characterized by a cellular spindle cell proliferation in a myxoid to collagenous stroma with a prominent inflammatory infiltrate composed primarily of plasma cells and lymphocytes, with occasional admixed eosinophils and neutrophils [12].

Conclusion

The acute intestinal obstruction by intussusception secondary to a small tumor is rarely seen in adults. Its symptoms are not specific. The diagnosis is facilitated by the COMPUTED TOMOGRAPHY scan. Surgical excision is the treatment of choice.

Consent Disclaimer:

As per international standard or university standard, patient’s consent has been was collected and is preserved by the authors.

References

1. Chiang JM, Lin YS. Tumor spectrum of adult intussusception. *J Surg Oncol.* 2008;98:444-7.

2. Begos D G, Sandor A, Modlin I M. The diagnosis and management of adult intussusception. *Am J Surg*. 1997;173:88-94
3. F.M.C.D.M. Fletcher, K. Unni, World health organization classification of tumours pathology and genetics of tumours of soft tissue and bone, *Cancer* 177 (2002) 1365–1376.
4. Marinis A, Yiallourou A, Samanides L, Dafnios L, Anastasopoulos G, Vassiliou L, Theodosopoulos T. Intussusception of the bowel in adults: A review; *World J Gastroenterol* 2009 January 28; 15(4): 407-411
5. Cerro P, Magrini L, Porcari P, De Angelis O. Sonographic diagnosis of intussusceptions in adults. *Abdom Imaging* 2000;25: 45-47
6. Kim YH, Blake MA, Harisinghani MG, Archer-Arroyo K, Hahn PF, Pitman MB, Mueller PR. Adult intestinal intussusception: CT appearances and identification of a causative lead point. *Radiographics* 2006; 26: 733-744
7. McKay R. Ileocecal intussusception in an adult: the laparoscopic approach. *JSLs* 2006; 10: 250-253
8. Susan M. Cera Intestinal Intussusception Clinics in colon and rectal surgery/volume 21, number 2 2008
9. Azar T, Berger DL. Adult intussusception. *Ann Surg* 1997;226: 134-138.
10. Wang LT, Wu CC, Yu JC, Hsiao CW, Hsu CC, Jao SW. Clinical entity and treatment
11. Zubaidi A, Al-Saif F, Silverman R. Adult intussusception: a retrospective review. *Dis Colon Rectum* 2006; 49: 1546-1551
12. Cláudia Paiva*, Filomena Soares, Raquel da Inez Correia, Vítor Valente. Inflammatory myofibroblastic tumor presenting as ileocecal intussusception—A case report. *International Journal of Surgery Case Reports* 24 (2016) 146–149
13. Bhbhavuray T, Madhu CP, Sudhir S, Shreeharsha MV. Ileo-ileal Intussusception in an Adult Caused by Vanek's Tumour: A Rare Case Report. *Journal of Clinical and Diagnostic Research*. 2013 Dec, Vol-7(12): 2994-2995.
14. Low HM, Chinchure D. Clinics in diagnostic imaging. *Singapore Med J* 2016; 57(12): 664-668
15. Feldis M, Dilly M, Marty M, Laurent F, Cassinotto C. An inflammatory fibroid polyp responsible for an ileal intussusception discovered on an MRI. *Diagnostic and Interventional Imaging* (2015) 96, 89—92.

- 169 16. Satoshi Idaa, Hosei Matsuzakib Shinichi Kawashimac Masayuki Watanabea Yasuhiro
170 Akiyamab Hideo Babaa. Adult Intestinal Intussusception Caused by an Inflammatory
171 Myofibroblastic Tumor
- 172 17. Onkendi EO, Grotz TE, Murray JA, Donohue JH: Adult intussusception in the last 25
173 years of modern imaging: is surgery still indicated? J Gastrointest Surg 2011;
174 15:1699–1705.
- 175