

Original Research Article

The Nature and Complications of Acute Traumatic Cervical Spinal Cord Injury in a University Hospital in Nigeria

Abstract

Background: Traumatic cervical spinal cord injury is one of the most devastating types of injuries. It results in varying degrees of disabilities that adversely affect patients' quality of life. This study aims to evaluate the nature and complications of traumatic cervical spinal cord injury in Benue State University Teaching Hospital, Makurdi, Nigeria.

Methods: A retrospective study of traumatic cervical spinal cord injury patients who presented to BSUTH from June 2012 to May 2017. Socio-demographic and clinical characteristics were extracted from hospital records and analyzed with SPSS version 21.0 (Armonk, NY: IBM Corp.).

Results: A total of 47 patients with traumatic cervical spinal cord injury were studied. There were 42 males (89.4%) and 5 females (10.6%). Their ages ranged from 17 to 77 years with a mean of 35 ± 16 . The common causes were road traffic injury (n=29, 61.7%), falls from height (n=8, 17.0%) and falling objects (n=5, 10.6%). Most patients were transported to the hospital in saloon cars (n=34, 72.3%). Of these patients, 3 (6.7%) had atlanto-axial (C1–C2) injuries while 44 (93.3%) had subaxial (C3–C7) injuries. Associated injuries were identified in 48.9% of patients with head injury being the most predominant. Four cases had established pressure sores on presentation. Twelve patients (19.7%) had skull traction while the rest had orthosis. Majority of the patients had ASIA class A (n=31, 66.0%) at discharge. One patient (2.1%) had no neurological deficits (ASIA class E). Complications were observed in 63.8% of the patients, with skin PU being the most frequent (36.6% of all complications). Other complications included UTI, respiratory tract infection, depression, neurogenic pain, spasms and contractures.

Conclusion: Traumatic cervical spinal cord injury predominantly affected young male patients. It mostly involved the subaxial spine. Majority of patients had American Spinal Injury Association Class A injury. Pressure ulcers and urinary tract infection were the most common complications encountered.

Key words: Cervical spine injury, pressure sores, urinary tract infection, developing countries, Nigeria

Introduction

Traumatic cervical spinal cord injury (TCSCI) accounts for 2–3% of all trauma patients and 8.2% of trauma-related deaths [1]. Spinal cord injury (SCI) is one of the most devastating types of injury resulting in varying degrees of paralysis, sensory loss, and bladder/bowel dysfunction [2]. Suspicion, early diagnosis of injury, preservation of spinal cord function and maintenance or restoration of spinal alignment and stability are necessary for successful treatment of the acute injury [1]. The best outcomes in treating traumatic SCI are achieved by managing the acute injury, providing comprehensive rehabilitation and ensuring adequate support to allow community reintegration [3]. Despite great improvements in the management of acute traumatic SCI, resource-constrained settings have not benefitted from this progress to the same extent as have countries with vast resources [4].

Paucity of dedicated spine injury centers in most Sub-Saharan countries has hampered favourable outcomes. Lack of ventilators, pressure-relieving surfaces, wheel chairs, home-care teams are some of the problems militating against effective management. This has resulted in avoidable complications which then increase morbidity and mortality. In-hospital mortality rates after acute traumatic SCI are higher in resource-constrained settings mainly due to pressure ulcers (PU) and urinary tract infections (UTI), respiratory complications, delays in admissions and multiple hospital presentations [5].

The cervical spine is the most predominant segment of the spine involved in injury [5], [6]. This is due to its greater mobility when compared to other segments of rendering it more vulnerable to dangerous loading following sudden application of forces [7]. TCSCI is associated with significantly more morbidity and mortality than SCI involving other segments [8], [9]. Possible involvement of respiration and all limbs is a major challenge to independent post-rehabilitation survival. Complications have a considerable impact on these patients and in order to optimize the individual rehabilitation process and outcome, it is important to predict and prevent complications in addition to managing them when they occur [10], [11].

This study aims to study the pattern and associated complications in patients with acute TCSCI managed at Benue State University Teaching Hospital, Makurdi, Nigeria (BSUTH).

Patients and methods

This was a retrospective observational study of all patients with acute TCSCI seen from June 2012 to May 2017 at BSUTH. The hospital is a tertiary care center and teaching hospital for the Benue State University. It is a 350-bed hospital and serves as the major treatment center for trauma patients over an area with a population of about 240 000.

All patients with TCSCI admitted within 6 weeks of injury were included in this study. Patients with incomplete medical records and those who presented later than 6 weeks following their injuries were excluded from the study.

Patients' case notes were reviewed for demographic and clinical variables including age, gender, cause of trauma, associated injuries, neurological status, treatment given and clinical complications and length of hospital stay.

Neurological status was recorded and classified using the American Spinal Injury Association (ASIA) guidelines [12]. Motor and sensory examinations were

performed routinely while on admission and at discharge. TCSCIs were defined as complete when there was no motor or sensory function in the lowest sacral segment, whereas an incomplete injury is characterized by the presence of sacral sparing.

The conservative treatment for TCSCI involved measures employed to maintain alignment of the cervical spine employing cervical collar or skull traction using Gardner-Wells or Crutchfield calipers. Patients treated using skull traction lie supine in bed for 6 weeks then rehabilitated while keeping a cervical collar on.

Statistical analysis was carried out using the software IBM SPSS Statistics for Windows, Version 21.0. (Armonk, NY: IBM Corp.) Descriptive statistics were used to display single variable quantities using means and standard deviations (SD) for continuous variables or proportions for categorical variables unless otherwise stated.


Results

Of the forty seven patients included in this study, 42 (89.4%) were male and 5 (10.6%) female making the ratio of men to women 8.4:1. Their ages ranged between 17 and 77 years with a mean of 35.15 ± 15.9 . Students constituted the majority of patients (n=21, 44.7%). They were followed by farmers (n=10, 21.3%), civil servants (n=7, 14.9%), traders (n=4, 8.5%), artisans (n=3, 6.4%) and retirees (n=2, 4.3%). Fig. 1 shows the distribution of patients according to gender and age groups.

The commonest cause of traumatic cervical SCI was road traffic injury (RTI) (n=29, 61.7%). The other causes were fall from height (n=8, 17.0%), falling objects (n=5, 10.6%), diving into shallow pools (n=3, 6.4%) and firearm injuries (n=2, 4.3%). Only 3 patients (6.4%) were transported to the hospital via ambulances; most patients were via saloon cars (n=34, 72.3%). Other modes of transport to hospital were via commuter buses (n=7, 14.9%) and motorcycle taxis (n=3, 6.4%). Of these patients, 3 (6.7%) had atlanto-axial (C1–C2) injuries while 44 (93.3%) had subaxial

(C3–C7) injuries. Associated injuries were identified in 48.9% of all patients, with head injury accounting for 69.2% of all associated injuries. Long bone fractures were the second most frequent (23.1% of all associated injuries). Other associated injuries included chest injury and facial trauma. Four cases had established pressure sores on presentation. Twelve patients (19.7%) had skull traction while the rest had orthosis. Majority of the patients had ASIA class A (n=31, 66.0%) at discharge. One patient (2.1%) had no neurological deficits (ASIA class E) Table 1 shows the demographic and clinical characteristics of patients. Complications were observed in 63.8% of the patients, with skin PU being the most frequent (36.6% of all complications). Other complications included UTI, respiratory tract infection, depression, neurogenic pain, spasms and contractures. Figure 2 shows the distribution of complications among patients. The length of hospital stay ranged from 1 to 84 days with a mean of 33.8 days. Seven patients died giving a case fatality rate of 15.2%.

141 **Figure 1: The distribution of patients according to gender and age groups**



142

143

144

145

146

147

148

149

150


151 **Table 1: Demographic and clinical characteristics of patients**

Variable	Frequency	Percentage
Occupation of patients, n=47		
Student	21	44.7
Farming	10	21.3
Civil service	7	14.9
Business	4	8.5
Artisanship	3	6.4
Retirees	2	4.3
Aetiology of TCSCI* , n=47		
RTI**	29	61.7
Fall from height	8	17.0
Falling objects	5	10.6
Diving into shallow pools	3	6.4
Firearm injuries	2	4.3
Means of transport of patients to A&E, n=47		
Saloon cars	34	72.3

Commuter buses	7	14.9
Ambulance	3	6.4
Motorcycle taxis	3	6.4
Type of TCSCI, n=47		
Atlanto-axial injuries	3	6.7
Subaxial injuries	44	93.3
SEVERITY OF INJURY (ASIA[#])		
Grade A	31	66.0
Grade B	5	10.6
Grade C	7	14.9
Grade D	3	4.9
Grade E	1	2.1

*Traumatic cervical spinal cord injury, **Road traffic injuries, #American Spinal Injury Association

159 **Figure 2: The distribution of complications among patients**



160
161
162

163

164

165

166

167

168

Discussion

TCSCI remains an important public health problem associated with significant morbidity and mortality [7], [8]. In this study, the majority of patients were predominantly young adult males. This is in keeping with findings from other studies [8], [13]. The high proportion of this group of patients may be due to the fact that young males are more mobile and participate in high risk-taking activities. Involvement of the economically productive age-group places a huge toll on national productivity. The multiplier effect on the society is evident considering the fact that the affected population, in a country with no social security, are usually breadwinners for their families. The adverse effects of SCI are not limited to the individual and family alone but the entire society [14]. There ought to be an urgent public policy response to address the causes of SCI in Nigeria.

RTIs were the most common cause of TCSCI in this study. This is similar to findings in other studies [7], [13]. RTIs are said to be the leading cause of injury-related mortality in both developed and developing countries with developing countries bearing a disproportionately high share of the burden [15]. Some of the reasons RTIs abound in developing countries are growth in motor vehicle numbers and poor enforcement of traffic safety regulations [16]. The absence of a formal pre-hospital trauma care in the study area means urgent airway management and protection of the injured spine to prevent aggravation of injury cannot be guaranteed. This is likely to be a major contributing factor to the morbidity and mortality of TCSCI.

Accidental falls were the next most important cause of cervical spine injury in this study. Falls play an important role in the causation of cervical SCI [8], [17]. While falls from height are responsible for most cases of fall-related SCI, low-level falls are of importance in populations with a lot of elderly patients [18]. Most falls in this study were from heights among palm wine tappers, electricians and carpenters. Some of the workers use inefficient locally-fabricated safety equipment. Establishment and enforcement of complete fall protection protocol

for people at risk of falling while working at heights may help reduce fall-related cervical SCI in Nigeria. The program should include training users, selecting, fitting, and inspecting equipment.

Firearm injuries were the least common cause of cervical SCI in this study. This may be because the neck presents a small surface area compared to other parts of the spine and that the injuries may be so severe that the patients do not make it to the hospital. The cavitation effect of high-velocity injuries explain some cases of complete cord injuries seen with no vertebral fractures.

The mean duration between injury and presentation to hospital was 4.6 days. Late presentation has been reported in similar studies done in developing countries [19], [20]. Patients remain in peripheral hospitals in sub-optimal conditions that may lead to the development of complications resulting in increased morbidity and mortality. Financial constraints have been cited as important impediment to effective management of cervical SCI in Nigeria [21]. Almost all patients cater for their hospital bills. The National Health Insurance Scheme at the moment is not comprehensive with regards SCI. Extending comprehensive insurance coverage to the entire population may help curb the menace of delayed presentation among these patients.

Majority of patients (93.7%) showed involvement of the subaxial region of the cervical spine with C 5 being the most commonly involved. Involvement of the atlanto-axial region may reflect poor initial survival precluding them from presenting to hospitals. There was no neurological deficit in 3 cases, 13 were motor and sensory incomplete and 31 were motor and sensory complete.

PU were the most common complication noted in this study. They are a frequent cause of morbidity among spinal cord-injured patients [6], [10], [19] The causation of PU is multifactorial. While direct pressure or shear over bony prominences is necessary, there are multiple systemic, internal, and external factors that contribute to the development of PU, including but not limited to such divergent factors as skin moisture level, nutritional factors, psychosocial, and cognitive issues [22]. They are costly to manage and result in prolonged hospitalizations,

229 delayed community reintegration, reduced quality of life, and loss of self-esteem
230 [22], [23]. Prevention of these ulcers is one of the cornerstone principle of
231 management. This involves education, involvement and education of family,
232 increased nurse-to-patient ratio, optimized nutrition, maintaining good skin
233 hygiene, having pressure relief surfaces and finally having 'turning teams' or other
234 ways of regular pressure relief [24]. Treatment of the ulcers when they occur is
235 paramount. However insufficient human and material resources are an
236 impediment to efficient implementation of these measures.

237 A SCI may interrupt the communication between the pontine micturition center
238 and the spinal cord in cervical SCI causing neurogenic lower urinary tract
239 dysfunction [25]. This may result in incontinence and bladder-emptying difficulties
240 which then predispose to, among others, UTI. UTI is a major cause of morbidity
241 and mortality among SCI patients. It was the second most prevalent complication
242 in this study. The neurogenic bladder in SCI patients is predisposed to an
243 increased risk of infection because of incomplete voiding, elevated intra-vesical
244 pressure and catheter use [26]. All the patients in the study had indwelling
245 urethral catheters as the method of urinary drainage during the spinal shock and
246 rehabilitation periods. The use of an indwelling urethral catheter (IUC) is an
247 important risk factor for UTI. Clean intermittent catheterization (CIC) is said to be
248 the method of choice of bladder management in SCI patients during the
249 rehabilitation phase. CIC (that is, self-catheterization) decreases complication
250 rates compared with IUC and is recommended for the management of neurogenic
251 bladder dysfunction [27]. However, IUC are used to a large extent in resource-
252 constrained settings because of a lack of catheters and staff to assist with
253 catheterization, and a resistance among patients to self-catheterize [26].

254
255
256 Conclusions: : TCSCI predominantly affects young male patients. It mostly involves
257 the subaxial spine. Majority of patients had ASIA Class A Injuries. Pressure ulcers
258 and urinary tract infection were the most common complications encountered.

References

- 1 Ning G.-Z., Wu Q., Li Y.-L., Feng S.-Q. Epidemiology of traumatic spinal cord injury in Asia: a systematic review. *Journal of Spinal Cord Medicine*. 2012;35(4):229–239.
- 2 Pagliacci MC, Franceschini M, Di Clemente B, Agosti M, Spizzichino L. A multicentre follow-up of clinical aspects of traumatic spinal cord injury. *Spinal Cord* 2007;45: 404–410.
- 3 Burns AS, O’Connell C. The challenge of spinal cord injury care in the developing world. *The Journal of Spinal Cord Medicine* 2012; 35(1): 3 -8.
- 4 Löfvenmark I, Hasselberg M, Nilsson Wikmar L, Hultling C, Norrbrink C Outcomes after acute traumatic spinal cord injury in Botswana: from admission to discharge. *Spinal Cord*. 2017; 55(2):208-212.
- 5 Nwankwo OE, Uche EO. Epidemiological and treatment profiles of spinal cord injury in southeast Nigeria. *Spinal Cord* 2013;51:448-52.
- 6 Obalum DC, Giwa SO, Adekoya-Cole TO, Enweluzo GO. Profile of spinal injuries in Lagos, Nigeria. *Spinal Cord* 2009;47:134-7.
- 7 Uche E O, Nwankwo O E, Okorie E, Nnezianya I. Cervical Spine Injury: A ten-year multicenter analysis of evolution of care and risk factors for poor outcome in southeast Nigeria. *Niger J Clin Pract* 2015;18:203-8
- 8 Santos EAS, Filho WJS, Possatti WL, Bittencourt LRA, Fontoura EAF, Botelho RV. Clinical complications in patients with severe cervical spinal trauma: a ten-year prospective study *Arq Neuropsiquiatr*. 2012;70(7):524-8
- 9 Yiltok SJ, Onche II, Obiano SK. Cervical Spine Injury: Nature and Complication *Nigerian Journal of Orthopaedics and Trauma* 2004; 3(2): 149-163
- 10 Kawu AA, Alimi FM, Gbadegesin AAS et.al. Complications and Causes of Death in Spinal Cord Injury Patients in Nigeria *WAJM* 2011; 30(4): 301–304.
- 11 White AA, Southwick WO, Panjabi MM. Clinical instability in the lower cervical spine. a review of past and current concepts. *Spine* 1976;1:15-27.

- 12 Maynard FM Jr, Bracken MB, Creasey G, et al. International standards for neurological and functional classification of spinal cord injury. American Spinal Injury Association. *Spinal Cord* 1997;35:266-274.
- 13 Frielingsdorf K, Dunn RN. Cervical spine injury outcome – a review of 101 cases treated in a tertiary referral unit. *SAMJ* 2007; 97(3): 203-207.
- 14 Obalum DC, Giwa SO, Adekoya-Cole TO, Enweluzo GO. Profile of spinal injuries in Lagos, Nigeria. *Spinal Cord* 2009;47:134-7.
- 15 World Health Organization. WHO: Global Status Report on Road Safety 2015. Available from http://www.who.int/violence_injury_prevention/road_safety_status/2015/en/ [Last accessed 2016 June 30]
- 16 Garrib A, Herbst AJ, Hosegood V, Newell M. Injury mortality in rural South Africa 2000-2007: rates and associated factors. *Trop Med Int Health*. 2011;16: 439–46.
- 17 Torretti JA, Sengupta DK. Cervical spine trauma. *Indian Journal of Orthopaedics*. 2007;41(4):255-267.
- 18 Feng HY, Ning GZ, Feng SQ, Yu TQ, Zhou HX. Epidemiological profile of 239 traumatic spinal cord injury cases over a period of 12 years in Tianjin, China. *J Spinal Cord Med* 2011;34(4):388–94
- 19 Yongu WT, Elachi CI, Mue DD, Kortor JA. Pattern of traumatic spinal cord injury in Makurdi, Nigeria. *Niger J Surg Res* 2016;17:53-7.
- 20 Gosselin RA, Coppotelli C. A follow-up study of patients with spinal cord injury in Sierra Leone. *Int Orthop* 2005;29(5):330–2.
- 21 Ogungbo B. Anterior decompression, fusion and plating in cervical spine injury: Early experience in Abuja, Nigeria. *Surgical Neurology International*. 2011;2:156.
- 22 Consortium for Spinal Cord Medicine Clinical Practice Guidelines. Pressure ulcer prevention treatment following spinal cord injury: a clinical practice guideline for health-care professionals. *J Spinal Cord Med* 2001; 24 (Suppl 1): S40–S101.
- 23 Dealey C, Posnett J & Walker A. The cost of pressure ulcers in the United Kingdom. *Journal of Wound Care* 2012; 21(6): 261–66.

- 321 24 Zakrasek EC, Creasey G, Crew JD. Pressure ulcers in people with spinal cord
322 injury in developing nations. *Spinal Cord* 2015; 53: 7–13.
- 323 25 Adriaansen JJE, van Asbeck FWA, Tepper M, et al. Bladder-emptying
324 methods, neurogenic lower urinary tract dysfunction and impact on quality
325 of life in people with long-term spinal cord injury. *The Journal of Spinal Cord*
326 *Medicine*. 2017;40(1):43-53.
- 327 26 MEG Leoni, Ruz AED. Management of urinary tract infection in patients
328 with spinal cord injuries Clin Microbiol Infect 2003; 9: 780-785
- 329 27 Di Benedetto P. Clean intermittent self-catheterization in neuro-urology.
330 Eur J Phys Rehabil Med 2011; 47: 651–659.

331