

Original Research Article

**OUTCOME OF CHILDHOOD TUBERCULOSIS AT A SPECIALIST HOSPITAL IN
GUSAU, NIGERIA**

ABSTRACT

Introduction

Tuberculosis (TB) is a chronic infectious disease that is preventable and curable, yet a major cause of childhood morbidity and mortality. Tuberculosis prevalence and mortality are underestimated in many high burden countries. Directly Observed Treatment Short course (DOTS) enhances treatment outcome with overall reduction in morbidity and development of multidrug resistant TB.

Objective

To determine the pattern and outcome of childhood tuberculosis managed at the DOTS clinic in Gusau, Nigeria.

Methodology

A retrospective study of children managed for TB at the DOTS clinic over a 30 months period. The clinic serves both children and adults. All children (≤ 18 years) treated for tuberculosis over the study period were included. Relevant information from the register were retrieved and analysed accordingly. Treatment outcomes were assessed according to World Health Organisation (WHO) and National Tuberculosis and Leprosy Control Programme (NTLCP) guidelines. "Cured" and "treatment completed" outcomes were classified as treatment successful.

Results

Of the 415 patients managed, 76(18.3%) were children; males were 30(39.5.2%), with a M: F ratio of 1:1.5. Mean $\pm$ SD age was 8.89 $\pm$ 5.38 years, with 29(38.2%) being in the 0-5 years age group. Pulmonary TB (PTB) was seen in 58(76.3%), more females had pulmonary TB than males, which was not significant ($\chi^2=1.350$, $p=0.245$). Seventy five (98.7) were new cases, with 1(1.3%) treatment after failure.

Acid fast bacilli (AFB) were positive in 12(15.8%) while GeneXpert MTB/RIF sensitivity was detected in 7(9.2%). Majority 51(67.1%) completed treatment, 12(15.8%) were cured, 9(11.8%) were transferred out, 3(3.9%) died, while 1(1.3%) was lost to follow up; with successful outcome of 82.9%.

Conclusion

Treatment outcome using DOTS strategy was good, with a success rate close to 85.0% of WHO benchmark. The proportion of childhood TB indicates that childhood TB as compared to the adult cases is still under diagnosed and under treated.

Key words- childhood, DOTS, Gusau, tuberculosis, outcome

39 1.0 Introduction

40 Tuberculosis (TB) is a chronic infectious disease that is preventable, treatable and curable;
41 yet still a major cause of childhood morbidity and mortality. In order to decrease transmission
42 of the disease, effective identification, diagnosis and treatment of infectious TB patients is
43 required.[1] A proven strategy to ensure patients' adherence to anti-tuberculous medication is
44 the use of DOTS therapy.[2]

45 DOTS involves mainly early diagnosis by quality ensured sputum-smear microscopy and
46 standardized short-course anti-TB treatment given under direct and supportive
47 observation.[3] Other components are a regular, uninterrupted supply of high quality anti-TB
48 drugs, standardized recording and reporting; and sustained political and financial
49 commitment. It aimed at ensuring that patient with TB complete treatment to cure and
50 prevent the development of drug resistant TB in the community.[3]

51 Evidence has suggested that TB prevalence and mortality are under-estimated in many high
52 burden countries including Nigeria.[4-7] DOTS is associated with significant improved
53 treatment outcome with overall reduction in morbidity and development of multidrug
54 resistant TB.[1]

55 1.1 Objective

56 To determine the pattern and outcome of childhood tuberculosis managed at the DOTS clinic
57 in Gusau, Zamfara State, Nigeria.

58 2.0 Methodology

59 This was a retrospective study of children managed for tuberculosis at the DOTS clinic of
60 Ahmad Sani Yariman Bakura Specialist Hospital (ASYBSH), Gusau, Zamfara State, Nigeria
61 over a 30 months period (1st Jan 2015 to 30th June 2017). All patients that have completed
62 treatment over the study period were included in the study. GeneXpert machine was acquired
63 in the hospital in October 2016, before then only sputum smear microscopy and gastric
64 washout were available for detection of Acid fast bacilli (AFB). Relevant information from
65 the register were retrieved and analysed accordingly. Tuberculosis treatment outcome was
66 assessed according to WHO [8] and NTLCP [9] guidelines.

67 Treatment outcomes: Tuberculosis treatment outcomes were classified as follows:

- 68 1. Cured- sputum smear positive patient who was sputum negative in the last month of
69 treatment and on at least 1 previous occasion.
- 70 2. Treatment completed – Patient who has completed treatment but who does not meet the
71 criteria to be classified as a cure or a failure.
- 72 3. Treatment failure- Any TB patient who is sputum smear positive at 5 months or later
73 during treatment.
- 74 4. Died- Patient who died from any cause during the course of treatment (regardless of the
75 cause of death).
- 76 5. Lost to follow up- Patient whose treatment was interrupted for two consecutive months or
77 more after registration.

78 6. Not evaluated- A TB patient for whom no treatment outcome is assigned (includes case of
79 transferred out to another treatment unit) where the treatment outcome is unknown.

80 7. Transferred out- A TB patient who has been transferred to another local government area
81 to continue his/her treatment and for whom treatment outcome is not known.

82 8. Removed from TB treatment register- Patient who became MTB detected/Rif resistance
83 detected at any point of their treatment and who is moved to 2nd line treatment register.

84 “Cured” and “treatment completed outcomes” were referred to as treatment successful while
85 outcomes such as death, default and treatment failure were considered unsuccessful.

86 3.0 Results

87 Of the 415 patients managed at the DOTS clinic over the study period, 76(18.3%) were
88 children ≤18 years. Males were 30(39.5.2%), with a M:F ratio of 1:1.5. Mean±SD age was
89 8.89±5.38 years (range of 7 months to 17 years). Majority were aged 5 years and below as
90 shown in table 1.

91 Table 1: Frequency distribution of various age groups

Age range (years)	Frequency (Percentage)
0-5	29(38.2)
6-10	17(22.4)
11-15	22(28.9)
16-18	8(10.5)
Total	76(100.0)

92 Pulmonary TB was seen in 58(76.3%) and extra pulmonary TB in 18(23.7%). More females
93 had pulmonary TB than males, which was not statistically significant ($\chi^2=1.350$, $p=0.245$).
94 Seventy five (98.7%) of the patients were new cases, while 1(1.3%) was treatment after
95 failure.

96 Acid fast bacilli were positive in 11(14.5%) while GeneXpert MTB/RIF sensitive was
97 detected in 7(9.2%) and only one (1.3%) was HIV positive as shown in table 2. However,
98 49(64.5%) children did not have GeneXpert done due to lack of availability of the machine.

99 Table 2- Results of investigation carried out on the children

Test	Positive result	Negative result	Not done/Not tested
HIV	1(1.3%)	75(98.7%)	0(0.0%)
AFB smear	11(14.5%)	51(67.1%)	14(18.4)
Gene-Xpert	7(9.2%)	8(10.5%)	61(80.3%)

101 Fifty one (67.1%) had their treatment completed as shown in table 3, with 82.9% successful
102 treatment outcome. All the 3(3.9%) deaths occurred in the 6-10 year age group.

103

104 Table 3- Treatment outcome of the cases

Outcome	Frequency (Percentage)
Treatment completed	51(67.1)
Cured	12(15.8)
Transferred out	09(11.8)
Died	03(3.9)
Lost to follow up	01(1.3)
Removed from register	00(0.0)
Total	76(100.0)

105 4.0 Discussion

106 Directly Observed Treatment Short course have been found to be effective and efficient in
107 treating TB.[1,2] The study showed more females treated than males, which is comparable to
108 what was reported by Panigatti et al[10] but contrasts reports Ramesh et al,[11] Nandimath et
109 al,[12] and Bandichhode et al.[13] This contrasts the Global TB report [4] which showed
110 more males being affected than females, reason for the disparity cannot be explained. The
111 under fives were mostly affected which was similar to other studies done by Ramesh et
112 al,[11] Nandimath et al [12] and Bandichhode et al,[13] however contrasts findings by
113 Adejumo et al.[14] This could be explained by such age group are prone to infections and
114 some of the children may not have been immunised against tuberculosis, however this was
115 not looked for in this study.

116 Majority of our patients had PTB, which is similar to reports from other studies,[11,13,14]
117 but contrasts findings by Panigatti et al.[10] This is because TB cases are primarily
118 pulmonary and is easier to diagnose than extra pulmonary which requires high index of
119 suspicion.

120 Most of the patients were new cases similar to findings by Adejumo et al [14] in Lagos and
121 Daemo et al [15] in Ethiopia. This can be explained by better awareness about the disease and
122 availability of drug treatment with increased presentation to the hospital.

123 The AFB detection rate in this study was low, however it was higher than 5.3% obtained in
124 Kerala [11] and 10.7% in Solapur,[12] but lower than 20.7% reported from Ethiopia.[15]
125 Reason for the low AFB detection rate may be because of rarity of smear positive TB in
126 children and difficulty getting sputum/gastric samples from children. The MTB detection
127 using GeneXpert was also low, and none of the available studies reviewed reported on the
128 yield of GeneXpert in children.

129 Only 1.3% had HIV/TB co infection, in contrast to 15.4% in Ethiopia [15] and 19.5% in
130 Abuja.[16] Even though the study is hospital based, reason for the low prevalence could be

attributable to the fact that Zamfara State has the lowest HIV seroprevalence in the country with 0.9%.[17] However, a previous study in Gusau, Zamfara State by Mado et al [18] reported HIV/TB co infection of 8.0%. The reason for the disparity could be explained by the fact that the previous study was conducted in 2008 when the seroprevalence rate was higher.

The outcome showed majority of the cases completed treatment, with an overall successful outcome close to the 85.0% WHO bench mark. Other studies also showed similar outcome and overall good success rate,[10,12-14,19] however our treatment success rate is higher than 77.4% reported by Adejumo et al[14] and 78.9% by Kebede et al.[19] The mortality rate seen in this study was low, which is similar to 3.5% reported from Ethiopia,[15] but higher than 1.1% reported from India.[12,13] It is however lower than 6.0% reported from Lagos[14] and 28.4% from Kano.[20] The reason for the differences could be due to difference in methodology (some studies were retrospective others prospective, sample size); timing of presentation; presence of complications and other associated medical problems.

4.1 Conclusion

Treatment outcome of children treated for TB in our centre was good, with a success rate close to WHO benchmark of 85.0%. The proportion of 18.3% childhood TB as compared to the adult cases indicates that childhood TB is still under diagnosed and under treated.

4.2 Limitations

The study was retrospective, hence timing of presentation and presence of complications were not looked at.

5.0 References

1. Sunday O, Oladimeji O, Ebenezer F, Akintunde B, Abiola TO, Saliu A *et al.* Treatment outcome of tuberculosis patients registered at DOTS centre in Ogbomosho, Southwestern Nigeria: A 4- year retrospective study. *Tuberculosis Research Treatment* 2014. doi.10.1155/2014201705.
2. Ebuenyi ID, Ikuabe PO, Jumbo J. Treatment outcome of tuberculosis at one year: A single centre's experience in Niger Delta, Nigeria. *IJTDH* 2016;1291):1-6.
3. WHO. What is DOTS (Directly Observed Treatment, Short Course). In WHO Tuberculosis. www.searo.who.int. Accessed 9/01/2018.
4. WHO. Global Tuberculosis Report 2016. ISBN 978 92 415 6539 4. Accessed 23/10/2017.
5. Joseph BN, Sariem CN, Dangiwa DA, David S, Joseph SI, Egoh DZ. Comparative assessment of Directly Observed Treatment Short course (DOTS) for tuberculosis in a primary and a tertiary health centre in Nigeria. *J Pharm Bioresources* 2015;12(1):22-9. doi.10.4314/jpb.v12:1.4.
6. Erhabor GE, Adewole O, Adisa AO, Olajolo OA. Directly observed treatment short course therapy for tuberculosis- a preliminary report of a three-year experience in a Teaching Hospital. *J Natl Med Assoc* 2003;95:1082-8.
7. Ogbudebe CL, Izuogu S, Abu CE. Magnitude and treatment outcomes of pulmonary tuberculosis patients in a poor urban slum of Abia State, Nigeria. *Int J Mycobacter* 2016;5(2):205-10.

- 172 8. World Health Organization (2013b) Definitions and reporting framework for
173 tuberculosis – 2013 revision 2013. WHO Geneva. Accessed 11/11/2017.
- 174 9. Federal Ministry of Health, Nigeria (2010) National Tuberculosis and Leprosy
175 Control Programme: Workers Manual. Revised 5th edition. Abuja: Federal Ministry
176 of Health. 1-119. Accessed 11/11/2017.
- 177 10. Panigatti P, Ratageri VH, Shivannard I, Madhu PK, Shepur TA. Profile and outcome
178 of childhood tuberculosis treatment with DOTS – An observational study. *Ind J Paed*
179 2014,81(1)9-14.
- 180 11. Ramesh H, Shoba A, Manjula VD. Treatment outcomes of childhood tuberculosis
181 with DOTS strategy in Kottayam, Kerala. *Ind J Comm Hlth* 2012,24(4):280-4.
- 182 12. Nandimath VA, Ukarande AR, Chaithra S. Outcome of paediatric DOTS treatment at
183 city tuberculosis centre, Solapur. *Ind J Curr Research* 2015,7(08):19683-5.
- 184 13. Bandichhode ST, Nandimath VA. Health profile of paediatric tuberculosis patients on
185 directly observed short course therapy. *Ind J Contemp Pediatr* 2016,3(4):1401-4.
- 186 14. Adejumo OA, Daniel OJ, Adebayo BI, Adejumo EN, Jaiyesimi EO, Akang G.
187 Treatment outcomes of childhood tuberculosis in Lagos, Nigeria. *J Trop Pediatr*
188 2016,62(2):131-8.
- 189 15. Daemo MD, Kelbore AG. Treatment outcomes and associated factors of childhood
190 tuberculosis: treated under DOTS program in health centers of Makelle town, Tigray
191 Regional State, Ethiopia. *Central Afr J Pub Hlth* 2016,2(1):11-7.
- 192 16. Okechukwu AA, Okechukwu OI. Clinical correlate of tuberculosis in HIV co infected
193 children at the University of Abuja Teaching Hospital, Gwagwalada, Nigeria. *Niger J*
194 *Clin Pract* 2011,14(2):206-11.
- 195 17. NACA Annual report 2016. www.naca.gov.ng. Accessed 6/11/2017.
- 196 18. Mado SM, Isa A, Abubakar U, Onazi SO, Gbemiga AO. Spectrum of tuberculosis in
197 children at Federal Medical Centre, Gusau, Zamfara State, North-western Nigeria.
198 *Sahel Med J* 2017, 20:8-12.
- 199 19. Kebede ZT, Taye BW, Matebe YH. Childhood tuberculosis; management and
200 treatment outcome among children in Northwest Ethiopia: a cross-sectional study.
201 *Pan Afr Med J* 2017,27:25.
- 202 20. Adamu AZ, Aliyu MH, Galadanci NA, Musa BM, Gadanya MA, Gajida AU *et al.*
203 Deaths during tuberculosis treatment among paediatric patients in a large tertiary
204 hospital in Nigeria. *PLoS* 2017,12(8):e0183270. doi.10.1371/journal.pone.0183270.