

MANUSSCRIPT NUMBER: 2017/AJMAH/34160

Bacteriology of Suya Meat Sold in Bonny Local Government Area, Rivers State, Nigeria.

Konne, Felix Eedee^{1*}, Wokem, Gloria Ngozika¹, Tetam, Jack Gbenene¹,
Monsi, Tombari Pius¹.

¹Department of Medical Laboratory Science, Rivers State University, Port Harcourt, Nkpolu -
Oroworukwo P.M.B. 5080, Rivers State, Nigeria

Original Research Article

ABSTRACT

Aim: To determine the bacteriology contamination of roasted Suya meat sold in Bonny local government area.

Place and Duration of study: The locations were, Tennis Hut, Golf Club, Ama Hausa, Hospital Road, (round about), Finima market junction and Akiama between November and December, 2016.

Materials and Methods: Total of five (5) different Suya meat sample were collected at random, from each of these six Suya location or stand in Bonny, which sum up the number to total of 30 Suya meat samples assayed. Culture characteristics of colonies were considered after 24hrs incubation period. Identifications of isolates were carried out using standard methods.

Results: The results shows that six genera of bacteria which include *Bacillus cereus*, *Salmonella species*, *Staphylococcus aureus*, *Klebsiella species*, *Enterococcus species* and *Proteus species* was isolated from the Suya meat. The total number of *Bacillus cereus* and *Salmonella species* presence were 10 (34%) and 5(17%) respectively, from the five Suya samples each, across the six Suya stand in Bonny. *Escherichia coli* have no growth. The total of 1(3.4%) and 2 (6.8%) of *Staphylococcus aureus* and *Klebsiella species* respectively were also isolated. Total value of 6(20%) and 5(17%) of *Enterococcus species* and *Proteus species* respectively were also isolated. However, the results also shows that Tennis Hut and Golf club have total of 3(9.6%) and 2(6.4%), of different types of microorganisms isolated. Finima Suya and Akiama have total of 7(22%) and 6(19%) respectively of different type of microorganisms isolation.

Conclusion: The study revealed that proper preparation and processing of our Suya meat should be adhered to by law.

Keywords: Bacteriology; Suya; meat; Bonny.

1. INTRODUCTION

Suya Meat is obtained from roasted or smoked beef and other boneless animal mostly. They are good, taste and rich in protein, vitamin B12, zinc, selenium and phosphorus. It is also rich in niacin, iron and vitamin (pyridoxine). Suya meat is commonly found in most streets as source of food in Nigeria and sub-Sahara Africa [1].

Based on that Suya meat contamination is associated with the exposure on the street. It is easily contaminated with bacteria, microorganisms and parasite. The meat (Suya meat) is produced from meat part from animals. The meat part from this animal is roasted smoked to dry making it attractive to eat and increasing the shelf life of the Suya meat. Therefore, it is necessary to carried out some microbiological quality assessment in order to ascertain their safety for consumption. Suya could be contaminated from air, handlers, equipments, water, papers, some kitchen utensils for example the spoon, trays, knives and sometimes the contamination could be due to nature of the vendors and the environments [4].

In the abattoirs during slaughtering, there is the chance of contaminating sterile tissue of the meat, with some intestinal flora which is associated with humans, animals and the environment [6].

It is believed that raw consumption of uncooked meat could result to food poisoning. This is possible based on the presence of some bacteria in the meat which has the ability to produce some toxins in human resulting to some pathological condition like diarrhea and vomiting. However, the infectious doses of some common contaminations of meat were still under debate or investigations [8].

Based on the fact that microorganisms could be found everywhere, it's believed that unhygienic preparation of Suya meat, could promote the presence of microorganism on Suya meat sold in Bonny, which could cause contamination to the consumer. It is based on these; my research will ascertain

some possibilities towards recommending ways to reduced possible health hazards that could be associated with the consumption of Suya meat sold in Bonny.

This work is aimed at determining the bacteria contamination of roasted Suya meat sold in Bonny local government area.

2.0 MATERIALS AND METHODS

2.1 Study area

This study was conducted within the City of Bonny (Ibani) Local Government Area of Rivers state in the month of November to December, 2016. Bonny is geographically located at latitude $40^{\circ} 26' N$ $7^{\circ} 10' E$, with lengthy raining season and short dry season temperature between $25-28^{\circ}C$. It is situated at the southern edge of Rivers State in the Niger Delta of Nigeria near Port Harcourt. The major occupation of the people is fishing and farming and they speak Ibani language.

The population for this study was drawn from six (6) selected Suya stands namely; Tennis Hut, Golf Club, Finima market Junction, Akiama, Ama-Hausa and Hospital Road (roundabout junction). A total of thirty (30) samples were collected for the study, five from each Suya stand location at different time and day.

2.2 Sample Collection

All Suya meat samples from each was aseptically collected into a sterile container, labeled and taking to Bonny General Hospital Medical Laboratory section and department of Medical laboratory Science, Rivers state University, Port Harcourt to analyzed and was cultured into various culture media, strictly following the protocol for media inoculation and organism identification guide for bacteriology culture. The materials used include: Sterile universal containers, Bijou bottles, sterile disposable wire loops, Petri-dishes, Media, Peptone water, Incubator, Autoclave and Test tubes.

The media was prepared according to the manufacturer's instruction. Based on the number of plates, sterilized by autoclaving at $121^{\circ}C$ for 15 minutes and dispensed aseptically into disposable Petri- dishes (about 20mls each) after cooling to about $50^{\circ}C$ [2].

2.3 Bacteriology Determination

The sample was incubated aerobically in the incubator for 24hours. Sub-culture was done the following day into various media. The plates were incubated at $37^{\circ}C$ for 12 -24 hours and examined for bacterial growth, biochemical and gram staining test for proper identifications was also considered [2].

2.4 BIOCHEMICAL TESTS

Number of Isolates	1	2	3	4	5	6
Morphology	Short rods	Short rods	Short rods	Cocci clusters in	Short rods	Short rods
Maltose	+	+	+	+	+	+
Glucose	+	+	+	+	+	+
Gram Staining	-	-	-	+	-	-

Indole	+	-	-	-	-	-
Citrate	-	-	+	-	-	-
Oxidase	+	+	-	-	-	-
Catalase	+	+	+	+	-	+
Coagulase	-	-	-	+	-	-
Urease	+	+	-	-	+	-
Bacteria species	<i>Proteus</i> spp	<i>Enterococcus</i> spp	<i>Klebsiella</i> spp	<i>S. aureus</i>	<i>Salmonella</i> spp	<i>Bacillus cereus</i>

Key: (-) Negative, (+) Positive

2.5 Data Analysis

The data were analyzed statistically using Excel, Software for the graph and the percentage calculations.



3. RESULTS

Result shows that from the 30 (thirty) samples of Suya meat brought from six different locations in Bonny Local Government Areas of Rivers state, were all positive with different microbial contaminations and distribution. The presence of *Bacillus cereus*, *Salmonella species*, *Staphylococcus aureus*, *Klebsiella species*, *Enterococcus species* and *Proteus species* that were isolated across the six Suya stands are 10(34%), 5(17%), 1(3.4%), 2(6.8%), 6(20%) and 5(17%) respectively, for **Tennis Hut**, **Golf Club**, **Finima**, **Akiama**, **Ama hausa** and **Hospital Road**.

Table. 1: Showing the Total Number of Microorganisms Isolated from Six different Suya Locations in Bonny Local Government Area.

Organisms Isolated	Tennis Hut (%)	Golf Club (%)	Finima (%)	Akiama (%)	Ama hausa (%)	Hospital Road (%)	Total (%)
<i>Bacillus cereus</i>	2(66%)	1(50%)	3(42%)	2(33%)	2(25%)	2(40%)	10(34%)
<i>Salmonella species</i>	1(33%)	00(00)	2(28%)	00(00)	2(25%)	00(00)	5(17%)
<i>Escherichia coli</i>	00(00)	00(00)	00(00)	00(00)	00(00)	00(00)	00(00)
<i>Staphylococcus aureus</i>	00(00)	00(00)	00(00)	1(16%)	00(00)	00(00)	1(3.4%)
<i>Klebsiella species</i>	00(00)	00(00)	00(00)	1(16%)	00(00)	1(20%)	2(6.8%)
<i>Enterococcus species</i>	00(00)	1(50%)	00(00)	1(16%)	2(25%)	2(40%)	6(20%)
<i>Proteus species</i>	00(00)	00(00)	2(28%)	1(16%)	2(25%)	00(00)	5(17%)
Total	3(9.6%)	2(6.4%)	7(22%)	6(19%)	8(25%)	5(16%)	

Percentage of Microorganisms Isolated from the Six different Locations; Tennis Hut, Golf Club, Finima, Akiama, Ama hausa and Hospital Road









Figure 1. Graph showing the percentage of organisms isolated from: a) Tennis Hut, b) Golf Club, c) Finima, d) Akiama, e) Ama Hausa, f) Hospital Road





Figure 2. a) Percentage a) organisms isolated from areas, b) Different microorganisms isolated.

1 4.0 DISCUSSIONS

2 *Bacillus cereus* and few other microorganisms isolated across the six Suya stands agreed with
3 other work earlier done [5], and the improper establishment of a Hazard Analysis Critical Control
4 Point (HACCP), [10] which is also essential for the control of *Salmonella species*, *Enterococcus*
5 *species* and *Proteus species*. This method could be used to achieved the potential food safety
6 hazard or could be taken to reduced or totally eliminates the presence of these organisms
7 contained in the meat, contaminated from the abattoir personnel or the environment [9].
8 The Golf Club Suya sample assayed indicates that 1(50%) was contaminated with *Bacillus*
9 *cereus* and 1(50%) of *Enterococcus species*, given a total of 2(6.4%) of different microorganisms
10 of Suya contamination in Bonny. It is believed that *Bacillus cereus* can cause two types of
11 illness, one characterized by diarrhea and vomiting with the aid of their enterotoxins. Though the
12 diarrhea enterotoxin is heat sensitive (could be inactivated by 56°C heat for 5 minutes). But the
13 emetic toxin which induces vomiting is very heat resistant [9]. *Bacillus cereus*, a gram positive,
14 facultative aerobic, spore forming rod which are different from other *Bacillus species* in their
15 biochemical test with their optimum growth temperature of about 30°C to 50°C, pH value of 4.3 to
16 9.3 and water activity (AW) values of 0.912. *Bacillus cereus* is commonly found in nature for
17 example in the soil and water. The infective dose of *B. cereus* could be more than 10 cfu (gm) of
18 visible cells consumed to cause disease [9].
19 The result gotten from Finima, shows that 3(42%) of *Bacillus cereus*, 2(28%) of *Salmonella*
20 *species* and 2(28%) of *Proteus species* were positives given a total of 7(22%) of different types
21 of microbial contaminations. *Salmonella species* could be gotten from uncooked cow meat, or
22 taking raw cow meat that is improper cooked could be harmful to human health [7]. The result
23 from others [5] shows that the isolation of *Salmonella species* from Suya meat sample also
24 agreed with the fact that 2(28%) of *Salmonella species* contamination can rendered the Suya
25 meat unsafe for consumption [5].
26 Akiama, also shows 2(33%) of *B. cereus*, *Staphylococcus aureus* 1(16%), *Klebsiella species*
27 1(16%), *Enterococcus species* 1(16%) and *Proteus species* 1(16%) given a total number of
28 6(19%) different microorganisms, confirming previous work [5]. The presence of *Staphylococcus*
29 *aureus* in Suya meat from Akiama indicates *Staphylococcus aureus*, gram positive coccal bacteria,
30 has high salt tolerance which enable its growth in Suya meat. It also can produce toxins which
31 are heat resistant and cannot be destroyed through cooking. It is commonly contracted through
32 food workers. An individual with weakened immune systems may also be vulnerable to it toxins
33 [8]. Because *Staphylococcus aureus* are commonly found on the skin, hair, nose and human
34 throats, the bacteria could be found in up to 25% of healthy individual and are more common
35 with those individual that has nose, skin, eye and throat infections [7]. Ama Hausa results show
36 contamination of Suya meat of 2(25%) *Bacillus cereus*, *Salmonella species* 2(25%),
37 *Enterococcus species* 2(25%) and *Proteus species* 2(25%) given a total of 8(25%) of different
38 microorganisms contamination in Bonny.
39 Finally, Suya meat picked from Hospital road shows that *Bacillus cereus* has 2(40%), *Klebsiella*
40 *species* shows 1(20%) and *Enterococcus species* 2(40%) positive out of the total of 5(16%) of
41 different microorganisms in Suya meat from Hospital road in Bonny Local Government area. The
42 presence of *Proteus species* agreed with the fact that, it is commonly found in the human
43 intestinal tract as part of normal flora, and it is also found in multiple environmental habitats, it is
44 a gram-negative bacillus which also colonized both the skin and oral mucosa. *Bacillus cereus*
45 with total of 10(34%) was isolated across the six Suya stand in Bonny making it the highest
46 among other microorganism found from the Suya sample. This agreed [10], that *B. cereus* has
47 the ability to produce spores which can also germinate if the temperature is lowed, make it to be
48 temperature tolerance.
49 *Salmonella species* and *Staphylococcus aureus* with the total of 5(17%) and 1(3.4%)
50 respectively indicates that both organisms could also cause poison to human when consumed in
51 their high infective dose. This also agreed [3], with the saying that their infective dose could be
52 as little as 1.0 nanogram of the toxin per gram of food or meat to produce illness. Ingestion of 1 -

5 microgram of the toxin has been shown to be associated with many outbreaks of diseases [3]. *Klebsiella species*, *Enterococcus species* and *Proteus species* have total of 2(6.8%), 6(20%) and 5(17%) respectively isolated from the six different Suya stand in Bonny local government area. But the infectious dose of *Klebsiella species* is still unknown and it is found widely through nature in soil and water so there is that likelihood that Suya contaminated with *Klebsiella species* could be from soil or water source, human skin and mouth [3]. Finally, *Escherichia coli* were negative across the six Suya meat stand in Bonny, with the use of Chromagen 0157 media for isolation and differentiation and there was no growth noticed.

4. CONCLUSION

In a way to reduce or total elimination *Bacillus cereus* organism which also have contributed to some resistant noticed in the used of our antibiotics, the authority appointed to over see the proper preparation of our abattoir meats and Suya processing should adhered to required principle by law. Also the need to stored perishable food like the Suya meat within the temperature “danger zone” at the range of 50°C to 60°C should be encouraged. Suya meat should be reheated to about 75°C before eating. Studies have also shown that food handles especially suya meat handless can also serve as a source of infection and the hand contact with the suya represent the means by which the contamination may occur.

4. RECOMMENDATIONS

Hazard Analysis and Critical Control Point (HACCP) which represents a systematic preventive approach to food safety should be encouraged. Effective measures should be in place in order to avoid cross contamination between raw and cooked foods. Regular hand washing practice with soap and a running tap water and lowering of finger nails thoroughly before cooking and serving the public with Suya meat should be encouraged. Employee with wounds or skin infection are strictly prohibit from Suya handling in Bonny local government areas. Routine environmental sanitations practice, educational symposiums and disinfection of equipment used in Suya meat processing in Bonny should be encouraged.

COMPETING INTERESTS

Authors have declared no competing interests.

REFERENCES

1. Abdullahi, O. I., Umoh, V. J. & Gaadima, M. Hazards associated with kilishi preparation in Zaria, Nigeria. *Nigeria Journal of Microbiology*, 2004, 8(2), 337 – 345.
2. Cheesbrough, M. *District Laboratory Practice in Tropical Countries Part 2*. Cambridge University Press, United Kingdom. 2000, Pp. 62 - 70.
3. Diana, G., Susana, D., Daniel, V., Beatrice, M., Marta, L. C., Ana, R., Juan, J. H. & Pilar, G. Incidence of *Staphylococcus aureus* and analysis of associated bacterial communities on food industry surfaces. *Applied and Environmental Microbiology*, 2012, 78(24), 8547 – 8554.
4. Edema, M. O., Osho, A. T. & Dials, C. I. Evaluation of microbial hazard associated with processing of Suya (a grilled meat product). *Science Research Essays*, 2008, 3(12), 621 – 626.

- 98 5. Egbebi, A.O & Seidu, K.T. Microbiological evaluation of Suya (dried smoked meat)
99 in Ado and Akure, south West Nigeria. *European Journal of Experimental biology*, 2011,1(4),
100 1 – 5.
- 101 6. Judge, D. M., Robert, A. M. & Morris, M. J. Preparation of Suya in Africa. *Journal of*
102 *African Foods*, 2002, 20, 52 – 55.
- 103 7. Ogbonna, I. O., Sunday, D. M. Oyekemi, A. & Odu, C. E. Microbiological safety and
104 proximate composition of (suya) stored at ambient temperature for six hours from Maidugari,
105 Northern Nigeria. *Internet Journal for Food Safety*, 2012, 14, 11 – 16.
- 106 8. Riser, E. T. & Marth, E. H. New foodborne pathogens of public health significance.
107 *American Journal of Dietary Association*, 1989, 89, 948 – 956.
- 108 9. Sonal, A. & Chagan, T. The bacteria with the lowest infections base to cause meat
109 food poisoning. *Journal of Interdisciplinary Science Topics*, 2016, 21, 13 – 20.
- 110 10. Sperder, W. H. & Richard, F. S. Happy 50th birthday to HACCP. Retrospective and
111 perspective. *Food safety Magazines*, 2010, 20, 42 – 46.
112

113
114