

Microalbuminuria As A Marker Of Cardiovascular Risks In Egyptian Hypertensive Patients

Abstract:

Hypertension (HTN) is one of the modifiable atrial fibrillation (AF) risk factors, and management of HTN may reduce the incidence of AF. Microalbuminuria (MAU) increases cardiovascular risk in hypertension.

Aim of the study: To determine the association of microalbuminuria in Egyptian hypertensive patients and its relationship with AF

Patients and methods: Five hundred hypertensive patients without a history of pre-existing kidney diseases participated in this study. A questionnaire was used for collecting information on demographics, lifestyle, and family history of cardiovascular and kidney disease, and spot morning urine samples were collected for albuminuria estimation.

Results: A total of 500 Egyptian hypertensive patients, aged 47 ± 7.3 years were enrolled in this study. Two hundred and ninety three were males and 107 were females. The mean duration of hypertension was 5.6 ± 2.7 years. Mean body mass index (BMI) was 27.1 ± 2.4 kg/m². There were 354 (70.8%) patients whose BP well controlled ($<140/<90$ mmHg), while 146 (29.2%) patients were not controlled. 249 (49.8%) patients had MAU. Logistic regression analysis revealed that male gender, lower glomerular filtration rates, higher mean BP, higher heart rates and ejection fraction less than 50% had significant effect on prevalence of MAU with an odds ratio > 1 and P value < 0.05 . Two hundred patients had AF, 173 (86.5%) of them had MAU. There was a positive correlation between MAU and AF and left ventricular hypertrophy (LVH) with P value < 0.05 .

Conclusion and recommendation: Screening for albuminuria may be useful in early risk assessment of cardiovascular disease in Egyptian hypertensive patients, identification of a patient at risk of CV events provides an opportunity for early treatment, to slow the progression of disease.

key words: Hypertension, AF, CV risks, micro albuminuria.

Introduction:

Hypertension is still the main risk factor for the development of atrial fibrillation (AF). Several pathophysiologic mechanisms (such as structural changes, neurohormonal activation, fibrosis, atherosclerosis, etc.) have been advocated to explain the onset of atrial fibrillation (1). AF is often hidden [paroxysmal], and is associated with an about fivefold risk of stroke. This risk

can be reduced by the use of appropriate therapy for hypertensive patients with increased cardiovascular risk(2).

Hypertension and microalbuminuria (MAU) commonly coexist. The mechanism is controversial but is thought to be a renal manifestation of generalised vascular endothelial dysfunction(3). Microalbuminuria is a well-recognised marker of cardiovascular complications and increased cardiovascular risk in hypertension(4).

However, reliable data about the association of microalbuminuria in hypertensive patients and its relation with cardiovascular and renal morbidity are limited in Egypt. There is still a debate concerning whether screening for microalbuminuria is cost-effective in risk assessment and prevention of cardiovascular disease, so **the aim of this study** is to determine the association of microalbuminuria in Egyptian hypertensive patients and its relationship with AF.

Patients and methods:

A cross-sectional study was performed from January to December 2016 at internal medicine outpatient clinic of El Fayoum University Hospital, on five hundred Egyptian hypertensive non diabetic patients. Subjects were excluded from the study if they had co-morbidities like cardiomyopathy, valvular heart diseases, diabetes mellitus(DM), pulmonary diseases, liver diseases, renal impairment, proteinuria, urinary tract infection (UTI), anaemia or thyrotoxicosis. All cases of secondary hypertension were excluded. All participants gave written informed consent. This study was approved by our ethical committee.

A structured questionnaire regarding the demographic data such as age, gender, height, body weight and family history of DM. BMI was calculated as weight (Kg) divided by height (m²). Blood pressure was measured with a suitable mercury sphygmomanometer. Readings were taken after the patients had been resting comfortably, back supported in the sitting or supine position, for at least 5 minutes and at least 30 minutes after smoking or coffee ingestion. Mean arterial pressure (MAP) was calculated as [systolic + (2 x diastolic) pressure]/3. High blood pressure was defined as systolic blood pressure (SBP) ≥ 140 mmhg and/ or diastolic blood pressure (DBP) of ≥ 90 mmhg or use of anti-hypertensive medications. Heart rate, cardiac examination, electrocardiogram (ECG) and echocardiography (ECHO) were done. Lipid profile, liver profile, fasting and 2 hours post prandial blood glucose, HA1C, renal function tests (glomerular filtration rate, serum creatinine), Estimated glomerular filtration rate (GFR) by the Modification of Diet in Renal Diseases (MDRD):

$GFR = 186 \times \text{serum CR}^{-1.154} \times \text{age} \times 1.212 (\text{if patient is black}) \times 0.742 (\text{if female})$. Morning mid-stream urine sample, negative for proteinuria by Albustix, was used to calculate microalbumin: creatinine ratio in mg/g. Microalbumin was carried out using ELISA assay. If the ratio was < 30 mg/g the patient was considered normo-albuminuric, ratios between 30-300mg/g were indicative of microalbuminuria and above 300mg/g revealed macro-albuminuria(3).

89

90 **Statistical analysis:**

91 All the collected data were expressed as mean $\pm$ SD and analyzed by using
 92 SPSS version 20 using the following tests: Student T, Chi square, Pearson
 93 correlation coefficient and logistic regression. $P > 0.05$ was considered non-
 94 significant, $P < 0.05$ was considered significant $P < 0.01$ was considered highly
 95 significant.

96 **Results:**

97 A total of 500 hypertensive patients, aged 47 ± 7.3 years were enrolled in this
 98 study. Two hundred and ninety three were males and 207 were females. The
 99 mean duration of hypertension was 5.6 ± 2.7 years. Mean body mass index
 100 (BMI) was 27.1 ± 2.4 kg/m². Beta blockers (BBs) were prescribed in 50.9%,
 101 followed by calcium channel blockers (CCBs) 23.5 %, Angiotensin converting
 102 enzyme inhibitors (ACEI) (21.1%) and thiazide type diuretic (4.5%). There
 103 were 354 (70.8%) patients whose BP well controlled ($<140/<90$ mmHg), while
 104 146 (29.2%) patients were not controlled. 249 (49.8%) patients had MAU.
 105 Logistic regression analysis revealed that male gender, lower glomerular fil-
 106 tration rates, higher mean BP, higher heart rates and ejection fraction (EF)
 107 less than 50% had significant effect on prevalence of MAU with an odds ratio
 108 > 1 and P value < 0.05 . Two hundred patients had AF (173 (86.5%) of them
 109 had MAU. There was a positive correlation between MAU and AF and LVH
 110 (diagnosed by ECG) in essential hypertension patients, there was no correla-
 111 tion between renal function and lipid profile, BBs, diuretics, as shown in ta-
 112 bles 1 and 2

113 **Table (1): Correlation between microalbuminuria and different parameters**

114

PARAMETER	P VALUE	PARAMETER	P VALUE
Age	<0.05	AF	<0.05
Male	<0.05	HR	<0.05
Female	>0.05	LVH	<0.05
BMI	<0.05	EF	<0.05
Duration of HTN	<0.05	GFR	<0.05
Control on HTN	<0.05	DIURETICS	>0.05
ACEI	<0.05	BB	>0.05
CCB	<0.05	lipid profile	>0.05

Table (2): Multivariate logistic regression in all patients and odds ratio (OR) of different predictors for presence of microalbuminuria.

Predictor	OR	95 CI%	P value
Age (middle vs young)	1.030	0.931–1.427	0.762
Gender (male vs female)	1.220	1.170-1.250	0.04
BMI (>25 vs <25)	1.066	1.021-1.322	0.235
GFR (<30 vs > 80)	1.598	1.498-1.683	0.023
GFR (30-60 vs > 80)	1.345	1.294-1.413	0.049
GFR (60-80 vs > 80)	1.531	1.481-1.601	0.034
Duration of hypertension	1.030	1.011-1.463	0.247
Mean blood pressure	1.395	1.298-1.410	0.037
HR (>100 vs < 80)	1.154	1.054-1.236	<0.001
HR (80-100 vs < 80)	1.396	1.376-1.412	0.028
EF (>50% vs <50%)	1.002	0.981-1.031	<0.001

Discussion

In this study, 249 (49.8%) patients had MAU, increase BMI, male gender, lower glomerular filtration rates, higher mean BP, higher heart rates and ejection fraction less than 50% had significant effect on prevalence of MAU with an odds ratio > 1 and P value < 0.05 .

Prevalence of microalbuminuria in hypertensive patients is ranging in the different countries between 53% and 71%, with the highest rate in patients with uncontrolled hypertension(5). This variation might be due to patient characteristics.

Microalbuminuria is independently associated with numerous modifiable CV risk factors and markers of CV diseases. These include: obesity; smoking; insulin resistance syndrome; left ventricular hypertrophy (LVH); left ventricular (LV) dysfunction and elevated C reactive protein levels(6,7).

Obesity, by several mechanisms (glomerular hyper-filtration, focal segmental glomerulosclerosis, insulin resistance and adiponectin) can lead to the development of albuminuria(8).

We found a positive correlation between MAU and AF and LVH in essential hypertension patients, there was no correlation between renal function and lipid profile, BBs, diuretics.

Many studies (9) have shown an association between microalbuminuria and risk of CV events. Microalbuminuria was shown to predict ischaemic heart disease(9).

Moreover **Wachtell et al.**, (10) reported that microalbuminuria is associated with persistent electrocardiographic (ECG) LVH independent of established risk factors for cardiac hypertrophy , multivariate analysis showed that LVH was associated with a 1.6-fold higher prevalence of microalbuminuria and 2.6-fold higher prevalence of macroalbuminuria (both p<0.001). This relationship was independent of age, systolic and diastolic blood pressure (SBP and DBP), diabetes, gender, race, serum creatinine or smoking status, and, there-

fore, it has been postulated that cardiac damage and albuminuria occur in parallel (10).

The main determinant of albumin excretion rate in subjects with mild hypertension and no cardiovascular complications seems to be the hemodynamic load, whereas in subjects with more severe hypertension and associated target organ damage, the augmented urinary albumin leak is probably the consequence of glomerular damage. Further evidence for the link between microalbuminuria and CV disease comes from the fact that microalbuminuria is considered a reliable indicator of endothelial dysfunction, this being associated with transendothelial escape rate of albumin and plasma levels of von Willebrand factor(11). Endothelial dysfunction is considered an early step of atherosclerosis and represents increased risk for CV events(12).

Conclusion and recommendation: Screening for albuminuria may be useful in early risk assessment of cardiovascular disease in Egyptian hypertensive patients. Early identification of a patient at risk of CV events provides an opportunity for early treatment, to slow the progression of disease.

References

1. Manolis AJ, Rosei EA, Coca A, et al. Hypertension and atrial fibrillation: diagnostic approach, prevention and treatment. Position paper of the Working Group 'Hypertension Arrhythmias and Thrombosis' of the European Society of Hypertension. *J Hypertens*. 2012;30(2):239-52.
2. Wayne Kuznar. Microalbuminuria, Retinopathy Predict AF November 22, 2013.
3. Alia Ali, Azeem Taj, Muhammad Joher Amin, et al. Correlation between Micro albuminuria and Hypertension in Type 2 Diabetic Patients *Pak J Med Sci*. 2014; 30(3): 511–514.
4. Palatini P1. Micro albuminuria in hypertension. *Curr Hypertens Rep*. 2003;5(3):208-14.
5. Böhm M, Thoenes M, Danchin N, et al. Association of cardiovascular risk factors with microalbuminuria in hypertensive individuals: the i-SEARCH global study. *J Hypertens* 2007; 25: 2317-24.
6. Barzilay JI, Peterson D, Cushman M et al. The relationship of cardiovascular risk factors to microalbuminuria in older adults with or without diabetes mellitus or hypertension: the cardiovascular health study. *Am J Kidney Dis* 2004; 44: 25-34.
7. Pongsathorn Gojaseni, Angkana Phaopha, Worawon Chailimpamontree, et al. Prevalence and risk factors of microalbuminuria in Thai nondiabetic hypertensive patients. *Vasc Health Risk Manag*. 2010; 6: 157–165.
8. Farzanehsadat Minoo, Mitra Mahdavi-Mazdeh, Mohamad-Reza Abbasi, et al. Impact of the severity of obesity on microalbuminuria in obese normotensive nondiabetic individuals *J Renal Inj Prev*. 2015; 4(2): 34–38.

- 194 9. Karalliedde J and Viberti G. Microalbuminuria and cardiovascular risk.
195 Am J Hypertens. 2014 ;17(10):986-93.
- 196 10. Wachtell K, Olsen MH, Dahlof B, et al. Microalbuminuria in hypertensive
197 patients with electrocardiographic left ventricular hypertrophy: the LIFE
198 study. J Hypertens 2002; 20: 405-12.
- 199 11. Deckert T, Feldt-Rasmussen B, Borch-Johnsen K et al. Albuminuria re-
200 flects widespread vascular damage. The Steno hypothesis. Diabetologia
201 2015; 32: 219-26.
- 202 12. Quyyumi AA. Prognostic value of endothelial function. Am J Cardiol
203 2013; 91:19H-24.