

1 ***Proximate composition and nutritional analysis of seeds and testas of Dacryodes***
2 ***edulis and Garcinia kola***

3
4 **ABSTRACT**

5 *Dacryodes eludis* and *Garcinia kola* are two plants amongst others that are rightly
6 regarded as underutilized. The aim of this study was therefore to examine the seeds
7 and testas of *G. kola* and *D. eludis* for the presence of nutrients, minerals, vitamins and
8 anti-nutrients. The analyses were done using standard techniques and the resulting
9 replicate readings subjected to analysis of variance (ANOVA). The results of the
10 proximate composition revealed that both seeds and testas of the studied plants were
11 very rich in nutrients. The moisture content was more in the testas than the seeds of
12 both plants but was highest in the testa of *D. eludis* (41.00 ± 0.01 g/100g dry matter). The
13 ash and protein contents were almost similar in both plants. Fat and fibre contents were
14 higher in the *G. kola* while the testa of *D. eludis* had more carbohydrate (71.88 ± 0.02
15 g/100g dry matter). Consistently, *G. kola* had more vitamins than *D. eludis*. However,
16 the most abundant vitamin was B which ranged from 213.07 to 224.70 (mg/dl) while the
17 least in both plants was vitamin E that ranged from 2.08 to 3.14 (mg/100g). Mineral
18 analysis showed the presence of minerals such as Na, K, Ca, Mg, Fe, Zn, Cu and P in
19 *G. kola* and K, Ca, Mg, P and N for *D. eludis*. Anti-nutrients analysis revealed the
20 presence of hydrocyanic acid, soluble and total oxalate, and phytate with *G. kola* having
21 the highest concentrations of all the anti-nutrients. Analysis of replicate readings for all
22 examined parameters showed significance ($p < 0.05$). There is a need to further studies
23 aimed at their utilization in human and livestock nutrition.

24 ***Keys words: Garcinia kola, Dacryodes eludis, anti-nutrients, vitamins, minerals,***
25

26 **INTRODUCTION**

27 Plants not only form the integral part of the world's food chains, they are a source of
28 medicine, raw material, food, and so on for man and animals. Most of the plants parts
29 such as fruits, seeds and vegetables, though very rich in amino acids, minerals and
30 vitamins have limited use in human and livestock nutrition because of unknown levels of
31 anti-nutrients they may contain. *Dacryodes eludis* and by extension *Garcinia kola* are

largely regarded as underutilized tree crops despite the fact that they are rich in nutrients and there is a renewed interest in their medicinal potentials across the world [1].

African pear (*Dacryodes eludis*) is a tree crop that is largely cultivated mainly for its oily fruit as it contains a lot of amino acids and triglycerides [2]. The plant is native to the equatorial and humid tropical climates of Central and West Africa, and Gulf of Guinea. There are two variants of *Dacryodes edulis*: *D. e. var. edulis* and *D. e. var. parvicarpa*. [3][4][5]. Extracts and secondary metabolites have been found to show biological activities such as antimicrobial, antioxidant and anti sickle-cell disease [4]. Studies abound that show that the seeds of *D. eludis* is rich in ash, dry matter, protein, fibre, metabolisable energy, ether extract, essential and non-essential amino acids, vitamins and minerals [1][6][7]. *Garcinia kola* on the other hand, is a medium sized tree that is extensively distributed across West Africa countries. In Nigeria, the seeds of *G. kola* are usually consumed but not the testa for various medicinal purposes [8]. Studies have also shown that it is rich in phytochemicals, nutrients, minerals and vitamins [9][10][11][12].

From antiquity, plants have always been a source of food and medicine to not just man but also to animals. African pear also known as *Dacryodes eludis* is so versatile that the seeds have been used in the production of biodiesel with an extractable yield of 59% that meet acceptable and allowable ASTM (American Society for Testing and Materials) and European Norm (EN) limits [13]. In Nigeria, although the seeds of *G. kola* are very valuable, the thin testa is usually thrown away. Interestingly, the seeds and testa unlike the edible pulp of *D. eludis* are also usually thrown away despite their potential nutritive and medicinal properties. The aim of this study was therefore to examine the seeds and testa of both plants for potential nutrients and anti-nutrients.

MATERIALS AND METHODS

Collection and identification of samples

Freshly harvested samples used in this study were purchased locally from dealer in Etim Ekpo Local Government Area of Akwa Ibom State and identified as *Dacryodes*

eludis and *Garcinia kola*. The seeds and testas were then obtained from both plants and stored separately at room temperature for further analysis.

Proximate Composition Analysis

The samples used in this study were examined for proximate nutrient composition such as moisture, ash, protein, fat and carbohydrate. This was done as previously described by the Association of Official Analytical Chemists [14].

Moisture

Exactly 5g each of the seeds and testas of both plants were weighed and oven dried to a steady temperature of 70°C. The amount of moisture in each samples were then expressed as loss in weight after cool weighing.

Ash Content

Five grams of each sample were placed in a crucible and heated to 550°C to eliminate organic components. The crucible and its contents were then cooled and weighed, and the ash evaluated as a proportion of the original dry weight of samples.

Crude Protein

This was done using the micro-Kjedahl method. The nitrogen proportion of the protein in 5 g of each of the sample was converted into ammonium sulphate by digestion with concentrated hydrogen tetraoxosulphate (VI) acid using copper sulphate as a catalyst. The liberated ammonia was collected in boric acid double indicator solution and the nitrogen quantified through standard hydrochloric acid titration until end point was reached. The amount of crude protein was then obtained by multiplying by a factor of 6.25.

Crude Fat

Crude fat was extracted from both plant samples using 5g of the plant samples, petroleum ether and soxhlet extractor apparatus. The weight of the fat obtained after

evaporating off the petroleum ether from the extract gave the crude fat in the samples and this was expressed as a percentage.

Crude Fibre

Five grams of the defatted samples were used to determine the fibre contents in samples via extraction by acid digestion, filtration and base digestion. The resulting residues were eventually ignited at 550°C. Fibre content was then expressed as a percentage lost on ashing and initial weight.

Carbohydrate

The amount of carbohydrate in each of the sample was then estimated as the difference from 100 of crude protein, fat, ash, and fibre.

Estimation of Anti-nutrients

The anti-nutrients examined in all the seeds and testas of both plants were hydrocyanic acid, phytic acid and oxalate. These were estimated using procedures previously described [15-17].

Hydrocyanic Acid (HCN)

Exactly 10 g of each of the sample was soaked in 300 ml of distilled water for 4hours to allow for the liberation of the cyanide. The liberated cyanide was steam distilled into 20 ml (2.5% w/v) NaOH. About 8 ml of NH₄OH was added to the distillate before titrating with 0.02 M AgNO₃ to a faint and permanent turbidity. The HCN was then estimated using 1 ml of 0.02 AgNO₃ as the equivalence of 1.08 mg of HCN.

Phytic Acid

Two grams of each of the samples were extracted with 0.5M HCl. Ferric chloride was used to precipitate the phytic acid to ferric phytate. NaOH solution was then used to convert the precipitate into sodium phytate and then digested with acid mixture containing equal portion of concentrated H₂SO₄ and HClO₄. The liberated phosphorus

was then quantified colorimetrically at 620 nm after colour development with molybdate solution.

Oxalate

About 2.5 g of the sample was extracted with dilute HCl. The oxalic acid in the extract was precipitated with calcium chloride as calcium salts. The precipitated oxalate was washed with 25% H₂SO₄ and dissolved in hot water before titrating with KMnO₄.

Determination of vitamins

Vitamins A, E, B and C were determined according to methods previously described by [18]-[21].

Determination of mineral elements

The mineral elements were determined by the dry ash extraction method of AOAC [22].

Statistical analysis

Replicate readings were subjected to one way analysis of variance (ANOVA) for each of the samples. The replicate readings for each of the samples were also compared using Student t-test for significance. Probability level was set at 95% (0.05).

RESULTS

The results of the study are presented in Tables 1 to 5. Table 1 shows the parts of the plants used, local names and scientific names. The proximate composition (g/100g) of seeds and testas of both plants are shown on Table 2. The results indicates that the seed of *D. edulis* was richer than the testa in carbohydrate 71.88±0.02, protein 9.62±0.01, ash 5.50±0.02 and fat while the testa was richer in moisture 41.10±0.01 and fibre 13.50±0.02. However, for *G. kola*, the result showed that the seed was richer protein, fat and carbohydrate than the testa which had more fibre, moisture and ash then the seed. Statistical analysis of replicate readings showed significance (p < 0.05).

Table 3 shows the results of the vitamin analysis. The seed and testa of *G. kola* was richer than the seed and testa of *D. eludis* for Vitamin B, C and E except for A which

was just slightly higher for seed and testa of *D. eludis*. Table 4 shows the anti-nutrients composition (mg/100g dry matter) for seeds and testa of both plants. The testa of *G. kola* has the highest concentration of all the anti-nutrients examined for both plants. Both the seed and testa of *D. eludis* had much lesser concentration compared to *G. kola*. The highest anti-nutrient recorded was soluble oxalate with a value of 210.40 ± 0.10 for the testa of *G. kola*. Statistical analysis showed significance ($p < 0.05$).

Table 5 shows the mineral composition analysis of both plants studied. The results indicate that the seed and testa of *G. kola* are particularly rich in potassium, calcium, phosphorus & magnesium compared to the seeds and testa of *Dacryodes edulis*. Sodium, iron, zinc, and copper were not determined for seed and testa of *D. eludis* and similarly, nitrogen was not determined for the seed and testa of *G. kola*.

Table 1: Plants and parts under study

Scientific names	Family	Local name	Plant part used
<i>Dacryodes edulis</i>	Burseraceae	Eben	Seed/Testa
<i>Garcinia kola</i>	Guttiferae	Efiat	Seed/Testa

Table 2: Proximate composition of seeds and testa of *Dacryodes edulis* in g/100g dry matter

Proximate composition	DE		Gk	
	Seed	Testa	Seed	Testa
Moisture	37.00 ± 0.01^a	41.10 ± 0.01^a	2.64 ± 0.02^b	4.60 ± 0.10^b
Ash	5.5 ± 0.02	5.0 ± 0.10	3.50 ± 0.10	4.84 ± 0.02
Protein	9.62 ± 0.01	8.75 ± 0.001	10.81 ± 0.01	9.70 ± 0.10
Fat	8.0 ± 0.10	6.5 ± 0.01	14.90 ± 0.10	12.60 ± 0.02
Fibre	5.0 ± 0.10	13.50 ± 0.02	12.60 ± 0.02	17.80 ± 0.02
Carbohydrate	71.88 ± 0.02	66.25 ± 0.02	57.94 ± 0.02	54.80 ± 0.02

^{a,b}ANOVA of replicate readings gave significant Mean $\pm$ SD ($p < 0.05$). DE = *D. eludis* and GK = *G. kola*.

171 Table 3: Estimation of Vitamins of seeds and Testa of *Dacryodes edulis* (Mean±SD)

Vitamins	DE		GK	
	Seed	Testa	Seed	Testa
A (µg/dl)	141.80 ±0.10 ^a	134.19 ±0.10 ^a	130.30 ± 0.10 ^b	124.70 ± 0.02 ^b
B (mg/dl)	213.07 ± 0.01	215.10 ± 0.10	224.70 ± 0.20	219.40 ± 0.02
C (mg/100ml)	5.12 ± 0.02	6.01 ± 0.01	127.10 ± 0.10	93.30 ± 0.10
E (mg/100g)	2.08 ± 0.02	2.10 ± 0.10	3.14 ± 0.01	2.94 ± 0.02

172 ^{a,b}ANOVA of replicate readings gave significant Mean±SD (p < 0.05). DE = *D. eludis*
 173 and GK = *G. kola*.

174

175

176 Table 4: Anti- Nutrients of seeds and Testa of *Dacryodes edulis* (mg/100g dry
 177 matter)

Anti-nutrients	DE		GK	
	Seed	Testa	Seed	Testa
HCN	1.58±0.01 ^a	1.75 ±0.01 ^a	6.10±0.10 ^b	7.30±0.20 ^b
Soluble oxalate	17.50±0.1	15.62±0.02	167.18±0.01	210.40±0.10
Total oxalate	30.02±0.02	24.62±0.02	44.80±0.10	101.70±0.10
Phytate	0.37±0.01	0.39±0.01	0.90±0.10	1.10±0.01

178 ^{a,b}ANOVA of replicate readings gave significant Mean±SD (p < 0.05). DE = *D. eludis*
 179 and GK = *G. kola*.

180

181

182

183

184 Table 5 Mineral Composition of seeds and testa of *Garcinia kola* in mg/100g Dry matter

Anti-nutrients	GK		DE	
	Seed	Testa	Seed	Testa
Na	10.10±0.1 ^a	18.20±0.02 ^b	ND	ND
K	730.01±0.1	499.19±0.2	0.48	0.56
Ca	200.40±0.2	140.70±0.1	0.32	0.96
Mg	170.49±0.2	166.30±0.	0.14	0.14
Fe	4.20±0.1	1.50±0.01	ND	ND
Zn	3.70±0.02	3.50±0.01	ND	ND
Cu	2.50±0.1	1.32±0.01	ND	ND
P	720.14±0.01	520.10±0.01	0.17	0.18
N	ND	ND	1.54	0.40

185 ^{a,b}ANOVA of replicate readings gave significant Mean±SD (p < 0.05). DE = *D. eludis*
 186 and GK = *G. kola*. ND= Not determined.

187 Discussion

188 African pear fruit pulp is a widely eaten delicacy especially in eastern and southern
 189 Nigeria as a result of its abundant nutrients and minerals [23]. However, despite a
 190 number of studies highlighting the nutritional potentials of the seeds [23][24], it is still
 191 discarded after the edible pulp is eaten [23].

192 In an earlier study, where the possibility of replacing maize with pear seeds in feeds
 193 was examined, they reported the presence of proximate nutrients which are agreeable
 194 to our findings [9]. Interestingly, the proximate composition of the seeds and testa of *D.*
 195 *eludis* in our study are comparable to those of the varieties of *D. eludis* already
 196 examined [1],[25]. Furthermore, they reported the presence of calcium, phosphorus,
 197 potassium, magnesium, sodium, iron, manganese, zinc, copper, selenium and iodine.
 198 They also reported the presence of vitamins like A, D, E, K, in addition to B₁, B₂, B₃, B₆
 199 and B₁₂, biotin, cholin and folic acid [26].

200 *Garcinia Kola* also called Bitter kola is a popular plant in Nigeria and is extensively used
 201 as food and as medicine. Unlike the seeds of *D. eludis*, the seeds of *G. kola* are edible.

In addition, medicinally, it is used to treat and suppress cough and also to relieve colic disorders [27]. Proximate composition (%) in an earlier study on *D. eludis* also confirms the presence of moisture 59.46, fat 4.48, protein 2.48, ash 0.88, fibre 5.01 and carbohydrate 27.69 and minerals such as magnesium, calcium, potassium, Phosphorus, sodium, iron, zinc, copper and cobalt [27].

Our findings for carbohydrate and fibre content were more similar to the proximate analysis (%) of *G. kola* by Mazi *et al* [28] which showed that the presence of moisture 9.28, ash 4.173, fibre 3.940, fat 1.03, crude protein 11.27 and carbohydrate 70.31. However, other components were different. Furthermore, they reported the presence of vitamins B₁, B₂ and B₃, C, E, A, and minerals calcium, magnesium, potassium, sodium and iron that agrees with our study. Furthermore, in another study, the proximate analysis of the nutrient composition of powdered *G. kola* seed showed the presence of moisture, ether extract (EE), crude fibre (CF), crude protein (CP), ash and nitrogen free extracts in the following proportion 7.40, 1.48, 2.94, 3.19, 4.39 and 80.58%, respectively [28]. The study also showed that the seeds also have good lipid lowering effects that could be useful in preventing cardiovascular diseases in humans [29].

From the vitamin analysis, vitamin A was the most abundant of all the vitamins examined. Apart from vitamin A that was the more abundant in the seeds and testa of *D. eludis* than in *G. kola*, consistently, vitamin B, C and E were more abundant in the seeds of *G. kola*. When compared to the vitamin composition of oyster mushroom in an earlier study, vitamin A, B and C in both of our study samples are atleast two times higher in concentration [30].

Studies have shown that anti-nutrients such as oxalate, phytate, hydrocyanic acid in excessive amounts can bring about some adverse health effects on humans [31]. The anti-nutrients levels in *G. kola* in our study are higher than those found in *D. eludis*. The amount of ant-nutrients in *D. eludis* is comparable to those of commonly eaten vegetables previously reported but not those of *G. kola* [31]-[34]. Interestingly, the levels of hydrocyanic acid in both plants samples were far less than those previously reported in cassava and *Citrus paradis* peels [35][36].

CONCLUSION

The findings in this study reveal that the seeds and testa of *D. eludis* and *G. kola* are indeed very rich in nutrients, minerals and vitamins, and also allowable levels of anti-nutrients. However, there is a need for further studies aimed at the utilization of these plants parts in human and animal nutrition.

REFERENCE

- [1] Ibanga OI, Ekpa OD. Minerals and Anti-Nutrients in Two Varieties of African Pear (*Dacryodes edulis*). Journal of Food Technology. 2009; 7(4):106-110.
- [2] Lam HJ, Omoti U, Okiy DA. Characteristics and Composition of the pulp oil and cake of the African Pear, *Dacryodes edulis* (G. don). Journal of Science Food Agriculture. 2006;38(1):67-72.
- [3] Emebiri LC, Nwufo MI. Effects of fruits types and storage treatment on the biodeterioration of African pear (*Dacryodes edulis*) (G. Don) H.J. Lam). Int. Biodeterior. 1990; 26(1): 43-50
- [4] Ajibesin KK. *Dacryodes edulis* (G. Don) H.J. Lam: A review on its medicinal, phytochemical and economical properties. Research Journal of Medicinal Plant. 2011; 5 (1): 32-41.
- [5] Okwu DE, Nnamdi FU. Evaluation of the Chemical Composition of *Dacryodes Edulis*, *Raphia Hookeri* Mann and *Wendl Exudates* used in Herbal Medicine in South Eastern Nigeria. African Journal of Traditional, Complementary and Alternative Medicine. 2008; 5(2), 194–200.
- [6] Iyawe HOT, Osagie VE, Esekheigbe A, Azihm MC. Nutrient composition and Effect of seed Extract of African Black Pear (*Dacryodes eludis*) on Rats. Asian Network for Scientific Information. 2007; 6(5):878-880.
- [7] Bratte L, Mmereole, FUC, Akpodiete OJ, Omeje SI. The Nutrient Composition of Seeds of the African Pear (*Dacryodes edulis*) and its Implications for Non-Ruminant Nutrition. Pakistan Journal of Nutrition. 2010; 9(3); 255-258.
- [8] Dike MC. Proximate, phytochemical and Nutrient compositions of some fruits, seeds and Leaves of some plant species at Umudike Nigeria. ARPN J. Agric. Biol. Sci. 2010; 5(1): 7-16.
- [9] Yete P, Ndayishimiye V, Agbangnan PC, Djenontin ST, Wotto VD, Sohounhloue DCK, Chemical composition of the Seeds and the Defatted Meal of *Garcinia kola* Heckel (Guttiferae) from Benin. Chemistry Journal. 2014; 4(5):13-19.

- [10] Odebunmi EO, Oluwaniyi OO, Awolola GV, Adediji OD. Proximate and nutritional composition of kola nut (*Cola nitida*), bitter cola (*Garcinia kola*) and alligator pepper (*Aframomum melegueta*). African Journal of Biotechnology. 2009; 8(2): 308-310.
- [11] Dah-Nouvlessounon D, Adjanooun A, Sina H, Noumavo PA, Diarrasouba N, parkouda C, Madode YE, Dicko, MH. Nutritional and Anti-Nutrient Composition of Three Kola Nuts (*Cola nitida*, *Cola acuminata* and *Garcinia kola*) Produced in Benin. Food and Nutrition Sciences. 2015; 6:1395-1407.
- [12] Dike MC, Asuquo ME. Proximate, phytochemical and mineral compositions of seeds of *Allanblackia floribunda*, *Garcinia kola* and *Poga oleosa* from Nigerian rainforest. African Journal of Biotechnology. 2012;11(50):11096-11098.
- [13] Ogunsuyi HO. Production of biodiesel using African pear (*Dacryodes edulis*) seed-oil as feedstock. Academic Journal of Biotechnology. 2015; 3(5):85-92.
- [14] AOAC (2002). Official Methods of Analysis -17th ed. Association of Official Analytical Chemist, Maryland.
- [15] Akwaowo EU, Ndon EU, Etuk Eu. Minerals and Anti-nutrients in fluted pumpkins (*Telfaria occidentalis Hook F.*). Food Chemistry. 2000;70:235-240.
- [16] Agbaire PO, Emoyan OO. Nutritional and anti-nutritional levels of some local vegetables from Delta State, Nigeria. African Journal of Food Science. 2012;1:8-11.
- [17] Edet UO, Ebana RUB, Ekanemesang UM, Ikon Gm, Edem E, Mbim EN. Phytochemical Screening, Nutrient Analysis, Antitermite and Antimicrobial Activity of *Citrus paradisi* Peel Powder. Journal of Applied Life Sciences International. 2016; 9(4):1-9.
- [18] Bessy OA. Lowry OH. Brocke MS. Lopper J A. Determination of Vitamin A in Small Quantities of Blood Stream. Journal of Biological Chemistry. 1946; 166:177.
- [19] Bessy OA. Determination of Ascorbic Acid by titrimetric Method. Journal Association Office Agric. Chemist. 1944; 27:537.
- [20] Bender AD. An Introduction to Nutrition and Metabolism. University College, London Press Limited, London; 1993: 182-182.
- [21] Bender AD. Nutritional Biochemistry of Vitamins. Cambridge University Press, London; 2003

- [22] AOAC. Official Methods of Analysis. 13th Edition. *Association of Official Analytical Chemist*, Washington, D.C;1995
- [23] Onuegbu NC, Nwuka MU, Ojukwu M, Kabu NO. Nutritional Properties of African Pear Seed and Performance of Defatted Cake in Poultry Feed Formulations . *Animal Nutrition*. 2016; 1(29):1-3.
- [24] Ajayi IA, Adesanwo O. Comparative study of the mineral element and fatty acid composition of *Dacryodes edulis* pulp and seed. *World J. Agric. Sci.* 2009; 5: 279-283.
- [25] Onuegbu NC, Ihediohanma NC. Some Proximate Analysis of African Pear (*Dacryodes Edulis*). *Journal of Applied Science and Environmental Management*. 2008; 12 (1): 83 – 85
- [26] Bratte L, Mmereole FUC, Akpodiete OJ, Omeje SI. The Nutrient Composition of Seeds of the African Pear (*Dacryodes edulis*) and its Implications for Non-Ruminant Nutrition. *Pakistan Journal of Nutrition*. 2010; 9(3):255-258.
- [27] Alaje DO, Owolabi KT, Olakunle TP, Oluti, OJ, Adetuberu IA. Nutritional, Minerals and Phytochemicals composition of *Garcinia cola* [Bitter cola] and *Aframomum melegueta* [Alligator pepper]. *Journal of Environmental Science, Toxicology and Food Technology*. 2014;8(1):86-91.
- [28] Mazi EA, Okoronkwo KA, Ibe UK. Physico-Chemical and Nutritive Properties of Bitter Kola (*Garcinia Kola*). *Journal of Nutrition Food Science*. 2013; 3(4):1-3
- [29] Omeh YN, Onoja SO, Ezeja MI, Uchendu WC, Okorie E, Raymond M. Quantitative phytochemical, proximate analysis and hypolipidemic effects of *Garcinia Kola*. *British Journal of Medicine and Medical Research*. 2014; 4(36): 5770- 5778.
- [30] Edet UO, Ebana RUB, Etok CA, Udoidiong VO. Nutrient Profile and Phytochemical Analysis of Commercially Cultivated Oyster Mushroom in Calabar, South-South Nigeria. *Advances in Research*. 2016; 7(3):1-6.
- [31] Ladeji O, Akin CU, Umaru. Level of anti-nutritional factors in vegetables commonly eaten in Nigeria. *Afri. J. Nt. Sci.* 2004; 7:71-73
- [32] Ebana RUB, Asamudo NU, Etok CA, Edet UO, Onyebuisi CS. Phytochemical Screening, Nutrient Analysis and Antimicrobial Activity of the Leaves of *Lasianthera africana* and *Dennettia tripetala* on Clinical Isolates. *Journal of Advances in Biology and Biotechnology*. 2016;8(4): 1-9.

- [33] Agbaire PO. Levels of anti-nutritional factors in some common leafy edible vegetables of southern Nigeria. African Journal of Food Science and Technology. 2012; 3(4):99-101.
- [34] Agbaire PO, Emoyan OO. Nutritional and anti-nutritional levels of some local vegetables from Delta State, Nigeria. African Journal of Food Science. 2012;(1): 8-11.
- [35] Edet UO, Ebanu RUB, Ekanemesang UM, Ikon GM, Edem EE, Mbim EN. Phytochemical Screening, Nutrient Analysis, Antitermite and Antimicrobial Activity of *Citrus paradisi* Peel Powder. Journal of Applied Life Sciences International. 2016; 9(4):1-9.
- [36] Alexander J, Guðjón AA, Diane B, Andrew C, Jean-Pierre C, Eugenia D. Carbamate and hydrocyanic acid in food and beverages scientific opinion of the panel on contaminants. EFSA Journal. 2007; 555:1-44.