

Original Research Article

Comparative Assessment of the Physiological and Granules Size Distribution of Starch from Variety of Cassava, Sweet Potato and Cocoyam

ABSTRACT (ARIAL, BOLD, 11 FONT, LEFT ALIGNED, CAPS)

SAMPLE ABSTRACT:

Aims: To produce starch from cassava variety varieties (30572, 419, and umu 37), sweet potato root (*x-igbariam*), cocoyam (*edeuhie* and NXS003). To sturdy and compare their functional, pasting and micro-structural properties.

Study Design: Multifactor randomized complete block design (RCBD) was used for this study. Each experiment was repeated in duplicate. Data obtained were subjected to analysis of variance (ANOVA) at the significant level of 5% ($p \geq .05$) using SPSS 20.

Place and Duration of Study: Department of food science and technology, Michael okpara university of agriculture, Umudike, between May 2016 and october 2016.

Methodology: A Total of 6 (six) starch samples were produced from cassava variety varieties (30572, 419, and umu 37), sweet potato root (*x-igbariam*), cocoyam (*edeuhie* and NXS003). The starch samples were analyzed for functional, pasting and micro-structural properties..

Results: The starch from different tuber crops analysed showed different functional, pasting and micro-structural properties. At $p \geq .05$ significant level, Pasting temperature was highest in cocoyam starch sample (ede) while cassava starch sample 419 was the lowest. Peak, trough and breakdown result was highest in sweet potato starch while peak was lowest in Cocoyam starch sample. solubility was highest in cocoyam (Ede) and lowest in cassava sample 419. The microscopic picture of starches showed spherical shape for all granules with cocoyam starch having smooth, fine and regular circle shape. Cassava starch sample 30572 was highest in granular size while 419 recorded lowest granular size.

Conclusion: Starch from different plants and varieties significantly differs in their functional, pasting and micro-structural properties. Different starch products requires different physiological properties and when starch source is rightfully selected, different starch source shows optimum functionalities in specific product.

Keywords: starch, microstructure, physiology, cassava, cocoyam sweet-potato.

1. INTRODUCTION

Tubers and roots are important sources of carbohydrates as an energy source and are used as staple foods in tropical and sub tropical countries (Liu *et al.*, 2006) [1]. Starch granules are insoluble in cold water but swell reversibly to a limited extent through hydrogen bonding. With additional heat, the granules continue to swell. As the temperature is increased, the thermal energy overcomes the inter- and intra-molecular hydrogen bonds and hydrophobic interactions, resulting in irreversible disintegration of granular order. In the gelatinization process, the point at which the native crystalline regions are melted, birefringence is lost (French 1984) [2]. Continued heating will result in a colloidal solution. Generally, the gelatinization of starch granules occurs over a temperature range that differs for each starch type. Examples include; normal maize, 64.4 to 80.4°C; waxy maize, 64.2 to 80.4°C; and normal potato, 59-68°C (Jane *et al.* 1997) [3]. Gelatinization temperatures of starch are affected by granule size (Banks and Greenwood 1975) [4], degree of crystallinity (Zobel, 1984) [5], relative amounts of amylose and amylopectin (Inouchi 1983) [6], presence of phosphorous derivatives (Lim and Seib 1993) [7], lipids (Morrison 1988) [8] and the amylopectin branch-chain length (Kasemsuwan and Jane 1995) [9]. Pasting

follows gelatinization and was defined by Atwell et al. (1988) [10] as "the phenomenon following gelatinization in the dissolution of starch. It involves the granular swelling, extrudation of molecular components from the granule and eventually total disruption of the granules." In both food and industrial applications, pasting is an extremely important attribute of the functionality of starch. Starch pastes are extensively used as thickeners, stabilizers and binders. Paste viscosity of starch, as measured by Brabender Visco-amylograph or Rapid Visco Analyzer, is often used as a critical acceptance measurement in quality control processes by both starch manufacturers and end-use industries. Also tubers and roots do not contain any gluten, which is an important factor when considering a carbohydrate source. Using tubers as a source of carbohydrate instead of gluten containing carbohydrates, may aid in a reduction in the incidence of cardiac disease (CD) or other allergic reactions. Selection and use of starches in food products require basic knowledge of food processing and starch properties (Moore et al. 1984) [11]. Therefore, the study of starch and its molecular structures and functional relationships is important. The understanding of starch molecular structures gives direction to breeders, starch producers and processors. Today, there is growing interest in natural starches with unique properties for use in foods to avoid starch modified" label statement]. Therefore, the present study was aimed to assess the Rheological, Micro-structure and functional properties of starch from tubers, in an attempt to broaden what applications they may be used for within the food industry. The tuber starches assessed in this study were cassava (419 um37 and 30572) sweet potato (*X-igbariam*) and cocoyam (*edeahie* and NX003)

2. MATERIAL AND METHODS / EXPERIMENTAL DETAILS / METHODOLOGY

2.1 SOURCE OF MATERIALS

Matured roots of four cassava varieties (30572, 419, and umu 37), one sweet potato root (*x-igbariam*) and two cocoyam (*edeuhie* and NXS003) were obtained from National Root Crops Research Institute (NRCRI), Umudike and processed into starch within 24hrs after harvest.

2.2 PROCESSING OF STARCH

The method described in Osunsami *et al.*, (1989) [12] with slight modification was used for the production of starch from cassava, sweet potato and cocoyam samples. Matured roots of cassava (30572, 419 and umu 37) sweet potato (*x-igbariam*) and cocoyam (*edeuhie* and Nx003) were peeled and washed with clean water and allowed to drain. They were subsequently grated with locally fabricated grater. The pulp was placed on a muslin cloth, sieved and flushed with clean water until the starch stream ceases. The crude starch milk was kept in a refrigerator overnight. The supernatant were decanted leaving the precipitated starch.

2.2.1 Purification of starch

Exactly 1000ml of clean water was added to 300g of crude starch and stirred until all the starch was suspended in the water. The suspension was undisturbed until sedimentation ceased. The supernatant was decanted leaving the sediments in a plastic bucket. An equivalent volume of fresh clean water was added to the volume of the starch sediments and stirred until the starch was suspended in the water. The starch slurry was filtered through a muslin cloth and flushed with water. The filtrates were allowed to sediment for 20min and the supernatant decanted. This procedure was repeated two (2) times yielding clean starch cake.

2.2.2 Quality control of the starch

A teaspoonful sample was taken from the top of the clean starch cake sediment and dissolved in a 15ml of water in a test tube. The tube and its content were centrifuged for 20min at 2000 rpm and a small sample was drawn from the boundary between the liquid and the sediment using a pipette. Observation was done under a microscope (magnification x400). Absence of particles in the samples under observation confirmed its purity.

2.2.3 Drying and storage

The starch flakes were spread on a table and allowed to sundry for two days. They were pulverized with a milling machine (Thomas Wiley mill-model Ed-5, Philadelphia, USA 1982), stored in bottles and preserved in a refrigerator (Haier thermacool, model HRF-350,2001) (Ubalua, 2014) [13].

2.2.4 Starch yield

Percent starch was calculated on dry bases as

$$\frac{\text{Weight of ground tuber} - \text{weight of starch}}{\text{Weight of ground tuber}} \times \frac{100}{1}$$




Figure 2.1: Flow Chart for Production of Starch from Root Crops

2.2.2 Rheological properties of starch

Paste viscosity/ Rheology were determined using the Rapid Visco Analyser (RVA). Measuring the viscosity of cooked paste of starch based product. Starch (3g) was dispersed in 25ml distilled water and quantitatively mixed and transferred into the canister of RVA model 4500 as recommended. The slurry was heated from to 95°C with a holding time of 2min. The rate of heating and cooling were at constant rate of 11.25°C per min. Peak viscosity, trough, breakdown, final viscosity, setback, peak time and pasting temperature were read from the pasting profile with the aid of thermocline for windows software connected to a computer (New port scientific, 1998) [14]. The viscosity was expressed in terms of Rapid visco unit (Rvu) which is equivalent of 10 centipoises.

2.2.3. Functional properties of starch

i. Water absorption capacity :

The procedure of Sathe *et al.* (1982) [15] was used. Exactly 10ml of water was added to 1.0g of each sample. The suspension was then stirred for 5min. The suspension was transferred into centrifuge tube and centrifuged at 3,500rpm for 30 min. The supernatant obtained was measured using 10ml measuring cylinder. The density of water was assumed 1 g/ml. The water absorbed was calculated as the difference between the initial water used and the volume of the supernatant obtain after centrifuging. The result was expressed as a percentage of water absorbed by the starch on g/ml basis.

ii. Oil absorption capacity:

The procedure of Sathe *et al.* (1982) [15] was used as described above. Instead of water used, refined soybean oil with density of 0.92g/ml was used. The oil and the starch blends (1g starch in 10ml oil) were mixed using a magnetic stirrer at 1,000rpm for 5 min and then centrifuged at 3,500rpm for 30 min. The amount of oil separated as supernatant was measured using 10ml cylinder. The difference in volume was taken as the oil absorbed by the samples. The oil absorbed was expressed as g/ml of oil absorbed.

iii. Swelling capacity:

The methods describe by Safa-dede and Saalia (1997) [16] was used. About 1g of the sample were weighed into 10ml graduated measuring cylinder and volume measured. The samples were mixed with water stirred and allowed to stand for 1h. Increase in volume was measured in g/ml.

$$\text{Swelling index} = \frac{\text{final weight} - \text{initial weight}}{\text{weight of sample}} \times \frac{100}{1}$$

iv. Solubility:

The total solubility of the starch samples at room temperature was determined using the method of Singh (1987) [17]. Exactly 1 g of the sample was weighed into a centrifuge tube containing 10ml distilled water. The mixture was stirred and allowed to stand for 1h before being centrifuged at 4,000rpm for 15min. The supernatant was evaporated in a previously clean, weighed moisture can. The solubility was measured as increase in weight of the can over the weight of the sample and expressed in percentage.

iv Bulk density:

The bulk density of the starch samples was done using the method described by AOAC (1990) [18] About 10ml capacity graduated measuring cylinder was weighed and gently filled with the samples. The bottom of the cylinder was tapped several times until there was no further diminution of the sample level after filling to the 10ml mark.

Calculation:

$$\text{Bulk density (g/ml)} = \frac{\text{weight of sample}}{\text{volume of sample}}$$

V Starch granule morphology

Starch granule morphology of native starch from cassava, sweet potato and cocoyam were studied using scanning electron microscopy (SEM). One gram of dry starch was suspended in 10ml of water at 25°C

and gently stirred for 5 min. Starch samples were stained with crystal violet, mounted on circular aluminum stubs using adhesive and then coated with thin layer of gold using Bio-Rad Sputter coating System. The samples were examined and photographed in a Joel scanning microscope (JSM-6400, TOKYO, Japan) at an accelerated voltage of 5kv and magnification of X400. Twenty (20) granules were selected randomly and their size measured.

Vi Gelatinization temperature

Exactly 10% starch dispersion were prepared in tube with distilled water. The suspension was mixed thoroughly and heated. The temperature at which the starch starts gelling is recorded as the gelatinization temperature (°C).

3. RESULTS AND DISCUSSION

3.1. PASTING PROPERTIES OF SELECTED ROOT CROP STARCHES

Amylograph pasting viscosity studies on the starches in Table 3.1 showed Pasting temperature rang of 71.5 to 86.45. Cocoyam starch sample (ede) was highest while cassava starch sample 419 was the lowest in Pasting temperature. There were significant differences ($p \geq .05$) among the samples analyzed. The lower pasting temperature, the better the gelling ability of starch. Pasting temperature is considered as a reflection of the degree or order arrangement of molecules in starch granules. Onset temperature is influenced by short amylopectrin branch-chains and low Pasting temperature is characteristics of starches with large proportions of short amylopectin branch chains (Jane *et al.*, 1999) [19]. Pasting temperature is an indication of granular architecture (crystalline quality) and high peak temperature might be due to higher proportion of longer chains in the amylopectins as these requires higher temperature to dissociate completely than required for shorter double helices. (Karim *et al.*, 2007) [20]. Higher Pasting temperature for cocoyam starches that compared to cassava starches suggests the presence of higher proportions of long amylopectin chains in cocoyam starch. Peak, trough and breakdown result was highest in sweet potato starch (pot = 630.33 Rvu, 300.25 Rvu Bu and 330.09 Rvu respectively) while peak was lowest in Cocoyam starch sample (ede = 305.05 Rvu), Cassava starch sample 419 was lowest for trough, (150.38 Rvu) and break down (228.33 Rvu). There were significant differences ($p \geq .05$) among the starches analyzed. Ricard (1991) [21] reported a similar result saying that *Xanthosoma* starch has low viscosity values. Cocoyam is not susceptible to retrogradation and this makes it suitable source of carbohydrate in complementary food production (Oti and Akobundu, 2008) [22]. The difference in pasting properties of starches are attributed to factors such as starch composition (amylose to amylopectin ratio, amount of lipid complex amylose chains, phosphorus content), molecular structure e.g. Amylopectin (unit chain and extent of branching and granular architecture (crystalline to amorphous ratio) (Gunaratine and Hoover, 2002) [23]. Thickening and swelling are desired functionalities governing the use of starch in food production such as sauce. These two functionalities depend on pasting characteristics of starch (Karim *et al.*, 2007) [20].


Table 3.1 Pasting properties of starch from cassava (419, um37, 30572), sweetpotato (x-igbariam) and cocoyam (nxs003, edeuhie) tuber.

Sample	Peak 1 (Rvu)	Trough 1 (Rvu)	Breakdown (Rvu)	Final viscosity (Rvu)	Setback (Rvu)	Peak time (sec)	Pasting temperature (°C)
419	378.71 ^c	150.38 ^e	228.33 ^d	220.25 ^d	69.88 ^b	4.13 ^b	74.35 ^d
um37	430.38 ^b	155.42 ^e	274.96 ^b	224.21 ^d	68.79 ^b	3.70 ^c	71.50 ^f
30572	436.67 ^b	177.63 ^c	259.05 ^c	221.58 ^d	43.84 ^d	3.74 ^c	72.73 ^e
Pot	630.05 ^a	300.25 ^a	330.09 ^a	355.33 ^a	55.09 ^c	4.24 ^c	78.33 ^c
Nxs	343.54 ^d	183.63 ^b	137.42 ^b	291.33 ^b	107.71 ^a	4.60 ^a	84.93 ^b
Ede	305.05 ^e	166.04 ^d	139.00 ^e	277.05 ^c	111.00 ^a	4.67 ^a	86.45 ^a
LSD	0.054	0.073	0.782	0.198	0.148	0.063	1.00


Means with different superscript within the same column are significantly different ($p \geq 0.05$) 419, um37 and 30572 are cassava starch, ede (*edeuhie*) and Nxs (*Xanthosoma*) are cocoyam starch, pot (*x-igbariam*) is white skinned sweet potato.

Plate 3.1 Rapid Viscoamylo-graph of starch from cassava (419, um37, 30572), sweet potato (*x-igbariam*) and cocoyam (nxs003, *edeuhie*) tuber


cassava 419 starch




cassava 30572 starch



247 **cassava um37 starch**




248
249
250
251
252 **cocoyam edeuhie starch**




253
254
255
256
257

258 Cocoyam nxs003 starch



259
260
261
262
263

Sweet potato (x-igbariam) starch



264
265
266
267
268
269

3.2 FUNCTIONAL PROPERTIES OF STARCH FROM CASSAVA (419, UM37, 30572), SWEET POTATO (X-IGBARIAM) AND COCOYAM (NXS003, EDEUHIE).

i Bulk density

From Table 3.2, the bulk density of the starches from cassava sample 419 was the highest (0.81g/ml) and lowest (0.28g/ml) in sweet potato starch samples, bulk density of starches analyzed were generally low. There were significant differences among the samples in terms of bulk density. High bulk density limits caloric and nutrient intake which result in growth faltering (Onimawo and Egbekun, 1998) [24]. Bulk density is affected by particle size and density of the sample (Karuna *et al.*, 1996) [25]. Plaami (1997) [26]. reported that bulk density is influenced by the structure of starch polymers and loose structure of starch results in low bulk density. Low bulk density observed in starch samples in the work is

an advantage. Low bulk density gives an indication of the greater ease of dispersability and reduction of paste thickness (Udensi and Eke, 2000) [27]. Low bulk density is easily carried and distributed to locations where they are required (Agunbiade and Sanni, 2003) [28].

ii Water absorption capacity (WAC)

Water absorption capacity result in Table 3.2 ranged from 150 for cassava sample 419 to 180 for cocoyam sample Nxs003. There were significant differences ($p \geq 0.05$) in WAC among the samples. Water absorption capacity is a useful indication of whether flour can be incorporated into aqueous food formulation, especially those involving dough handling (Circle and Smith, 1972) [29]. Water absorption capacity is important in bulking and consistency of products as well as baking application (Niba *et al.*, 2001) [30]. Better water absorption capacity suggests better performance in texture (Okezie and Bello, 1988) [31]. Increase in water absorption capacity implies increase in digestibility of starches. The difference might depend on the amount and nature of hydrophilic constituent (Ayele and Nip, 1994) [32]. Water absorption capacity reveals the inter-molecular associations between starch polymers (Rincon *et al.* 1999) [33].

iii Oil absorption capacity

Oil absorption capacity of the starch samples were between 1.2ml/g to 2.4ml/g. Cassava starch sample 30572 was lowest while cocoyam starch sample Nxs were the highest. There was significant difference ($p \geq 0.05$) among the samples in terms of oil absorption capacity. The ability of starch of these roots crops to bind with oil makes it useful in food system where optimum oil absorption is desired. The high oil absorption capacity suggests that cocoyam starch may contain more hydrophobic proteins than cassava and sweet potato. More hydrophobic proteins demonstrate superior binding of lipids (Lawal and Adebawale, 2004).

iv Swelling index

Swelling index of starches in Table 3.2 ranged from 0.105g/ml to 0.505g/ml. cassava starch 419 was the lowest while sweet potato (pot) was the highest. There was significant difference ($p \geq 0.05$) among the starches analyzed in terms of swelling index.

Swelling index is an indication of the water absorption index of the granules (Loos *et al.*, 1981) [35]. Sanni *et al.*, (2005) [36] reported that the swelling index of the granules reflects the extent of associative force within the starch, the lower the association the lower the swelling power (Sanni *et al.*, 2001) [37]. Swelling index of starch depend on the capacity of starch molecule through hydrogen bonding (Tang *et al.*, 2005) [38]. High swelling index in sweet potato and cocoyam sample (Nxs003) may be as a result of differences in the amylopectin content of the various starches since swelling index has been describes as amylopectin property (Li and yeh 2001, Wrolstad, 2012) [39, 40] .

v Solubility

The result of solubility in Table 3.2 recorded highest in cocoyam Ede (1.71g/m) and lowest in cassava sample 419 (1.5g/m). There were no significant differences in the solubility ($p \leq 0.05$) among the starch analyzed. Low fat content leads to increase in solubility. Kim and Wiesenborn (1996) [40] observed that increase in solubility was attributed to the low fat content and weak internal starches. Brambridge *et al.* (1996) [41] stated that good quality starch with high starch content and paste viscosity will have a low solubility, high swelling volume and swelling power. This was observed in this work. The difference in solubility and swelling power indicates difference in structure and solubility of starch and also could be attributed to difference in chain length/distributions in starches (Bellow-Perez *et al.*, 2000) [42].

Vi Gelatinization temperature

The result of gelatinization temperature in Table 3.2 ranged from 69.25 to 90.1. Cassava starch sample 30572 was lowest while sweet potato starch sample recorded highest gelatinization temperature. There was significant difference ($p \geq 0.05$) in the gelatinization temperature among the starch samples analyzed. Gelatinization affects digestible and texture of starch content of foods (Ricard, 1991) [43]. The gelatinization temperature gives an indication of the strength of associative force within the starch granules, so higher associative force leads to lower gelation temperature. Cassava starch samples

showed lower gelatinization temperature than cocoyam and sweet potato. This is in line with the works of Erikson *et al* (2014) [44] which reported lower gelatinization temperature for cassava starch which they translated into shorter cooking time than wheat flour.

Table 3.2: Functional properties of starch from cassava (419, um37, 30572), sweet potato (*X-igbariam*) and cocoyam (Nxs003, *edeuhie*).

Sample	Bulk density ^(g/ml)	Swelling Index ^(g/ml)	solubility	Gelatinization temperature(°C)	Visual color	gel	WAC ^(g/ml)	OAC ^(g/ml)
419	0.81 ^a	0.105 ^e	1.50 ^a	72.5 ^d	Transparent		150.0 ^c	180.0 ^b
um37	0.43 ^b	0.200 ^d	1.68 ^a	67.15 ^g	Milky		160.0 ^c	160.0 ^c
30572	0.41 ^b	0.410 ^b	1.65 ^a	69.25 ^e	White		170.0 ^b	120.0 ^e
Pot	0.28 ^c	0.505 ^a	1.62 ^a	90.10 ^a	Golden		190.0 ^a	160.0 ^c
Nxs	0.71 ^a	0.325 ^c	1.62 ^a	85.05 ^b	Dirty brown		180.0 ^a	240.0 ^a
Ede	0.57 ^b	0.475 ^{ab}	1.71 ^a	83.25 ^c	Light purple		170.0 ^b	140.0 ^d
LSD	0.054	0.051	0.05	1.00			0.267	1.00

Means with different superscript within the same column are significantly different (p≥.05). 419, um37 and 30572 are cassava starch, ede (*edeuhie*) and Nxs (*xanthosoma*) are cocoyam starch, pot (*x-igbariam*) is white skinned sweet potato. WAC = water absorption capacity, OAC = oil absorption capacity

3.3 GRANULES SIZE DISTRIBUTION OF STARCH FROM CASSAVA (419, um37, 30572), SWEET POTATO (*x-igbariam*) AND COCOYAM (Nxs003, *edeuhie*)

The result of granule size distribution are shown in Table 3.3 The granule size ranged from 27.3µm to 140Nm. Cassava starch sample 30572 was highest in granular size (273Nm) while 419 recorded lowest granular size (140Nm). Sweet potato starch was highest in terms of the size range (220-300Nm). Sweet potato starch also showed a high particle size of 254Nm.

The surface of the starch granules of samples analysed appeared to be smooth and fine with no evidence of fissures. This is because no treatment was given to the starch prior to analysis. The starches used for this study had a fine particle size distribution. Particle size affects the rates of hydration during processing as very fine (<180Nm) particle size have greater tendency to absorb more water during hydration (Hatcher *et al.*, 2009) [45] Hou and Kruk (1998) [46] reported that large particle requires time for water to incorporate and tend to form larger dough lumps. Tian *et al.*, (1991) [47] suggested that small granules have higher solubility and hence enhanced water absorption capacity which have positive implication for functionality of flour during processing, often creates more cohesion in most baking systems larger granules on the other hand would be insufficiently hydrated. Optimum dough mixing would thus require fine and evenly distributed particles size flour at most 300Nm for easy hydration. (Tian *et al.*, 1991) [48].

3.3.1 Starch microscopy of starch from cassava (419, Um37, 30572), sweet Potato (*x-igbariam*) and cocoyam (Nxs003, *edeuhie*)


The microscopy results in plate 3.1 showed spherical shape for all the starches visualized. Cocoyam starch particles showed a smooth, fine and regular circle shape. This result is in line with the findings of Benes (2005) [49] which reported mostly round or oval shapes with flat surface on one side containing conical pit which extends to all wall for cassava starches. Other researchers reported Round truncated, cylindrical and spherical for cassava starch granules (Gunaratine and Hoover, 2002; Sriroth *et al.*, 1999; Mishra and Rai, 2006) [23] [50] [51] Hatcher *et al.* (2009) [52] reported that particle size distribution of milled flours affects the rate of hydration during processing, as very fine (<180µm) particle-sized flours have greater tendency of absorbing more water during hydration.

375 **Table 3.3 Granules size distribution of starch from cassava (419, um37, 30572), sweet**
 376 **potato(*x-igbariam*) and cocoyam (Nxs003, *edeuhie*)**
 377


Sample	Size range Nm	Mode Nm	Mean Nm
419	120-160	130	140
um37	200-350	210	244
30572	180-300	180	273
Pot	220-300	300	264
NXS	100-200	100	154
Ede	100-200	100	154

378 **419, um37 and 30572 are cassava starch, ede (*edeuhie*) and Nxs (*xanthosoma*) are cocoyam**
 379 **starch, pot (*x-igbariam*) is white skinned sweet potato**
 380
 381
 382
 383
 384
 385
 386
 387
 388
 389
 390
 391
 392
 393
 394
 395
 396
 397
 398
 399
 400
 401
 402
 403
 404
 405
 406
 407
 408
 409
 410


Plate 3.1 Microscopic structure of starch from selected root crops




Cassava 419 starch




sweet-potato (*x-Igbariam*) starch




Cassava 30572 starch



cocoyam edeuhie starch



Cassava um37 starch



cocoyam Nxs003

Key: 419, um37 and 30572 are cassava starch, cocoyam (*edeuhie*) and Nxs003 (*xanthosoma*) are cocoyam starch, pot (*x-Igbariam*) is white skinned sweet potato

4. Conclusions

The starch from different tuber crops analyzed showed different physiological and micro-structural properties. Pasting temperature was highest in cocoyam starch sample (ede) while cassava starch sample 419 was the lowest in Pasting temperature. This suggests the presence of higher proportions of long amylopectin chains in cocoyam starch. Peak, trough and breakdown result was highest in sweet potato starch while peak was lowest in Cocoyam starch sample. Solubility was highest in cocoyam (Ede) and lowest in cassava sample 419. The microscopic picture of starches showed spherical shape for all granules with cocoyam starch having smooth, fine and regular circle shape. Cassava starch sample 30572 was highest in granular size while 419 recorded lowest granular size.

Starch from different plants and varieties significantly differs in their functional, pasting and micro-structural properties. Different starch products require different physiological properties and when starch source is rightfully selected; different starch source shows optimum functionalities in specific product.

REFERENCES

- [1] Liu, Q., Donner, E., Yin, Y., Huang, R.L. and Fan, M.Z.. The physicochemical properties and in vitro digestibility of selected cereals, tubers, and legumes grown in China. *Food Chemistry*. 2006, 99: 470-477
- [2] French D. Organisation of Starch Granules In: Whistler, R. S., BeMiller J.N. and Paschall, E.F. (eds.) second edition, *Starch Chemistry and Technology*, Academic Press, New York. 1984, 183-247
- [3] Jane, J., Wong, K.S. and McPherson, A.E. Branch-structure difference in starches of A- and B- type X-ray patterns revealed by their Naegali dextrans. *Carbohydrate Research*. 1997, 300: 219-227.
- [4] Banks, W., and C. T. Greenwood: *Starch and its Components*. Edinburgh University Press, Edinburgh (England.) 1975. 342
- [5] Zobel, H. F.. Starch Gelatinization and Mechanical Properties of Starch Pastes. in: *Starch: Chemistry and Technology*. R. L. Whistler, J. N. Bemiller, and E. F. Paschall, eds. Academic Press: Orlando, FL. 1984, 285
- [6], Inouchi, N., Glover, D. V., Sugimoto, Y., and Fuwa, H. Developmental changes in starch properties of several endosperm mutants of maize. *Starch/Staerke*, 1984, 36, 8-12
- [7] Lim S. and Seib, P. A. Preparation and Pasting Properties of Wheat and Corn Starch Phosphates, *Cereal Chem*. 1993, 70:137-144 Copyright 1993 by the American Association of Cereal Chemists, Inc.
- [8] Morrison, W.R. Lipids in Cereal Starches: A Review *J. cereal sci.* 1988, 8:1
- [9] Kasemsuwan, T., and Jane, J. Location of amylose in normal starch granules. Q. Location of phosphodiester cross-linking revealed by phosphorous-31 nuclearmagnetic resonance. *cereal Chem*. 1995, 71:282-287.
- [10] Atwell, W. A., Hood, L. F., Lineback, D. R., Varriano-Marston, E., and Zobel, H. F. The terminology and methodology associated with basic starch phenomena. *CerealFoods World*, 1988 33:306-310.
- [11] Moore, C. O., Tuschoff, J. V., Hastings, C. W., and Schanefelt, R. V. Applications of starches in foods. 1984.575-591 in R. L. Whistler, J. N. BeMiller, and E. F. Paschall, eds. *Starch Chemistry and Technology*. Second Edition. Academic Press, London
- [12]. Osunsani, A. T., Akingbala, Y.O., Oguntimehin, G. B. Effect of storage on starch content and modification of cassava, *starch/starch starke* 1989. 41: 54–57.
- [13] Ubalua, A.O. The use of corn starch for growth and production of α -amylase from *Bacillus subtilis*. *Journal of Microbiological Research*, 2014, 4(4): 153-160.
- [14] Newport Scientific. Interpretation, In: *Applications Manual for the Rapid Visco Tm Analyser* . Australia Newport Scientific Pty. Ltd. 1998. 13-16
- [15] Sathe SK, Deshpande SS, Salunkhe DK (1982). Functional properties of winged bean (*Psophocarpus tetragonolobus*, L) proteins. *J. Food Sci.* 47: 503-506.
- [16] Safa-dedeh, S., and Saalia, F.K. (1997).Extrusion of maize cowpea blend in a modified oil expeller.*Journal of Science of Food and Agriculture*, 73: 160-168.
- [17] Singh, K.B. Chickpea breeding. 1987, 127-162. In: M.C. Saxena and K.B. Singh (eds.), *The Chickpea*. CAB International, UK.

- 488 [18] AOAC. Official Method Of Analysis, 13th Ed., Association Of Official Chemists USA 1990, 233-250
- 489 [19]. Jane J. Chenyy, Lee L. F., Mcpherson A. E., Wong K. S., and Radosavjevic M. (1999). Effect Of
490 Amylobeetin Branch Chain Length and Amylose Content On The Gelatinization And Pasting Properties
491 Of Starch. *Cereal Chemistry*. 76(5):629 – 637
- 492 [20] Karim A.A., Toon L.C., Lee V.P., Ong W.Y., Fazilah A. and Noda T. (2007). Effects of Phosphorus
493 Contents on The Gelatinization and Retrogradation Of Potato Starch. *Journal of Food Science*. 2007.
494 72(2): 132 _ 138
- 495 [21] Ricard, J.E., M. Asoaka and M.V. Blanshard, The physico-chemical properties of cassava starch.
496 *Tropical Science*, 1991. 31: 189-207.
- 497 [22]. Oti, E and Akobundu, E.N.T, Physicochemical and Functional Properties of Some Cultivars of
498 Cassava. *www.scholar.google.com*, 2008. retrieved 19/3/2012
- 500 [23] Gunaratine, A., and Hoover, R. Effect of heat moisture treatment on the structure and
501 physiochemical properties of tuber and root starches. *Carbohydrate polymers*, 2002. 49, 425-437.
- 502 [24]. Onimawo, A.I. and Egbekun K.M. (1998) *Comprehensive Food Science and Nutrition Revised*
503 *edn Grace foundation pub, Jos 1998. 42-64*
- 504 [25] Karuna D., Noel G. and Dilip K. *Food and Nutrition Bulletin*, 1996.17: 2, United Nation University
- 505 [26]. Plaami, S., Content of dietary fiber in food and its physiological effect. *Food Review International*,
506 1997.13: 27-76
- 507 [27]. Udensi, A., and Eke, O., Proximate composition and functional properties of flour produced
508 from *Mucuna cochinensis* and *Mucuna utles*. In: *Proceedings of the 1st Annual Conference of the College*
509 *of Agriculture and Veterinary Medicine, Abia State University. 10–13th Sept. 2000. 170–174*
- 510 [28]. Agunbiade, S. O. and Sanni, M. O. The effect of ambient temperature of cassava tubers on
511 Starch quality. In: *Root Crops. The small processor and development of Local Food Industries for market*
512 *economy. Proceedings of the Eight Triennials Symposium of the International Society for Tropical Root*
513 *Crops. African Branch (ISTRAC-AB) 12-14 Nov. IITA, Ibadan Nigeria. 2003. 180-194.*
- 514 [29]. Circle, S. J. and Smith, A.K. (1972). Functional properties of commercial edible soybean Protein
515 production. In: *Symposium of Seed and Proteins (Inglett, G. E. ed). Avi Pub.Co.Inc. Westport*
- 516 [30]. Niba L.L, Bokanga M.M, Jackson F.L, Schlimme D.S, Li, B.W. physicochemical properties and
517 starch granular characteristics of flour from various *manihot esulenta*(cassava) genotypes. *Journal of*
518 *Food Science*, 2001. 67:1701-1705.
- 519 [31]. Okezie B.O. and Bello A.B. Physicochemical and functional properties of winged Bean flour and
520 isolate compared with soy isolate. *Journal of Food Science*. 1988. 53(2): 450-454
- 521 [32]. Ayele, T., and Nip, W.K. Functional properties of raw and precooked taro (*Colocasia esculenta*)
522 flours. *International Journal of Food Science and Technology*, 1994. 29: 457-4
- 523 [33] Rincon-Saab G., del-Rio, G., Santamaria, R.I., Lopez-Munguia, A. and Soberon, X. (1999) *FEBS Lett.*,
524 453, 100±106.
- 525 [34]. Lawal OS, Adebawale KO (2004). Effect of acetylation and succinylation on solubility profile, water
526 absorption capacity, oil absorption capacity and emulsifying properties of muncuna bean (*Mucuna*
527 *pruiens*) protein concentrate. *Nahrung/Food*, 48(2): 129-

- 542 136.
- 543 [35]. Loos, D. S., Schrman, D.J. and Feldman, D. A corticosteroid binding protein and endogenous
- 544 ligand in *C. albicans* indicating a possible steroidreceptor system. *Naturescience*, London 1981. 293, 477-
- 545 479
- 546 [36] Sanni SA, Okeleye KA, Soyode AF, Taiwo OC . Physiochemical Properties of Early and Medium
- 547 Maturing Nigerian rice varieties. *Nig. Food J.* 2005, 23: 148-152.
- 548
- 549 [37]. Sanni, L.O., Ikuomola, D.P. and Sanni, S.A.,. Quality of gari (roasted cassava mash) in Lagos State,
- 550 Nigeria. 2001, 208 -211.
- 551
- 552 [38]. Tang J., Naitow H., Gardner N.A., Kolesar A., Tang L., Wickner R.B. and Johnson J.E. The
- 553 structural basis of recognition and removal of cellular mRNA 7-methyl G 'caps' by a viral capsid protein: a
- 554 unique viral response to host defense. *Journal of Molecular Recognition* 2005. 18(2):158-68
- 555
- 556 [39] Li, J.-Y., & Yeh, An-I.. Relationships between thermal, rheological characteristics and swelling power
- 557 for various starches. *Journal of Food Engineering*, 2001, 50(3), 141–148
- 558
- 559 [40] Wrolstad R, *Food carbohydrate chemistry*. John Wiley and sons Inc. and IFT press, Iowa. 2012. 107-
- 560 131
- 561
- 562 [40] Kim Y.S. and Wiesenborn D. P. Starch Noodle Quality as Related to Potato Genotypes, *J. Food Sci.*
- 563 61,1: 1996 244-252.
- 564
- 565 [41] Bainbridge Z, Tomlins K, Willings K, Westby A. Methods for assessing quality characteristics of non
- 566 grain starch staple. Part 4 advanced methods. *National Ressources Institute*. University of Greenwich, UK
- 567 ISBN 0 – 85954 – 400 1996. 1:43–79
- 568
- 569 [42]. Bello-Perez, L.A., Contreras-Romos, S.M., Jimenez-Aparicio and Paredes-Lopez, O. Acetilation and
- 570 characterization of banana (*musa paradisiaca*) starch. *Actacient Venezolana*, 2000. 51: 143-149.
- 571
- 572 [43]. Ricard, J.E., M. Asoaka and M.V. Blanshard, The physico-chemical properties of cassava starch.
- 573 *Tropical Science*, 1991. 31: 189-207.
- 574
- 575 [44] Eriksson, E., Koch, K., Tortoe, C., Akonor, P., Oduro-Yeboah, C. (2014): Evaluation of the Physical
- 576 and Sensory Characteristics of Bread Produced from Three Varieties of Cassava and Wheat Composite
- 577 Flours. *Food Public Health* 4 (5), 214-222.
- 578
- 579 [45] Hatcher S, Atkins KD and Safari E. Phenotypic aspect of lamb survival in Australian merino Sheep,
- 580 *Journal of Animal Science*, 2009. 87(10) 2781-2790.
- 581
- 582 [46] Hou, G., and Kruk, M. Aslant Noodle Technology, Aslan Noodles Technical Bulletin XX 1998.12, 1-
- 583 10.
- 584
- 585 [47] Tian Q., Streuli M., Saito H., Schlossman S.F. and Anderson P. A polyadenylate binding protein
- 586 localized to the granules of cytolytic lymphocytes induces DNA fragmentation in target
- 587 cells. *Cell*.1991;67:629–639
- 588
- 589 [48]. Tang J., Naitow H., Gardner N.A., Kolesar A., Tang L., Wickner R.B. and Johnson J.E. The
- 590 structural basis of recognition and removal of cellular mRNA 7-methyl G 'caps' by a viral capsid protein: a
- 591 unique viral response to host defense. *Journal of Molecular Recognition*. 2005. 18(2):158-68
- 592
- 593 [49] Benesi I.R. Characteristics Action Of Malawian Cassava Germ Plasm For Diversity, Starch
- 594 Estraction And Its Nature And Modified Properties. *Bioemfontein: Phd Thesis*, Department Of Plant
- 595 Science, University of the Free State, South Africa. 2005. 74-123

596

597 [50] Sriroth k., Chollaku P.R, Chotineeranat S, piyachomkwam K., Oates C. G. Processing of cassava
598 waste for improved biomas utilization. Bioresource Technology 1999. 7: 63-69
599

600 [51] Mishra S, and Rai T, Morphology and Functional Properties of Corn, Potato and Tapioca Starches,
601 Food Hydrocolloid. 2006. 20. 557-592.

602 [52] Hatcher, D. W Bello, G. G. and Anderson, M. J. "Flour particle size, starch damage, and alkali
603 reagent: Impact on uniaxial stress relaxation parameters of yellow alkaline noodles," *Cereal Chemistry*,
604 2009. 86, 361-368
605

606 **APPENDIX**

607