

Original Research Article**Antioxidant and anticancer activities of extracts and compounds isolated from
Terminalia nigrovenulosa plant grown in Vietnam****Abstract**

This study was to isolate and identify antioxidant and anticancer compounds from extracts of bark and leaf of *Terminalia nigrovenulosa*. The EtOAc fraction of bark and n-BuOH fraction of leaf exhibited the highest DPPH (2,2-diphenyl-2-picrylhydrazyl hydrate) radical scavenging activity. Nuclear magnetic resonance (NMR) and mass spectra results showed that gallic acid, ethyl gallate, ellagic acid, catechin and luteolin isolated in EtOAc and n-BuOH fractions were the main components possessed DPPH radical scavenging activity. These fractions and their isolated compounds reduced human fibrosarcoma (HT1080) cell viability in a dose-dependent manner. In addition, these fractions and their isolated compounds significantly increased caspase-3 activity. Therefore, the reduction of cell viability might be due to the induction of apoptosis via caspase-3 pathway. These findings could be useful for the development of new chemotherapeutic agents for the treatment of malignant cancers from *T. nigrovenulosa* extracts and isolated compounds.

Keywords: *T. nigrovenulosa*, DPPH, HT1080 cells, Caspase-3, WST.

1. Introduction

There are about 250 species belonging to the genus *Terminalia* distributed in tropical region of the world. Some of them have been used as a traditional medicine in some Asian

countries. Such as *Terminalia catappa* and *Terminalia chebula* in China for diarrhea [13], *Terminalia bellerica*, *Terminalia chebula* and *Emblica officinalis* (Triphala) in India prescribed for symptoms of inflammation, infection, obesity, fatigue, candida, poor digestion, assimilation, tuberculosis, pneumonia and AIDS [9]. *Terminalia nigrovenulosa* Pierre ex Laness (*T. nigrovenulosa*) is a Vietnamese traditional medicinal plant belonged to *Combretaceae* family and grows wild in deciduous forests in the southern part of Vietnam. The previous researches showed that the extracts of *Terminalia* species possessed a variety of biological activities such as *T. nigrovenulosa* bark and leaf extracts [21]; methanol extracts of *T. chebula* fruits [6], [17], [26]. Tanaka et al. (1986) [29] isolated 12 phenolic compounds from *Terminalia catappa* L. Pfundstein et al. (2010) [22] identified and quantitated 34 phenolic compounds belonged to gallic acid and gallate esters; ellagic acid and its derivatives; chebulic ellagitanins and non-chebulic ellagitanin groups in methanol extracts of fruits of *T. bellerica*, *T. chebula* and *T. horrid*. Of the compounds isolated from *T. chebula* Retz fruit, chebulic acid was the most growth inhibitory against HOS-1 cell lines [27]. Gallic acid and methyl gallate from *T. superba* showed significant inhibition of α -glucosidase activity [30]. Extract of *T. catappa* L inhibited the growth of LLC cells [7]. However, there have been few data on the biological activities of *Terminalia nigrovenulosa* Pierre ex Laness. Therefore, the objective of this study was to isolate and identify the antioxidative and anticancer compounds from *T. nigrovenulosa* extracts in human fibrosarcoma (HT1080) cells.

2. Materials and methods

2.1 Materials and chemicals

Terminalia nigrovenulosa Pierre ex Laness bark and leaf were collected in Chu Yang Sin National Park, Daklak province, Vietnam. After collection, the different parts of fresh plants

were cut and dried at ambient temperature in a room with active ventilation, packed in PE bags and stored at -80°C before use.

Sephadex LH-20 resin (25 – 100 µm bead size), DPPH (2,2-diphenyl-2-picrylhydrazyl hydrate) and Ac-DEVD-AMC (N-acetyl-Asp-Glu-Val-Asp-7-amino-4-methylcoumarin) were obtained from Sigma-Aldrich (St. Louis, MO, USA); methanol d₄ (CD₃OD) and DMSO-d₆, silica gel 63 – 200 µm particle size were obtained from Merk (Darmstadt, Germany); ODS-A gel (3 x 40 cm, 120Å pore size) was purchased from YMC Co. LTD (Kyoto, Japan). Other chemicals were of analytical grade.

2.2 Extraction and isolation of active compounds

Dried leaves (3 kg) and bark (1.5 kg) of *T. nigrovenulosa* were separately extracted with MeOH (20 L (leaves) or 10 L (bark) × 3) at room temperature for 24 h. The combinations of each MeOH extract from bark or leaves were evaporated to produce 300 g extracts. The extracts were then re-suspended in distilled water (3 L for each) and separately partitioned with hexane (2L × 3), chloroform (2L × 3), EtOAc (2L × 3), and n-butanol (2L × 3); detailed in Fig 1.

The ethyl acetate (EtOAc) fraction of bark (112 g) was chromatographed over a silica gel column (10 x 40 cm; 63 – 200 µm particle size) eluting with a chloroform-EtOAc gradient (10 : 0, 8 : 2, 6 : 4, 5 : 5, 3 : 7; each 3 L) to give 5 fractions (F1: 150 mg; F2: 1 000 mg; F3: 12 000 mg; F4: 10 000 mg; F5: 15 250 mg). Fraction 3 was further chromatographed on a silica gel column (5 x 70 cm; 63 – 200 µm particle size) eluting with chloroform-EtOAc-formic acid (4 : 6 : 0.1). Total 20 sub-fractions of 250 ml each were collected and combined on the basis of TLC spraying with 0.1 % of DPPH solution. Sub-fractions 1-10 were then purified on a LH-20 column (4 x 40 cm) eluting with 80% MeOH to yield compound **1** (250 mg). Sub-fractions 13-20 were crystallized in water and then applied to a sephadex LH-20 column (4 x 40 cm) eluting

with 70% MeOH to give compound **2** (**570 mg**). Fractions 4 and 5 were combined and then applied to a ODS-A column (3 x 40 cm, 120Å pore size) eluting with 30% MeOH to give 20 sub-fractions of 200 ml each. Sub-fractions **12-19** were combined on the basis of TLC spraying with 0.1 % of DPPH solution and subsequently chromatographed over a silica gel column (4 x 80 cm, 63 – 200 µm particle size) using chloroform-EtOAc-formic acid (8 : 2 : 0.1) to give compound **4** (**350 mg**). Combination of Sub-fractions 3-8 was further chromatographed on a silica gel column (3 x 80 cm, 63 – 200 µm particle size) using chloroform-EtOAc-formic acid (6 : 4 : 0.1) to yield compound **3** (**250 mg**).

The n-BuOH fraction of leaf (150 g) were applied to a silica gel column (10 x 40 cm; 63 – 200 µm particle size) eluting with a hexane-EtOAc gradient (9 : 1 to 1 : 9) to give 9 fractions of 1.5 L each. The combination of fractions 1 – 4 (**12 300 mg**) was chromatographed on a sephadex LH-20 column (4 x 40 cm) eluting with 70% MeOH to give 10 sub-fractions of 350 mL each. After that, sub-fractions 3 to 4 (**1 720 mg**) was combined on the basis of TLC and then subjected to column chromatography on a silica gel (4 x 100 cm, 63 – 200 µm particle size) eluting with chloroform-EtOAc-formic acid (6 : 4 : 0.1) to give compound **5** (**195 mg**). The combination of sub-fractions **8-10** (**2 350 mg**) was applied to a silica gel column (4 x 100 cm, 63 – 200 µm particle size) eluting with chloroform-EtOAc-formic acid (6 : 4 : 0.1) and then purified in a sephadex LH 20 column (4 x 40 cm) eluting with 80% MeOH to yield compound **1** (**320 mg**). The main compounds in fractions **5-9** (**14 230 mg**) were compound **2** (**320 mg**) and compound **4** (**175 mg**) which were isolated and purified by the method mentioned above.

2.3 Structure analysis

Nuclear magnetic resonance (NMR) spectra were obtained on a Varian Unity Inova 500 and 600-MHz spectrometer (Varian, Walnut Creek, CA, USA) with TMS as the standard at the

Korea Basic Science Institute (KBSI, Gwangju Center, Korea). The mass spectra were measured by a Micromass mass spectrometer (QTOF2).

2.4 Evaluation of antioxidant activity

2.4.1 DPPH radical scavenging activity

Free radical scavenging activity of the extracts against stable DPPH radical (2,2-diphenyl-2-picrylhydrazyl hydrate) was determined by a spectrophotometer using the method described by **Nguyen and Eun** (2011) [21]. Extracted solutions were prepared in a range of concentration (0.075, 0.125, 0.25, 0.5 mg/ml). The solution of DPPH radical in methanol (6×10^{-5} M) was prepared daily before the UV measurements. Three milliliters of this solution was then mixed with 77 μ l of extract solution. The samples were kept in the dark for 15 min at room temperature, after which the decrease in absorption was measured. Absorption of a blank sample containing the same amount of methanol and DPPH radical solution was measured daily. The experiment was carried out in triplicate. Radical scavenging activity was calculated by the following formula:

$$\% \text{ Inhibition} = [(A_B - A_A)/A_B] \times 100,$$

where A_B and A_A stand for absorption of the blank sample ($t=0$ min) and absorption of the tested extract solution ($t=15$ min), respectively. The extract that could scavenge 50% of the DPPH radicals (IC_{50}) was calculated from a plot of scavenging effect versus extract concentration.

2.4.2 Antioxidative activity of fraction in purification procedure

The assay for antioxidative activity of fractions was performed by spraying the 0.1% of DPPH methanolic solution on TLC plate. Each fraction was spotted on TLC plate and developed by a suitable mixture solvent. After spraying the DPPH solution, active fraction will reduce $DPPH^*$, causing a colour change from deep-purple to light yellow.

2.5 Cell culture

Human fibrosarcoma (HT1080) cells were purchased from the American Type Culture Collection (ATCC, USA). The cells were grown in Dulbecco's modified eagle medium (DMEM; Gibco, USA) containing penicillin (100 U/ml), streptomycin (100 µg/ml) (Sigma-Aldrich, St Louis, MO, USA) and 10% fetal bovine serum (FBS; PAA, Canada) at 37°C in 5% CO₂ air. The medium was changed 3 times a week.

2.6 Cell viability assay

Cell viability was measured using Cell Counting Kit-8 (CCK-8; Dojindo, Japan). The procedure is slightly modified method of the instruction of technical manual. Briefly, HT1080 cells were seeded onto a 96-well culture plate at a density of 10⁴ cells/well in 100 µl of DMEM supplemented with 10 % fetal bovine serum and 1% streptomycin-penicillin. After 24 h incubation, the media was replaced with 100 µl of fresh medium and treated with the different concentrations of EtOAc bark and n-BuOH leaf fractions, GA, Cat, EG and luteolin (0 - 100 µg/ml), and EA (0 – 2 µg/ml). Afterward, the cells were incubated for 24h and then 10 µl of CCK-8 was added to each well and incubated for 2 h. Absorbance was measured at 450 nm using a microplate reader.

2.7 Determination of Caspase-3 activity

The caspase-3 assay was based on the hydrolysis of the peptide substrate N-acetyl-Asp-Glu-Val-Asp-7-amino-4-methylcoumarin (Ac-DEVD-AMC) by caspase-3, resulting in the release of the fluorescent 7-amino-4-methylcoumarin (AMC). HT1080 cells were seeded onto a cell culture plate at a density of 2 x 10⁶ cells/plate in DMEM supplemented with 10 % fetal bovine serum and 1% streptomycin-penicillin. After 24 h incubation, the cells were treated with extract fractions and isolated compounds for 24 h. The cells were collected and lysed by 50 µl

137 cold lysate buffer/ 10^6 cells (130 mM NaCl, 10 mM Tris- HCl, 10 mM phosphate buffer pH 7.4,
 138 10 mM. sodium pyrophosphate, 1% Triton X-100). The cell lysate was incubated on ice for 10
 139 min and then centrifuged at $14,000 \times g$ for 5 min. Supernatant was collected and kept on ice.
 140 Addition of 50 μ l of 2X reaction buffer (20 mM HEPES (pH 7.5), 10% glycerol, 2 mM DTT) to
 141 50 μ l of cell lysate in a 96 wells black plate and then mixed with 5 μ l of caspase-3 fluorogenic
 142 substrate (Ac-DEVD-AMC: N-acetyl-Asp-Glu-Val-Asp-7-amino-4-methylcoumarin). The
 143 mixture was incubated at 37°C for 1.5 h. The AMC liberated from Ac-DEVD-AMC was
 144 measured by a spectrofluorometer with an excitation wavelength of 380 nm and an emission
 145 wavelength of 440 nm.

146 **2.8 Statistical analysis**

147 Results were expressed as mean \pm standard deviation of three replicated. The significant
 148 differences between the means of parameters were determined by LSD test ($p < 0.05$) using
 149 Statgraphics centurion XV statistical software.

150

151 **3. Results**

152 **3.1 DPPH radical scavenging activity of extracts and fractions**

153 The DPPH radical scavenging activity of extracts and various fractions of *T. nigrovenulosa*
 154 leaf and bark were presented in **Table 1**. The results indicated that the effective radical
 155 scavengers were concentrated in EtAOc fraction of bark and n-BuOH fraction of leaf. Moreover,
 156 the patterns of EtAOc fraction of bark and n-BuOH fraction of leaf on TLC were simpler than
 157 that of other fractions (data not shown). Therefore, these fractions were used for isolation of
 158 antioxidative compounds.

159 3.2 Yields and structure of isolated compounds

160 The structure of compounds showed in **Fig 2** was elucidated by NMR and MS analysis as
161 follows:

162 **Compound 1:** ellagic acid (EA), yellow (Light), **¹H-NMR** (300 MHz, CD₃OD): δ 7.46 (2H, s,
163 H-13, H-14), 10.58 (2H, s, OH-19, OH-20), 10.795 (2H, s, OH-21, OH-22). **¹³C-NMR** (150
164 MHz, DMSO): δ 107.69 (C-9, C-4); 110.25 (C-8, C-3); 112.32 (C-13, C-14); 136.39 (C-10, C-
165 5); 139.55 (C-11, C-16); 148.12 (C-12, C-15); 159.15 (C-2, C-7). Electrospray ionization-MS
166 (**ESI-MS**) (negative mode) *m/z* 300.9984 [M - H]⁻

167 **Compound 2:** catechin (Cat), brown amorphous powder, **¹H-NMR** (600 MHz, CD₃OD): δ 2.49
168 (1H, *J* = 24, dd, H-3), 2.84 (1H, dd, *J* = 18, H-3), 3.97 (1H, m, *J* = 24, H-2), 4.55 (1H, d, *J* = 6,
169 H-1), 5.85 (1H, s, H-6), 5.92 (1H, s, H-8), 6.71 (1H, dd, *J* = 6, H-6'), 6.75 (1H, d, *J* = 12, H-5'),
170 6.83 (1H, s, H-2'); **¹³C-NMR** (150 MHz, CD₃OD): δ 28.67 (C-3), 68.96 (C-2), 83 (C-1), 95.62
171 (C-8), 96.4 (C-6), 100.94 (C-4), 115.38 (C-2'), 116.2 (C-5'), 120.17 (C-6'), 132.35 (C-1'),
172 146.36 (C-4'), 146.39 (C-3'), 157.05 (C-9), 157.75 (C-7), 157.98 (C-5). **ESI-MS** (negative
173 mode) *m/z* 289.0710 [M-H]⁻.

174 **Compound 3:** gallic acid (GA) white amorphous powder; **¹H-NMR** (600 MHz, CD₃OD) δ 7.05
175 (2H, s, H-2, H-6). **ESIMS** (negative mode) *m/z* 169 [M - H]⁻.

176 **Compound 4:** Ethyl gallate (EG) white amorphous powder, **¹H NMR** (600 MHz, CD₃OD); δ 7.05
177 (2H, s, H-2, H-6); 4.27 (2H, q, *J* = 24, CH₂); 1.34 (3H, t, *J* = 12, CH₃). **ESI-MS** (negative mode) *m/z*
178 197.008 [M - H]⁻

Compound 5: Luteolin, bright yellow amorphous powder; **ESI-MS** (negative mode) m/z 285.021 $[M - H]^-$

The **Table 2** showed that catechin was the highest content in both bark and leaf fractions. Ethyl gallate was found relatively high amount in EtOAc bark fraction, but without in n-BuOH leaf fraction and luteolin was only found in n-BuOH leaf fraction.

3.3. Cell viability

Cell Counting Kit 8 (CCK-8) was used for determination of toxic concentrations of extract fractions and their isolated compounds to HT1080 cancer cells (**Fig 3**). The results showed that EtOAc bark and n-BuOH leaf fractions did not inhibit the viability of HT1080 cells in a concentration range of 0 to 25 $\mu\text{g/mL}$, however, it significantly decreased the viability of cells of about 22.7 to 50.5 % and 13.5 to 51.2 % at a concentration range of 50 to 100 $\mu\text{g/mL}$ of bark and leaf fractions, respectively (**Fig. 3A**). These results were also similar trend in cells treated with Cat, EG and luteolin, with the cell reduction of about 35, 38 and 25 % at the treatment concentration of 100 $\mu\text{g/mL}$, respectively (**Fig. 3A**). GA inhibited the growth of about 27 % of HT1080 cells at a concentration of 25 $\mu\text{g/mL}$ and 64% at 100 $\mu\text{g/mL}$ but without any effect at a concentration ranges of 0 - 12.5 $\mu\text{g/mL}$ (**Fig. 3A**). EA had no significant cytotoxicity in HT1080 cells up to a concentration of 2 $\mu\text{g/mL}$ (**Fig. 3B**). The results indicated that extract fractions and their isolated compounds (GA, EG Cat and luteolin) could reduce the viability of HT1080 cells in dose dependent manner.

3.4 Effect of fractions and isolated compounds on caspase-3 activity in HT1080 cells

Recent work has revealed that caspase-3 plays an important role in the signal transduction pathway leading to apoptosis [23], [31]. Our data mentioned above exhibited that treatment of

HT1080 cells with EtOAc bark and n-BuOH leaf fractions, GA, EG, Cat and luteolin induced cytotoxicity in HT1080 cells (Fig. 4). However, whether this cytotoxicity leads to apoptosis or necrosis, the activity of caspase-3 was measured. The results indicated that the activity of caspase-3 increased together with increasing treatment concentration. At low concentration, there was no significant difference in caspase-3 activity between cells treated with fractions or compounds and control (without any treatment). However, there was a significant increase in the activity of caspase-3 in HT1080 cells treated with EtOAc bark and n-BuOH leaf fractions, EG, Cat, and luteolin at a concentration of 80 µg/mL by about 2.38, 2.69, 2.88, 2.17 and 2.09 fold higher than control cells, respectively. GA made an increase in caspase-3 activity in HT1080 cells by 2.33 fold at the concentration of 30 µg/mL. The data exhibited that the cytotoxic effect of the fractions and compounds (GA, EG, Cat and luteolin) could induce apoptosis in a population of HT1080 cells.

4. Discussions

Several studies have reported that the compounds isolated in *Terminalia* species were ellagic acid, luteolin, gallic acid, ethyl gallate, luteolin, tanic acid, catechins and ellagintannins [27], [5], [22]. The results of present study demonstrated that the main compounds possessed DPPH radical scavenging activity were GA, EG, Cat, luteolin and ellagic acid concentrated in EtOAc bark and n-BuOH leaf fractions of *T. nigrovenulosa* extracts. These compounds have reported to possess high antioxidant activity [15], [25], [12], [19]. Moreover, our data also showed that EtOAc bark and n-BuOH leaf fractions, GA, EG, Cat and luteolin exerted cytotoxicity to HT1080 cells in a type of fractions or compounds with dose dependent manner (Fig. 3A). As shown in Fig. 4, fractions and their isolated compounds increased caspase-3 activity

224 depending on type of fractions or compounds. Therefore, the reduction of cell viability might be
 225 due to the induction of apoptosis via caspase-3 pathway. Previous reports have indicated that
 226 several *Terminalia* species induced cytotoxicity via apoptosis in several cancer cells. *T. chebula*
 227 retz fruit extract induced the cell death by apoptosis at low concentration and necrosis at high
 228 concentration [27]. Acetone extract of Triphala (fruits of *Terminalia bellerica*, *Terminalia*
 229 *chebula* and *Emblica officinalis*) induced S115, MCF-7, PC-3 and DU-145 cells death by
 230 apoptosis [16]. Moreover, phenolic compounds of plants including phenolic acids and flavonoids
 231 are well known as dietary antioxidants. They also exhibit the contrasting pharmacological effects
 232 such as prooxidant toxicity at high doses or present of metals ions [24], [2] and inducing
 233 apoptosis [28], [10]. For instance, gallic acid induced apoptosis in A549 cells via intrinsic
 234 pathway by caspase-3 induction [18] or in fibroblast cells via both intrinsic and extrinsic
 235 apoptotic pathways [8]. The cytotoxic, antioxidant and anticarcinogenic potential of gallic acid
 236 and its derivatives are believed due to their three adjacent hydroxyl groups [14]. Luteolin, a
 237 common flavonoid that exists in many types of plants including fruits, vegetables, and medicinal
 238 herbs, has reported to induce apoptosis in oral squamous cancer cells (OC2) via increasing
 239 caspase-3 and -9 [32]. Furthermore, luteolin increased levels of caspase-3 and the expression of
 240 the pro-apoptotic protein Bax but decreased the expression of the anti-apoptotic protein Bcl-2 in
 241 three human pancreatic carcinoma cell lines [3]. The hydroxyl moieties and 2–3 double bond in
 242 structure features of luteolin that are associated with its biochemical and biological activities [4].
 243 Catechin, a natural flavonoid isolated from several plants, especially in tea, has been shown to
 244 exhibit cytostatic properties in many tumor cells [11], [20]. Alshatwi (2010) [1] indicated that
 245 catechin hydrate suppressed MCF-7 cell proliferation by induction of apoptosis via increasing the
 246 expression of caspase-3, -8, -9 and TP53. Therefore, gallic acid, ethyl gallate, catechin and

luteolin isolated from EtOAc bark and n-BuOH leaf fractions could be responsible for the reduction of cell viability and the increase in the activity of caspase-3 in HT1080 cells.

5. Conclusions

Our study found that the antioxidative compounds concentrated in EtOAc bark and n-BuOH leaf fractions of *T. nigrovenulosa* methanol extracts. Gallic acid, ethyl gallate, ellagic acid, catechin and luteolin were the main antioxidative components isolated from these fractions. The fractions, gallic acid, ethyl gallate, catechin and luteolin could reduce the viability of human fibrosarcoma (HT1080) cells. The reduction of HT1080 cell viability might be due to the induction of apoptosis in HT1080 cells via caspase-3 pathway. Further investigation of anticancer effects on HT1080 cells should continue to figure out the relation between the inhibition of cell growth and apoptosis.

References

1. Alshatwi AA. (2010). Catechin hydrate suppresses MCF-7 proliferation through TP53/Caspase-mediated apoptosis. J Exp Clin Canc Res 29:167-176.
2. Bouayed J, Bohn T. (2010). Exogenous antioxidants—Double-edged swords in cellular redox state. Health beneficial effects at physiologic doses versus deleterious effects at high doses. Oxid Med Cell Longev 3(4); 228-237.
3. Cai X, Lu W, Ye T, Lu M, Wang J, Huo J, Qian S, Wang X, Cao P. (2012). The molecular mechanism of luteolin-induced apoptosis is potentially related to inhibition of angiogenesis in human pancreatic carcinoma cells. Oncol Rep 4:1353-1361

4. Chan TS, Galati G, Pannala AS, Rice-Evans C, O'Brien PJ. (2003). Simultaneous detection of the antioxidant and pro-oxidant activity of dietary polyphenolics in a peroxidase system. *Free Radic Res* 37:787–794.
5. Chen LG, Huang WT, Lee LT, Wang CC. (2009). Ellagitannins from *Terminalia calamansanai* induced apoptosis in HL-60 cells. *Toxicol In Vitro* 23(4):603-609.
6. Cheng HY, Lin TC, Yu KH, Yang CM, Lin CC. (2003). Antioxidant and free radical scavenging activities of *Terminalia chebula*. *Biol Pharm Bull* 26:1331–1335.
7. Chu SC, Yang SF, Liu SJ, Kuo WH, Chang YZ, Hsieh YS. (2007). In vitro and in vivo antimetastatic effects of *Terminalia catappa* L. leaf on lung cancer cells. *Food Chem Toxicol* 45:1194–1201.
8. Chuang CY, Liu HC, Wu LC, Chen CY, Chang JT, Hsu SL. (2010). Gallic acid induces apoptosis of lung fibroblasts via a reactive oxygen species-dependent ataxia telangiectasia mutated-p53 activation pathway. *J Agric Food Chem* 58(5):2943–2951.
9. El-Mekkawey M, Merelhy M. (1995). Inhibitory effects of Egyptian folk medicines on human immunodeficiency virus (HIV) reverse transcriptase. *Chem Pharm Bull* 43:641–648.
10. Galati G, O'Brien P. (2004). Potential toxicity of flavonoids and other dietary phenolics: significance for their chemopreventive and anticancer properties. *Free Radic Biol Med* 37:287-303.
11. Graham HN. (1992). Green tea composition, consumption, and polyphenol chemistry. *Preventive Medicine* 21:334-350.
12. Han DH, Lee MJ, Kim JH. (2006). Antioxidant and apoptosis-inducing activities of ellagic acid. *Anticancer Res* 26(5A):3601-3606.

13. Huang KC. (1993). The Pharmacology of Chinese Herbs. New York, USA: CRC Press
194.
14. Inoue M, Suzuki R, Sakaguchi N, Li Z, Takeda T, Ogihara Y, Jiang BY, Chen Y. (1995).
Selective induction of cell death in cancer cells by gallic acid. Biol Pharm Bull
18(11):1526-1530.
15. Kalaivani T, Rajasekaran C, Mathew L. (2011). Free radical scavenging, cytotoxic, and
hemolytic activities of an active antioxidant compound ethyl gallate from leaf of *Acacia*
Nilotica (L.) Wild. Ex. Delile Subsp. Indica (Benth.) Brenan. J Food Sci 76 (6): 144-149.
16. Kaur S, Michael H, Arora S, Harkonen PL, Kumar S. (2005). The in vitro cytotoxic and
apoptotic activity of Triphala--an Indian herbal drug. J Ethnopharmacol. 97(1):15-20.
17. Lee HS, Jung SH, Yun BS, Lee KW. (2007). Isolation of chebulic acid from *Terminalia*
chebula Retz. and its antioxidant effect in isolated rat hepatocytes. Arch Toxicol 81:211–
218.
18. Maurya DK, Nandakumar N, Asir Devasagayam TP (2011). Anticancer property of
gallic acid in A549, a human lung adenocarcinoma cell line, and possible mechanisms. J
Clin Biochem Nutr 48(1):85–90.
19. Nanjo F, Goto K, Seto R, Suzuki M, Sakai M, Hara Y. (1996). Scavenging effects of tea
catechins and their derivatives on 1,1-diphenyl-2-picrylhydrazyl radical. Free Radic Biol
Med 21(6):895-902.
20. Nakachi K, Suemasu K, Suga K, Takeo T, Imai K, Higashi Y. 1998. Influence of
drinking green tea on breast cancer malignancy among Japanese patients. Jap J Cancer
Res 89:254-261.

21. Nguyen QV, Eun JB. (2011). Antioxidant activity of solvent extracts from Vietnamese medicinal plants. *J Med Plants Res* 5(13):2798-2811.
22. Pfundstein B, El Desouky SK, Hull WE, Haubner R, Erben G, Owen RW. (2010). Polyphenolic compounds in the fruits of Egyptian medicinal plants (*Terminalia bellerica*, *Terminalia chebula* and *Terminalia horrida*): Characterization, quantitation and determination of antioxidant capacities. *Phytochem* 71(10):1132-1148.
23. Porter AG, Janicke RU. (1999). Emerging roles of caspase-3 in apoptosis. *Cell Death Differ* 6:99-104.
24. Raza H, John A. (2005). Green tea polyphenol epigallocatechin-3-gallate differentially modulates oxidative stress in PC12 cell compartments. *Toxicol Appl Pharmacol* 207:212-220.
25. Reddy MK, Gupta SK, Jacob MR, Khan SI, Ferreira D. (2007). Antioxidant, antimalarial and antimicrobial activities of tannin-rich fractions, ellagitannins and phenolic acids from *Punica granatum* L. *Planta Medica* 73:461–467.
26. Saleem A, Ahotupa M, Pihlaja K. (2001). Total phenolics concentration and antioxidant potential of extracts of medicinal plants of Pakistan. *Z Naturforsch* 56:973–978.
27. Saleem A, Husheem M, Härkönen P, Pihlaja K. (2002). Inhibition of cancer cell growth by crude extract and the phenolics of *Terminalia chebula* retz. fruit. *J Ethnopharmacol* 81(3):327-336.
28. Sergediene E, Jonson K, Szymusiak H, Tyrkowska B, Rietjens IMCM, Cenas N. (1999). Prooxidant toxicity of polyphenolic antioxidants to HL-60 cells: description of quantitative structure-activity relationships. *FEBS lett* 462:392-396.

29. Tanaka T, Nonaka GI, Nishioka I. (1986). Tannins and related-compounds isolation and characterization of 4 new hydrolyzable tannins, terflavin-A and terflavin-B, tergallagin and tercatain from the leaf of *Terminalia catappa* L. Chem Pharm Bull 34:1039–1049.
30. Wansi JD, Lallemand MC, Chiozem DD, Toze FAA, Mbaze LM, Naharkhan S, Iqbal MC, Tillequin F, Wandji J, Fomum ZT. (2007). α -Glucosidase inhibitory constituents from stem bark of *Terminalia superba* (Combretaceae). Phytochem 68:2096–2100.
31. Woo M, Hakem R, Soengas M.S, Duncan GS, Shahinian A, Kagi D et al. (1998) Essential contribution of caspase-3/CPP32 to apoptosis and its associated nuclear changes. Genes Dev 12:806–819.
32. Yang SF, Yang WE, Chang HR, Chu SC, Hsieh YS. (2008). Luteolin induces apoptosis oral squamous cancer cells. J Dent Res. 87(4):401-406.

Table 1. The DPPH radical scavenging activity of extracts and fractions of *T. nigrovenulosa* leaf and bark

Extract and fraction	DPPH radical scavenging (IC ₅₀) mg/ml
Bark extract	0.273 ± 0.003 ^a
Leaf extract	0.408 ± 0.006 ^b
EtOAc bark fraction layer	0.162 ± 0.039 ^c
n-BuOH bark fraction layer	0.546 ± 0.014 ^d
Water bark fraction layer	0.698 ± 0.006 ^e
n-BuOH leaf fraction layer	0.258 ± 0.016 ^f
Water leaf fraction layer	0.606 ± 0.007 ^g

Results are means ± SD of triplicate measurements. Different labels (a-g) indicate a significant difference at $P < 0.05$.

351 **Table 2.** Yields of compounds isolated from bark and leaf fractions of *T. nigrovenulosa* (mg/g
 352 dry wt of fractions)

Compounds	Yield (mg/g dry wt of fractions)	
	EtOAc fraction of Bark	n-BuOH fraction of leaf
Ellagic acid	2.332	2.133
Catechin	5.089	2.133
Gallic acid	3.125	1.167
Ethyl gallate	2.323	0
Luteolin	0	1.300

353

354




Figure 1. Extraction fractionation scheme of bark and leaf of *Terminalia nigrovenulosa*




Figure 2. The molecular structure of isolated compounds from *T. nigrovenulosa* fractions





Fig. 3. Effect of EtOAc bark and n-BuOH leaf fractions of *T. nigrovenulosa* extracts, ethyl gallate (EG), catechin (Cat), luteolin, gallic acid (GA) (A) and ellagic acid (B) on viability of HT1080 cells using WST-8 kit. Results are means \pm SD of triplicate measurements. Different labels (a-d) above the bars for the same extract or compound indicate a significant difference at $P < 0.05$.





Fig. 4. Effect of EtOAc bark and n-BuOH leaf fractions, ethyl gallate, catechin and luteolin (A) and gallic acid (B) on activity of caspase-3 in HT1080 cells. Results are means \pm SD of triplicate measurements. Different labels (a-d) above the bars indicate a significant difference at $P < 0.05$.