

**SUSTAINABLE MANAGEMENT OF ROOT KNOT NEMATODE
MELOIDOGYNE INCOGNITA THROUGH ORGANIC
AMENDMENT ON *SOLANUM LYCOPERSICUM* L.**

ABSTRACT

Aims: To evaluate the nematicidal potential of wild spinach powder in combination with fresh chopped leaves of different plants viz., Indian mallow, Mexican poppy, Ivy gourd, Trailing eclipta, Wild eggplant and Black pigweed against root knot nematode, *Meloidogyne incognita* on plant growth characters of tomato cv. K-21.

Place and Duration of Study: A glasshouse pot experiment was carried out in the Department of Botany, Aligarh Muslim University, Aligarh.

Methodology: A mixture of soil and organic manure was prepared in the ratio 3:1. The pots were treated with fresh chopped leaves of different plants viz., Indian mallow, Mexican poppy, Ivy gourd, Trailing eclipta, Wild eggplant and Black pig weed applied 50g of fresh chopped leaves of plants were applied combined with seed powder 10g of seed powder of Black nightshade".

Results: Combined application of wild spinach powder along with the fresh chopped leaves of all the plants suppressed pathogenic effect of nematode and thereby resulted in significant reduction in root-knot nematode infestation and population density of *M. incognita* in soil. The highest reduction in root-knot nematode infestation was reported in plants employed with wild spinach powder@10g combined with Mexican poppy leaves@50g. Plants applied wild spinach powder @10g combined with Mexican poppy leaves@50g were the best followed by Trailing eclipta@50g, Wild eggplant@50g, Black pigweed@50g, Indian mallow@50g, Ivy gourd@50g in the descending order. However the lowest reduction was recorded with the application of wild spinach powder@10g plus fresh chopped Ivy gourd@50g wild spinach. Organic additives of wild spinach powder along with the fresh chopped leaves would work for sustainable management by increasing the nutrient status of the soil which further enhances the plant growth.

Conclusion: The application of Mexican poppy, Trailing eclipta and Wild eggplant in combination with wild spinach powder are quite efficient alternatives of chemical nematicides for the *M. incognita* management and yield enhancement. Although the utilization of chopped

leaves of Indian mallow and Ivy gourd along with wild spinach powder is not potentially active replacement of synthetic nematicides however further characterization is needed.

Key words: Tomato, Chopped leaves, Powder, *Meloidogyne incognita*, Sustainable management.

Introduction

Tomato (*Solanum lycopersicum* L. Family- Solanaceae) is the most important vegetable crop widely used throughout the world. Tomatoes are consumed in the form of juice, paste, ketchup, soup etc. Indian contribution to the annual world production was 10.26 kg with an area of 5.72 kg in 2006 (Mane, *et al.*, 2010). Root knot nematode was found constantly associated with the root of tomato where population of nematode was influenced by clay and silt percentage (Asif *et al.*, 2015). The *Meloidogyne incognita* are sedentary endoparasites and are among the most damaging agricultural pests attacking a wide range of crops (Sahebani and Hadavi, 2008), particularly in vegetables, causing dramatic yield losses mainly in tropical and sub-tropical agriculture (Kiewnick and Sikora, 2006). Plant parasitic nematodes attack plants and caused significant damages and yield losses in crops (Luc *et al.*, 2005). In the absence of effective control *M. incognita* can cause total crop failure (Sikora and Fernandez, 2005). Current management of nematodes includes plant resistance, crop rotation (Chitwood, 2002) and biocontrol agents (Mukhtar and Pervaz, 2003; Rahoo *et al.*, 2011). Most important way of nematode control is mainly the nematicides. Most of the nematicides are banned due their toxic nature and carcinogenic effect on the fauna; beside this they disturbed the ecological equilibrium of the soil. Thus the development of alternative control strategies and long-term integrated approaches is urgently needed in order to replace chemical nematicides (Martin, 2003; Ganai *et al.*, 2013; Hussain *et al.*, 2011b; Asif *et al.*, 2013).

The incorporation of organic material (plant parts and plant products) into the soil reduces *M. incognita* densities, resulting in an increase in tomato yield (Muller and Gooch, 1982 Ansari *et al.*, 2016). Various medicinal and antagonistic plants have been found effective for the control of root-knot nematodes (Ahmad *et al.*, 2004; Ibrahim *et al.*, 2006; Abo-Elyousr *et al.*, 2010). Therefore, this study was conducted with the

purpose to evaluate the efficacy of Wild spinach powder in combination with the fresh chopped leaves of different plant species against *Meloidogyne incognita* on tomato.

Material and Methods

SITE FOR THE EXPERIMENT

A mixture of soil and organic manure was prepared in the ratio 3:1. Glasshouse pot culture experiment was carried out in Department of Botany, Aligarh Muslim University, Aligarh. It has average temperature ranges from 28–38 °C (82–100 °F) with the scanty rainfall of 650 to 750 mm is recorded throughout the year.

TEST PLANT AND PATHOGEN

Tomato (*Solanum lycopersicum* L.cv.K-21, Family Solanaceae) was selected as a test plant and *Meloidogyne incognita* (Kofoid and White, 1919; Chitwood 1949), was chosen as test pathogen to evaluate the effect of various organic soil amendments on the management of *M. incognita*.

Seed powder of Wild spinach (*Solanum nigrum* L., Family Solanaceae), Fresh chopped leaves of Mexican poppy (*Argemone mexicana* L., Family Papaveraceae), Trailing eclipta (*Eclipta alba* L., Family Asteraceae), Wild eggplant (*Solanum xanthocarpum* Schrad. & Wendl, Family Solanaceae), Black pig weed (*Trianthema portulacastrum* L., Family Aizoaceae), Indian mallow (*Abutilon indicum* L. (Sweet), Family Malvaceae), Ivy gourd (*Coccinia grandis* (L.) Voigt Family Cucurbitaceae) were taken for the study from the campus Aligarh Muslim University, Aligarh. Collected leaves were washed in running tap water to remove dust particles and then processed for chopping with the help of knife.

COLLECTION OF INFECTED ROOT

M. incognita infected roots were collected from Mathura road, Aligarh, UP, India. Infected roots of the plants gently removed from the soil and kept in polythene bags and then labeled. Further all these samples were brought to the laboratory for the examination.

SOURCE OF MELOIDOGYNE INCOGNITA JUVENILES

Juveniles of *Meloidogyne incognita* (Kofoed and White, 1919; Chitwood, 1949) were prepared from a pure culture that was previously cultured by eggmasses and propagated on eggplant (*Solanum melongena* L.) in the glasshouse of Section of Plant Pathology and Plant Nematology, Department of Botany, Aligarh Muslim University, Aligarh, India. Surface attached eggmasses were detached by using sterilized forceps from the infected roots. These eggmasses were placed in 15 mesh sieves (8 cm in diameter) having crossed layer of tissue paper. These were kept in petridishes full of water so that eggmasses may remain in contact with water. These petridishes were then incubated at $28 \pm 2^{\circ}\text{C}$ for hatching and for freshly hatched second stage juveniles (J2) of *M. incognita*.

MELOIDOGYNE SPECIES IDENTIFICATION

Root knot nematode *Meloidogyne* spp. was identified on the basis of perineal pattern from infected root samples and were examined under stereomicroscope according to Eisenback *et al.*, 1981.

MAINTENANCE OF SEEDLINGS

The seeds of tomato cv. K-21 were surface sterilized in 0.01% HgCl_2 for three minutes and then rinsed with Double Distilled Water (DDW) three times. For nursery preparation seeds were sown in autoclaved clay pots(30cm diameter) along with the soil. Three weeks after germination of proper seedlings of tomato cv. K-21 were transplanted to each 15 cm diameter clay pots filled with 1 kg autoclaved soil. These pots were treated with fresh chopped leaves of different plants viz., Indian mallow, Mexican poppy, Ivy gourd, Trailing eclipta, Wild eggplant and Black pig weed applied 50g of fresh chopped leaves of different plant combined with seed powder of Black nightshade @ 10g per pot. The pots were watered regularly for proper decomposition of the organic additives for two weeks. As the seedling get established each of them was inoculated with 1500 hatched second stage juveniles (J2) of *Meloidogyne incognita*. Each treatment was replicated four times.

INOCULATION TECHNIQUE

Inoculation was done by making 3 holes in the pots soil nearby to the roots at the same distance in the manner so that root don't get damage. Then requisite amount of

suspension having necessary number of second stage juveniles was procured into the holes and then covered them with the soil.

OBSERVATION

After three months of the inoculation, roots of tomato cv. K- 21 were uprooted carefully from the pots and were washed in running tap water to wash off the soil particles adhered with the root is removed. The water present in the plants was eliminated by pressing them in between the blotting sheets. The plant growth parameters including the length (cm), fresh and dry weight (g) of shoot and root, yield/plant (g) were measured. Biochemical and pathological parameters including percent pollen fertility, chlorophyll content (mg/g), carotenoid content (mg/g), eggmasses /root, eggs/eggmass, nematode population (250g) and root knot index (RKI) were also examined respectively. The number of plants with galled root system and the root-knot index were evaluated on a 0-5 scale (0 = no galling; 1 = 1-2 galling; 2 = 3-10 galling; 3 = 11-30 galling; 4 = 31-100 galling; and 5 = more than 100 galling per root system according to Taylor and Sasser,1978). The nematode populations present in the soil was estimated by Cobb's sieving and decanting technique followed by modified Baermann's funnel technique by processing 250 g of soil sample.

STATISTICAL ANALYSIS

The data presented in the tables was analyzed by one-way analysis of variance (ANOVA) using SPSS 17.00 software (SPSS Inc., Chicago, IL, USA). Least Significant Differences (LSD) were calculated at $P= 0.05$ to test for significant differences between the treatment means.

RESULTS

The present experiment was carried out under glasshouse conditions to test the effectiveness and nematostatic potential of organic soil amendments with chopped leaves of Indian mallow, Mexican poppy, Ivy gourd, Trailing eclipta, Wild eggplant and Black pig weed applied @ 50g combined with seed powder of Black nightshade @ 10g per pot on the root-knot development caused by *Meloidogyne incognita* and plant growth character of tomato cv. K-21. It was noted that there was significant improvement in plant growth parameters in all the treatments but chopped leaves of *Argemone mexicana* combined with Black nightshade seed powder elucidate highest

improvement in plant growth parameters and maximum reduction in nematode population and root-knot indices. The total length of tomato cv. K -21 was found highest as 53.5 cm when tomato plants treated with leaves of Mexican poppy followed by 51.7, 50.0, 48.2, 46.4 and 44.7 cm in Wild eggplant, Black pig weed, Indian mallow and Ivy gourd respectively as compared to 36.5 cm in untreated inoculated and 78.2 cm in untreated uninoculated control plants. The total fresh weight of tomato cv. K -21 was found highest as 41.0g when tomato plants treated with leaves of Mexican poppy followed by 39.6, 38.4, 37.2, 36.0 and 35.2g in Trailing eclipta, Wild eggplant, Black pig weed, Indian mallow and Ivy gourd respectively as compared to 36.5g in untreated inoculated and 78.2g in untreated uninoculated control plants. The total dry weight of tomato cv. K -21 was found highest as 14.2g when tomato plants treated with leaves of Mexican poppy followed by 13.5, 12.9, 12.3, 11.5 and 11.0g in Trailing eclipta, Wild eggplant, Black pig weed, , Indian mallow and Ivy gourd respectively as compared to 8.5g in untreated inoculated and 24.0g in untreated uninoculated control plants (Table-1a).

The total chlorophyll and carotenoid content was significantly increased. The chlorophyll and carotenoid content was 1.73 & 0.495 mg/g, 1.68 & 0.474 mg/g, 1.62 & 0.446 mg/g, 1.55 & 0.418mg/g, 1.51 & 0.400 mg/g and 1.45 & 0.390 in Mexican poppy, Trailing eclipta, Wild eggplant, Black pig weed, Indian mallow and Ivy gourd respectively as compared to 1.12 & 0.252mg/g in untreated inoculated and 2.78 & 0.880 mg/g in untreated uninoculated control. The percent pollen fertility was significantly increased in all the treatments. Maximum pollen fertility (67.0%) in treated plants, when the tomato was treated with Mexican poppy it was followed by as 65.6, 64.0, 63.5, 62.0 and 61.0 % in Trailing eclipta, Wild eggplant, Black pig weed, Indian mallow and Ivy gourd respectively as against 47.0 % in untreated inoculated and 89.5% in untreated uninoculated control. The chopped leaves along with the seed powder significantly increased the yield. Maximum yield (228g) was observed in treated one, when the tomato was treated with Mexican poppy it was followed by as 200, 186, 167.0, 160.0 and 144.0g in Trailing eclipta, Wild eggplant, Black pig weed, Indian mallow and Ivy gourd respectively as against 112.0g in untreated inoculated and 355.0g in untreated uninoculated control (Table-1b). Significant reduction was observed in eggmasses,

187 eggs and nematode population of *M. incognita* when tomato was treated with chopped
188 leaves in combination with seed powder of different plant species. Highest reduction in
189 eggmasses, eggs and nematode population 114.0, 188.0 & 1042 was observed in
190 Mexican poppy leaves followed by 118.0, 193.0 & 1069 in Trailing Eclipta, 123.0, 197.0
191 & 1100 in Wild eggplant, 128, 202 & 1142 in Black pig weed, 132, 209 & 1170 in Indian
192 mallow and 140, 215 & 1204 in *C. grandis* respectively as against 148.0, 254.0 &
193 1558.0 in untreated inoculated control. Fresh chopped leaves in combination with seed
194 powder significantly reduced the root knot development. The highest reduction in the
195 root-knot development was detected when tomato plants were treated with leaves of
196 Mexican poppy and root knot index was recorded as 2.0 followed by 2.2, 2.3, 2.5, 2.8
197 and 3.0 in Trailing eclipta, Wild eggplant, Black pig weed, Indian mallow and Ivy gourd
198 respectively. In untreated inoculated pots the root-knot development was highest (5.0)
199 (Table-1b).

200 **Table-1a. Effect of Chopped leaves of different plant species in combination with leaf powder of Wild spinach on growth**
 201 **of tomato cv.K-21 in relation to root-knot development caused by *Meloidogyne incognita* in pots.**

Treatment	Length(cm)			Weight(g)					
	Shoot	Root	Total	Fresh			Dry		
				Shoot	Root	Total	Shoot	Root	Total
Indian mallow	31.4ef	15.0cd	46.4de	25.7e	10.3bc	36.0ef	8.3de	3.2cd	11.5ef
Mexican poppy	36. b	17.3b	53.5b	29.0b	12.0b	41.0b	10.0b	4.2b	14.2b
Ivy gourd	30.5f	14.2d	44.7 e	25.2e	10.0c	35.2f	8.0e	3.0d	11.0f
Trailing eclipta	35.0bc	16.7bc	51.7bc	28.0bc	11.6 b	39.6bc	9.5bc	4.0b	13.5bc
Wild eggplant	33.8cd	16.2bc	50.0bcd	27.2cd	11.2b	38.4cd	9.2bc	3.7bc	12.9cd
Black pig weed	32.4de	15.8bc	48.2cde	26.de	10.7bc	37.2 de	8.8cd	3.5bc	12.3de
UUC	53.6a	24.6a	78.2 a	52.0a	20.0a	72.0a	18.0 a	6.0a	24.0a
UIC	24.7g	11.8e	36.5f	20.0f	8.0d	28.0g	7.1f	1.4g	8.5g

202 Each value is the mean of four replicates; Initial inoculum = 1500 (J₂) of *Meloidogyne incognita* per pot; Means in each column
 203 followed by same letter are not significantly different according to Duncan's Multiple Range Test (DMRT) at $P \leq 0.05$; UUC-
 204 Untreated Uninoculated Control; UIC- Untreated Inoculated Control.

205

206

207 **Table1b.Effect of Chopped leaves of different plant species in combination with leaf powder of Wild spinach on**
 208 **tomato cv. K-21 In relation to root knot development and multiplication of *Meloidogyne incognita* in pots.**

Treatment	Chlorophyll content (mg/g)	Carotenoid content (mg/g)	Pollen fertility (%)	Yield /plant(g)	Eggmasses /root	Eggs/ Eggmass	Nematode Population (250g)	Root-knot Index
Indian mallow	1.51cd	.400 e	62.0c	160ef	132bc	209bc	1170 c	2.8bc
Mexican poppy	1.73 b	.495 b	67.0b	228b	114e	188e	1042 g	2.0d
Ivy gourd	1.45 d	.390f	61.0c	144g	140 b	215 b	1204 b	3.0b
Trailing eclipta	1.68b	.474 c	65.6b	200 c	118de	193de	1069 f	2.2cd
Wild eggplant	1.62bc	.446 d	64.0bc	186 d	123cd	197de	1100e	2.3cd
Black pig weed	1.55cd	.418 e	63.5bc	167e	128c	202cd	1142 d	2.5bcd
UIC	1.12e	.252g	47.0f	112.0g	175 a	266 a	1608 a	5.0 a
UUC	2.78 a	.880 a	89.5 a	355 a	0 f	0 f	0 g	0 e

209 Each value is the mean of four replicates; Initial inoculum = 1500 (J_2) of *Meloidogyne incognita* per pot; Means in each column f
 210llowed by same letter are not significantly different according to Duncan's Multiple Range Test (DMRT) at $P \leq 0.05$; UUC- U n tr
 211eated Uninoculated Control; UIC- Untreated Inoculated Control.

DISCUSSION

Results presented in the study clearly indicated that all the treatment were found to be potent for the tomato plants as they promoted of growth by enhancing the shoot and root growth of the plants thereby results in decreasing the nematode infestation. Among all the treatments Mexican poppy combined with wild spinach leaf powder recorded highest improvement in plant growth and maximum reduction in the nematode multiplication and root knot indices. Other treatments viz., Trailing eclipta, Wild eggplant, Black pig weed, Indian mallow and Ivy gourd combined with spinach leaf powder also showed significant enhancement in plant growth of tomato and reduction in the population of root-knot nematode, *Meloidogyne incognita*. A beneficial relationship was detected in the combined treatment of fresh chopped leaves plus wild spinach leaf powder with the tomato plants in terms of morphological and biochemical parameters as compared to inoculated control. The potential of all treatments to inhibit nematode activity resulted in increased number of fruits and yield of tomato. This beneficial relationship and improvement in plant growth parameters may be due to release of allochemicals and nutrients on the decomposition of the amendment that enhance the beneficial microflora of the soil rhizosphere that act as promoter of growth activity. While the reduction in nematode infestation may be due to release of some toxic chemicals which either disrupt the life cycle of the nematode or lesser down the infectivity of the juvenile. Yujil *et al.* (2002) reported that incorporation of rock fleabane plant (*Inula viscosa*) powder at a concentration of 0.1 % in soil reduced second stage juveniles of root- knot nematodes (*M. javanica*). The mechanisms of plant extract action may include denaturing and degrading of proteins, inhibition of enzymes and interfering with the electron flow in respiratory chain or with ADP phosphorylation (Konstantopoulou *et al.*, 1994). Sensitivity of plant-parasitic nematodes varies from plant to plant. Generally, J2s of *Meloidogyne* spp. are very sensitive to plant-derived nematicides (Oka *et al.*, 2001).

Organic farming is an area of great concern and interest for growing safe and healthy food in sustainable manner without disturbing the ecological diversity of the soil and maintaining the environmental and soil behaviour. The outcome of the study state that the application of Mexican poppy, Trailing eclipta and Wild eggplant in combination

243 with wild spinach powder is quite efficient alternative of chemical nematicides for the
244 nematode management and enhancing the yield. The leaf powder of *I. viscosa*, at a
245 lower concentration, showed nematicidal activity against *Pratylenchus mediterraneus*,
246 although this activity was much lower than that against *M. javanica* J2 (Oka *et al.*,
247 2001). Saxena (1989) reported that neem seed cake improved growth of tomato and
248 reduced the reproduction factor and root galling of nematodes. Some herbal powders
249 and their aqueous extracts result in increasing plant growth and reduction in the
250 infection rate (Moosavi, 2012). The observation of the findings are in conformity with
251 Asif *et al.*, 2013; Ganai *et al.*, 2013, Ansari *et al.*, 2016. Addition of soil amendments
252 results in a considerable increase in the liberation of CO₂ through the saprophytic
253 activities of soil saprophytes which can suppress the activities of pathogen (Papavizas
254 and Davey, 1992). The use of phytochemical compounds and natural plant extracts has
255 substantially reduced the number of galls and the extent of nematode reproduction on
256 tomato plants (Naz *et al.*, 2013a, b). More specifically, the decomposition of organic
257 residues from some oily plant residues, such as cottonseed meal, or from meal from
258 certain types of mustard have been reported as releasing toxic ammonia, organic acids
259 and other compounds as a byproduct that can kill nematodes (Oka, 2010; McSorley,
260 2011; Thoden *et al.*, 2011). The use of plants as nematicidal or nematostatic products
261 has been regarded as effective, economical and eco-friendly by numerous researchers
262 (Chitwood, 2002). Amendments not only change physical and chemical soil properties,
263 but also support a wide variety of antagonistic microorganisms like fungi, bacteria, etc.
264 (Jaffee *et al.*, 1998; Timm *et al.*, 2001) that through the competition and parasitism may
265 reduce the population of nematode. Usage of *A. indica* (neem) caused maximum
266 reductions in the number of galls, egg masses and reproduction factor (Rf) of the
267 nematode and increases all growth parameters of okra (Usha and Sundararaj, 1995;
268 Hussain *et al.*, 2011). The plant parts/products are safer, friendly than synthetic
269 nematicides and may be economic and easily available. Moreover, they are also
270 responsible to increase in soil nutrient status, soil loosening and texture, and to maintain
271 the soil fertility. The present study also elucidate that the reduction in nematode
272 population could be due to the decomposition of plant material in soil and release of
273 secondary metabolites.

CONCLUSION

The application of Mexican poppy, Trailing eclipta and Wild eggplant in combination with wild spinach powder is quite efficient alternative of chemical nematicides for the nematode management and enhancing the yield. Although the utilization of chopped leaves of Indian mallow and Ivy gourd along with wild spinach powder is not potentially active replacement of synthetic nematicide however for this further characterization is needed. Exaggeration of the fertility of the soil and enhancement of microbial diversity of the rhizosphere may be the results of the suppression of root knot nematode infestation due to the release of nematotoxic chemical constituents after decomposition thereby ameliorate the plant growth character of the crop. All the plants explored during the course of studies are locally available at large scale without any toxicological impact on the environmental flora and fauna. It can work for fulfill the demand of global food production through sustainable nematode management by the involvement of organic farming system.

REFERENCES

- Ansari T., M. Asif and M.A. Siddiqui, 2016. Potential of botanicals for root knot nematode management on tomato. Lambert academic publishing. ISBN NO. 9783659910920: p: 1-121
- Abo-Elyousr, K.A., Z. Khan, M. El-Morsi Award, and M.F. Abedel-Monei, 2010. Evaluation of plant extracts and *Pseudomonas* spp. for control of Root-knot nematode, *Meloidogyne incognita* on tomato *Nematropica*, 40.pp. 289–299.
- Agnihotri, N.P., S. Walia and V.T. Gajbhyie, 1999. Green pesticides, protection and safety evaluation. Indian Agricultural Research Institute, New Delhi.
- Ahmad, M.S., T. Mukhtar, and R. Ahmad, 2004. Some studies on the control of Citrus nematode (*Tylenchulus semipenetrans*) by leaf extracts of three plants and their effects on plant growth variables *Asian J. Plant Sci.*, 3.pp. 544–548.
- Alam, M.M., S.K Saxena, and A.M. Khan, 1978. Suitability of crops to certain ectoparasitic nematodes. *Acta Botanica Indica*, 6: (supplement) 205-208.
- Andrés, M. F., A., Gonzáles-Coloma, J., Sanz, J. Burillo And P. Sainz, 2012. Nematicidal activity of essential oils: a review. *Phytochemistry Reviews*, **11**: 371-390.

304 Asif, M., K., Parihar, B., Rehman, M.A., Ganai, A. Usman, and M.A. Siddiqui, 2014.
 305 Bio- efficacy of some leaf extracts on the inhibition of egg hatching and mortality of
 306 *Meloidogyne incognita*. *Archives of Phytopathology and Plant Protection*, 47:8, 1015-
 307 1021.

308 Asif, M., B., Rehman, K., Parihar, M.A Ganai, and M.A. Siddiqui, 2015. Effect of
 309 Various Physico-Chemical Factors on the Incidence of Root Knot Nematode
 310 *Meloidogyne* spp. Infesting Tomato in District Aligarh (Uttar Pradesh) India. *Journal of*
 311 *Plant Sciences* 10 (6): 234-243, 2015

312 Chitwood, D.J. 2002. Phytochemical based strategies for nematode control. *Annu Rev.*
 313 *Phytopath.*, 40: 221–249.

314 Chitwood, D.J. 2002. Phytochemicals based strategies for nematode control *Annu. Rev.*
 315 *Phytopathol.*, 40.pp. 221–249.

316 Claudius-Cole, A.O., A.E. Aminu, and B. Fawole, 2010. Evaluation of plant extracts in
 317 the management of root-knot nematode (*Meloidogyne incognita*) on cowpea (*Vigna*
 318 *unguiculata* (L.) Walp.) *Mycopathologia*. 8:53–60.

319 Eisenback, J.D., H ., Hirschmann ,J.N ., Sasser, and A.C. Triantaphyllou, 1981. A more
 320 complete characterization of the four most common species of root-knot
 321 nematodes (*Meloidogyne* spp.) with pictorial Key. IMP publication Raleigh, N.C.U.S.A

322 Ganai, M.A., B., Rehman, K., Parihar, M. Asif, and M.A. Siddiqui, 2013.
 323 Phytotherapeutic approach for the management of *Meloidogyne incognita* affecting
 324 *Abelmoschus esculentus* (L.) Moench. *Archives of Phytopathology and Plant*
 325 *Protection*, 47:15, 1797- 1805.

326 Ganai, M.A., B., Rehman, K., Parihar, M. Asif, and M.A. Siddiqui, 2013.
 327 Phytotherapeutic approach for the management of *Meloidogyne incognita* affecting
 328 *Abelmoschus esculentus* (L.) Moench. *Archives of Phytopathology and Plant*
 329 *Protection*, 47:15, 1797- 1805.

330 Grewal, P.S, 1989. Effects of leaf-matter incorporation on *Aphelenchoides composticola*
 331 (Nematoda), mycofloral composition, mushroom compost quality and yield of *Agaricus*
 332 *bisporus* *Ann. Appl. Biol.*, 115. pp. 299–312.

333 Hussain, M. A., T. Mukhtar, and M. Z. Kayani, 2011. Efficacy evaluation of *Azadirachta*
 334 *indica*, *Calotropis procera*, *Datura stramonium* and *Tagetes erecta* against root-knot
 335 nematodes *Meloidogyne incognita*. *Pak. J. Bot.*, 43: 197-204.
 336 Hussain, M.A., T. Mukhtar and M.Z. Kayani, 2011. Efficacy evaluation of *Azadirachta*
 337 *indica*, *Calotropis procera*, *Datura stramonium* and *Tagetes erecta* against root-knot
 338 nematodes *Meloidogyne incognita* *Pak. J. Bot. (Special Issue)*, 43 .pp. 197–204.
 339 Ibrahim, S.K., A.F. Traboulsi and S. El-Haj, 2006. Effect of essential oils and plant
 340 extracts on hatching, migration and mortality of *Meloidogyne incognita* *Phytopathol.*
 341 *Medit.*, 45.pp. 238–246.
 342 Jaffee, B.A., H. Ferris and K. M. Scrow, 1998. Nematode-trapping fungi in organic and
 343 conventional cropping systems. *Phytopathol.*, 88:344-350.
 344 Kashyap, N.P., R.M., Bhagat, D.C. Sharma and S.M. Suri, 1992. Efficacy of some
 345 useful plant leaves for the control of potato tuber moth, *Phthorimaea operculella* Zell. *in*
 346 *stores J. Entomol. Res.*, 16.pp. 223–227.
 347 Khan, A.M., M.M. Alam and S.K. Saxena, 1974. Mechanism of control of plant parasitic
 348 nematode as a result of the application of oil cakes to the soil. *Indian J. Nematol.*, 4: 93-
 349 96.
 350 Kiewnick, S. and R.A. Sikora, 2006. Biological control of the root-knot
 351 nematode *Meloidogyne incognita* by *Paecilomyces lilacinus* strain 251. *Biological*
 352 *Control*. 38:179–187.
 353 Konstantopoulou , I., L., Vassilopoulou, P.P. Mawogantisi and G. Scouras, 1994.
 354 Insecticidal effect of essential oils: a study of essential oils extracted from eleven
 355 Greek aromatic plants on *Drosophila auroria*. *Experientia*. 48:616–619.
 356 Luc, M., R.A. Sikora, and J. Bridge, 2005. Plant parasitic nematodes in subtropical and
 357 tropical agriculture. Luc M, Sikora RA, Bridge J (Editors), CABI Wallingford, UK. 871 pp.
 358 Mane, A.V., B.A. Karadge and J.S. Samant, 2010. Salinity induced changes in
 359 photosynthetic pigments and polyphenols of *Cymbopogon nardus* (L.) Rendle. *J. Chem.*
 360 *Pharm. Res.*, 2, 338-347.
 361 Martin, F.N, 2003. Development of alternative strategies for management of soil borne
 362 pathogens currently controlled with methyl bromide. *Annu. Rev. Phytopathol.* 41, 325
 363 350.

364 McSorley, R, 2011. Overview of organic amendments for management of plant-parasitic
 365 nematodes, with case studies from Florida. *J. Nematol.*, 43: 69–81.

366 Moosavi, M.R, 2012. Nematicidal effect of some herbal powders and their aqueous
 367 extracts against *Meloidogyne javanica*. *Nematropica*, 2012; 42: 48-56.

368 Mukhtar, T. and I. Pervaz, 2003. *In vitro* evaluation of ovicidal and larvicidal effects of
 369 culture filtrate of *Verticillium chlamydosporium* against *Meloidogyne javanica* *Int. J. Agri.*
 370 *Biol.*, 5 .pp. 576–579.

371 Muller, R. and P.S. Gooch, 1982. Organic amendments in nematode control. An
 372 examination of the literature. *Nematropica*, 12: 319-326.

373 Naz, I., J.E., Palomares-Rius, V., Saifullah, M.R. Blok, S. Ali Khan and S. Ali, 2013a.
 374 *In vitro* and in planta nematicidal activity of *Fumaria parviflora* (Fumariaceae) against
 375 the southern root-knot nematode *Meloidogyne incognita*. *Plant. Pathol.*, 62: 943–962.

376 Naz, I., Saifullah and M.R. Khan, 2013b. Nematicidal activity of nonacosane-10-ol and
 377 23a- homostigmast-5-en-3 β -ol isolated from the roots of *Fumaria parviflora*
 378 (Fumariaceae). *J. Agric. Food. Chem.*, 61: 5689–5695.

379 Oka Y, B, Ben-Daniel Y, Cohen, 2001. Nematicidal activity of powder and extracts of
 380 *Inula viscosa*. *Nematology* 3, 735–42.

381 Oka, Y., 2010. Mechanisms of nematode suppression by organic soil amendments – a
 382 review. *Appl. Soil. Ecol.*, 44: 101–115.

383 Pandey, R, 2000. Additive effect of three organic materials and nematicides on the
 384 reproduction of *M. incognita* and yield of *Mentha arvensis*. *Nematropica*, 30 (2): 154 –
 385 160.

386 Papavizas, G. C. and C. B. Davey, 1992. Activity of *Rhizoctonia* in soil as affected by
 387 carbon dioxide. *Phytopathology*, 52: 759-766.

388 Rahoo, A.M., T., Mukhtar, S.R. Gowen, and B. Pembroke, 2001. Virulence of
 389 entomopathogenic bacteria *Xenorhabdus bovienii* and *Photorhabdus luminescens*
 390 against *Galleria mellonella* larvae *Pak. J. Zool.*, 43 .pp. 543–548.

391 Sahebani, N. and Hadavi, N, 2008. Biological control of the root-knot
 392 nematode *Meloidogyne javanica* by *Trichoderma harizianum*. *Soil Biology and*
 393 *Biochemistry*. 40:2016–2020.

394 Saxena, R.C, 1989. Insecticides from neem. In: Arnason JT, Philogene BJR, Morand P,
 395 editors. Proceedings of the ACS Symposium on Insecticides of Plant Origin. ACS
 396 Symposium Series 387, Washington, DC: American Chemical Society; p. 110–135.

397 Sayre, R. M, 1980. Promising organism for bio-control of nematodes. *Plant Dis.*, 64:
 398 527-532.

399 Sikora, R.A. and E. Fernandez, 2005. Nematode Parasites of Vegetables. In: Plant
 400 Parasitic Nematodes in Subtropical and Tropical Agriculture. Luc M, Sikora RA, Bridge J
 401 (Editors) 2nd Edition, CABI Wallingford, UK. pp 319-392.

402 Thoden, T.C., G.W. Korthals and A.J. Termorshuizen, 2011. Organic amendments and
 403 their influences on plant-parasitic and free-living nematodes: a promising method for
 404 nematode management. *Nematology*, 13: 133–153.

405 Timm, L.; D. Pearson and B. Jaffee, 2001. Nematode-trapping fungi in conventionally
 406 and organically managed corn–tomato rotations. *Mycologia*, 93: 25-29.

407 Usha, M. and P. Sundararaj, 1995. Efficacy of some new soil amendments for the
 408 control of the lesion nematode in sugarcane. *Nematol. Medit.*, 23: 321-323.

409 Yuji Oka., S., Nacar, Y., E. Putievsky, U., Ravid, Z. Yaniv, and Spiegel, 2010.
 410 Nematicidal activity of Essential oils and their components against the root-knot
 411 nematode. *Phytopathology.*, 90 (7): 710-715