

Abstract

*In the present study, an agricultural soil artificially amended with different concentrations of palm oil mill effluent was analysed to determine its impact on the soil microflora and other properties such as pH, organic matter, electrical conductivity, and mineral elements. In the polluted soils, the total heterotrophic bacteria, total heterotrophic fungi and total heterotrophic actinomycetes decreased significantly ($p \leq 0.05$) with increase in the concentration of the pollutant. The total heterotrophic bacteria and total heterotrophic fungi showed significant reduction ($p \leq 0.05$) with increase in the duration of the pollution while total heterotrophic actinomycetes showed no significance difference ($p \geq 0.05$) over the duration of pollution. Suspected bacterial isolates from the polluted soils were *Pseudomonas* sp., *Bacillus* sp., *Staphylococcus* sp., *Micrococcus* sp., *Flavobacterium* sp., *Serratia* sp., *Proteus* sp., while the suspected fungal isolates were *Mucor* spp., *Penicillium* spp., *Aspergillus* spp and *Rhizopus* spp. There were Significant differences ($p \leq 0.05$) in soil pH, organic matter, total nitrogen, potassium, magnesium, available phosphorus and base saturation of the polluted soil and that of the control while the values of sodium, calcium, exchangeable acidity and electrical conductivity of the polluted soil did not show significant difference with that of the control. There was rather a steady increase which was not statistically different from the control with increase in concentration of the pollutant. The results of this study revealed that soil polluted with little quantities of palm oil mill effluent could enhance microbial proliferation and thus, increase soil fertility, while a heavy application inhibits the same.*

Keywords: Effect, Effluent, Palm oil, Microflora, Fertility

Introduction

The world has entered a period of unprecedented environmental change as evidenced in the rapid growth of natural and man-made changes in the biosphere in recent times. Environmental Pollution resulting in changes in environmental quality is one of the issues raising widespread alarm. An increase in socio-economic activities worldwide has been accompanied by an even faster growth in pollution stress. This is because the majority of human needs can only be met with goods and services provided by industries. Industries have the capacity to improve as well as degrade the environment. Raw materials, which have been taken from the natural resource base, are converted into products by the industry. Lang [16] reported that during the production process pollutants are often released into the environment.

Soil is an efficient purifying medium with a great capacity to receive and decompose wastes matter by its microflora yielding nutrients in the process [23]. However, if the input of the pollutants exceeds the soil purifying limit, the effectiveness of soil microbial activity is reduced considerably. As a result, there will be an adverse change in the soil physicochemical properties which consequently affect the growth and development of the crop plants.

Industrial effluents as pollutants contain a large number of both known and unknown substances formed during the production process. There is a direct impact of pollutants on minerals, organic matter and microbial community of soil [23]. The discharge of industrial effluents especially without treatment may have profound influence on physicochemical and biological properties of soil related to soil fertility. The palm oil industry is a major agro based industry in Nigeria especially in the Southern part where palm oil trees are found both in the wild and plantations. During processing of the ripe fruits to extract the oil, a lot of waste water called palm oil mill effluent (POME) is generated. It is estimated that for 1 tonne

of crude palm oil produced, 5 - 7.5 tonnes of water ends up as POME [3]. The solid waste products that result from the milling operation are empty fruit bunches, palm fibre, and palm kernel. Much of the POME results from water used in processing [30][1]. Reports indicate that most of the POME is not treated before being discharged into the surrounding environment especially by the small scale mills, causing pollution problems.

POME, if untreated, contains high amounts of fatty acids, proteins, carbohydrates and other plant materials [27][6][11]. It alters the environmental parameters such as biological oxygen demand, dissolved oxygen, carbon/nitrogen ratio and chemical oxygen demand as well as moisture content and general soil quality [30]. This has been supported by many researchers [21][2]. These parameters affect soil microbial flora which in turn affects soil fertility [5]. Soil fertility is a very important factor, considering the type of use a soil can be put into agriculturally. The raw or partially treated POME has an extremely high content of degradable organic matter, which is due in part to the presence of unrecovered palm oil [3]. This highly polluting wastewater can, therefore, cause pollution of waterways due to oxygen depletion and other related effects as reported by some researchers [3]. Thus, while enjoying a most profitable commodity, palm oil, the adverse environmental impact from the palm oil industry cannot be ignored. It has been observed that most of the POME produced by the small-scale traditional operators undergoes little or no treatment and is usually discharged into the surrounding environment.

In recent years, economic boom in fast developing countries has been witnessed with spectacular progress in industrialization which has resulted in an increased generation of industrial wastes such as liquid, solid and volatile organic gases [3]. In an attempt to dispose of these, man has carelessly polluted the environment. These reasons have made it justifiable to assess the effects of POME on the microbial populations of the soil which enhances soil fertility.

The general objective of this study was to assess the effects of POME on microbial population and physicochemical properties of an agricultural soil.

Materials and methods

Experimental design

The soil sample used in this study was an agricultural soil collected from the botanical garden in the University of Calabar. The pollution of the soil was performed artificially with four concentrations (0 ml, 250 ml, 500 ml, 1000 ml and 2000 ml) of palm oil mill effluent in a completely randomized manner. And the duration of this study was sixteen weeks.

Collection of samples

Soil sample

Soil samples of 0-15cm depth were collected (5 kg) from five locations on an agricultural soil within university of Calabar by excavation using spade. The collected samples from all locations were thoroughly mixed on the spot in order to obtain composite sample. 5 kg of this soil was weighed into four different polythene bags [29].

Palm Oil Mill Effluents Samples

Palm Oil Mill Effluents (POME) were collected from Obasanjo's palm oil mill factory Mfamosing-Akamkpa, in Cross River State. The sample was collected with clean plastic containers rinsed several times with the same sample and transported within 24 h to the Department of Soil Science, University of Calabar for analyses and treatment.

Pollution of the Soil Sample

Pollution of the soil was achieved by employing standard methods [31]. 0 ml, 250 ml, 500 ml, 1000 ml and 2000 ml of the effluent was added to 2 kg of already dried soil in each polythene and mixed thoroughly for even distribution. These polluted soil and the unpolluted soil (control) were left outside under normal environmental condition for the duration of 16 weeks (Four months)

Microbial analysis

Microbial analysis was carried out before and after pollution. 10 g of soil sample was collected aseptically, labelled and stored in ice packed plastic coolers and transported to the Microbiology Department Laboratory University of Calabar where microbial analysis was carried out within 24 h of sampling so as to maintain the stability of the sample without significant alteration in the microbial population [31].

Dilution

Serial dilution was carried out by weighing 10 g of soil in to 90 ml of sterile saline water contained in a stoppered 200 ml volumetric flask and agitated to dislodge the microorganisms from the soil particles. From this initial dilution, a ten-fold serial dilution was prepared.

Enumeration of heterotrophic Bacteria

The counts of total heterotrophic bacteria in the soil samples were determined by pour plating 1 ml of 10^{-5} dilutions into nutrient agar (NA). The medium was incorporated with antifungal agent (50 µg/ml Nystatin), in order to prevent the growth of fungal contaminants. Bacterial colonies were counted after 24 h of incubation at room temperature and reported as colony forming units (cfu/g) of soil [4].

Enumeration of heterotrophic Fungi

The total heterotrophic fungi count was measured by pour plating 1 ml of 10^{-3} dilution into Sabouraud Dextrose Agar (SDA) supplemented with antibacterial agents (50 µg/ml of streptomycin and 30 µg/ml of penicillin) to inhibit the growth of bacterial contaminants. Fungal counts were reported after 72 h of incubation [4].

Enumeration of heterotrophic Actinomycetes

Enumeration of total actinomycetes was achieved by pour plate technique. 1 ml of 10^{-2} dilution was plated onto sodium caseinate agar, 50 µg/ml of nystatin and 30 µg/ml of tetracycline was added to inhibit fungal and bacterial growth. An actinomycetes count was reported 7 days after incubation at room temperature [22][30].

Maintenance of pure isolates

Bacterial colonies were repeatedly transferred to freshly prepared nutrient agar plates by the streak-plate method and allow growing for 48 h before stocking. Similarly, distinct fungal and actinomycetes colonies were subcultured repeatedly on freshly prepared Sabouraud Dextrose Agar plates and sodium caseinate agar, respectively. Pure isolates of the microorganisms were maintained on agar slants as stock, which were preserved in the refrigerator for further use.

Characterization and Identification of isolates

Various methods were used to characterize and identify the isolates [4][15][40]. The test results for bacteria were evaluated using characteristics presented in *Bergey's Manual of Determinative Bacteriology* [15].

Representative colonies of fungal isolates were characterized and identified based on their cultural and morphological features. The characterizations were achieved through staining techniques-using lactophenol in cotton blue [4][15][40].

Soil Physicochemical analysis

Physicochemical analysis of the pristine soil was carried out before and after pollution with varying concentrations of the oil mill effluent. Also physiochemical analysis of each soil sample polluted with different concentrations of each effluent was also carried out bimonthly (every 8 weeks). Physicochemical analysis was assessed in terms of the following parameters:

Particle size and textural class analysis

In carrying out this test, the Bouyoucos-type hydrometer method was used [12].

Soil pH

Soil pH was determined in water 1:2 soils: water ratio using pH meter with glass electrode. 20 g of air-dried soil was weighed into a 50 ml beaker, and 20 ml of sterile saline water was added and allowed it to stand for 30 mins. The electrode of the pH meter was inserted into the 1:2 soil /water partly settled suspension and measured the pH. The result was recorded as soil pH measure in water [40].

Electrical conductivity (EC)

In the same soil solution (1:2 soil /water solution) for pH determination, electrical conductivity electrode was inserted into the partly settled suspension and the EC was measured [40]

Organic matter

This was determined by the dichromate wet-oxidation method [26].

Total nitrogen

Total nitrogen was determined by the micro- Kjeldahl method [9].

Available phosphorus

Available phosphorus was extracted with acid fluoride using Bray P-1 method [8]

Exchangeable cations

The bases were extracted with neutral NH_4OAc . Calcium and magnesium were determined in the extract by EDTA titration, and potassium and sodium by the use of flame photometer [39].

Exchangeable acidity

A^+ and H^+ were obtained by leaching the soil with NH_4Cl solution and the and the extract titrated with standard NaOH as described by I.I.T.A (1979).

Exchangeable acidity ($\text{Al} + \text{H}^+$) – Exchangeable $\text{Al} =$ Exchangeable H

Effective cation exchange capacity

This was determined by calculation. That is, total exchangeable bases ($\text{Ca} + \text{Mg} + \text{Na}$) + Exchangeable acidity (EA).

Percentage base saturation

This will be achieved by dividing the total exchangeable bases by exchangeable cation capacity and multiplied by 100 [6].

$$\% \text{ base saturation} = \frac{\text{summation of exchangeable bases}}{\text{exchangeable cation capacity}} \times 100$$

Statistical analysis

All statistical analysis of data from various treatments was carried out using analysis of variance (ANOVA) test using factorial experiment. Means were separated using least significance difference (LSD).

Results

Microbial analyses

Table 1 below shows the enumeration of total heterotrophic bacteria (THB), total heterotrophic fungi (THF) and total heterotrophic actinomycetes (THA) in the pristine soil. The counts obtained were as follows $1.90 \pm 1.41 \times 10^7$ Cfug, $1.29 \pm 1.25 \times 10^5$ cfug and $9.2 \pm 1.25 \times 10^3$ cfug respectively.

TABLE 1

Total microbial counts of the soil before the commencement of the study

THB (cfu/g)	$1.90 \pm 1.41 \times 10^7$
THF (cfu/g)	$1.29 \pm 1.25 \times 10^5$
THA (cfu/g)	$9.2 \pm 1.25 \times 10^3$

Key: THB = Total heterotrophic bacteria, THF= Total heterotrophic fungi, THA= Total heterotrophic actinomycetes, cfu/g = Colony forming unit/gram

Table 2 below shows the effect of different concentrations of palm oil mill effluent on microbial population of POME- polluted soil. The results obtained for THB in 250 ml, 500 ml, 1000 ml and 2000 ml were $1.59 \pm 1.84 \times 10^7$ cfu/g, $1.24 \pm 1.40 \times 10^7$ cfu/g, $7.0 \pm 1.38 \times 10^6$ cfu/g and $5.9 \pm 1.22 \times 10^6$ cfu/g respectively. The results showed that there was no significant difference ($p \leq 0.05$) in the mean counts of THB obtained in 250 ml and 500 ml, and no significant difference ($p \leq 0.05$) was also observed in THB counts of 1000 ml 2000 ml of the polluted soil (Table 2).

The results for the counts of THF in different concentrations showed that there was a significant increase ($p \leq 0.05$) in total heterotrophic fungi (THF) counts of 250 ml and 500 ml with mean counts of $2.02 \pm 3.00 \times 10^5$ cfu/g, $1.71 \pm 1.89 \times 10^5$ cfu/g respectively over the control, and a significance reduction ($p \geq 0.05$) in THF counts of 1000 ml and 2000 ml with mean counts of $1.07 \pm 1.26 \times 10^5$ cfu/g and $5.1 \pm 1.42 \times 10^4$ cfu/g respectively (Table 2).

The total heterotrophic actinomycetes (THA) counts in different concentrations showed that, there was a significant difference ($P \leq .05$) in THA counts among the concentrations. There was a significant increase ($p \leq 0.05$) in THA counts in 250ml and 500ml with mean counts of $1.33 \pm 2.09 \times 10^4$ cfu/g, $1.41 \pm 1.84 \times 10^4$ cfu/g respectively over the control with mean count of $8.9^b \pm 1.73 \times 10^3$ cfu/g while there was no significance ($p \geq 0.05$) in THA in 1000 ml and 2000 ml with mean counts of $4.8 \pm 1.16 \times 10^3$ cfu/g and $3.6^b \pm 1.09 \times 10^3$ cfu/g respectively over the control (Table 2).

TABLE 2
Effects of concentrations of pollution on microbial population in palm oil mill polluted soils

	0ml	250lm	500ml	1000ml	2000ml
THB (cfug ⁻¹)	1.89 ^a ±2.90x10 ⁷	1.59 ^a ±1.84x10 ⁷	1.24 ^a ±1.40x10 ⁷	7.0 ^b ±1.38x10 ⁶	5.9 ^b ±1.22x10 ⁶
THF (cfug ⁻¹)	1.26 ^c ±2.51x10 ⁵	2.02 ^b ±3.00x10 ⁵	1.71 ^b ±1.26x10 ⁵	1.07 ^c ±1.26x10 ⁵	5.1 ^d ±1.42x10 ⁴
THA (cfug ⁻¹)	8.9 ^b ±1.73x10 ³	1.33 ^a ±2.09x10 ⁴	1.14 ^a ±1.84x10 ⁴	4.8 ^b ±1.16x10 ³	3.6 ^b ±1.09x10 ³

Key: means with the same letter along the horizontal arrays indicates significant difference ($P \leq .05$), and means with different letter along the column indicates no significance difference ($P \geq .05$)

THB = Total heterotrophic bacteria, THF= Total heterotrophic fungi, THA= Total heterotrophic actinomycetes, cfu/g = Colony forming unit/gram

Table 3 below shows the effect of POME and duration of pollution on microbial populations. The total heterotrophic bacteria (THB) counts after 8 weeks and 16 weeks of pollution were $1.62 \pm 4.21 \times 10^7$ cfu/g and $7.84 \pm 0.98 \times 10^6$ cfu/g respectively. Significant reduction ($p \geq .05$) was observed in 16 weeks of pollution (Table 3). Also the counts obtained for total heterotrophic fungi (THF) after 8 weeks of pollution produced a mean count of $1.56 \pm 3.81 \times 10^5$ cfu/g was significantly higher than the THF counts after 16 weeks of pollution with a mean count of $1.07 \pm 2.06 \times 10^5$ cfu/g (Table 3). And for the effect of duration of pollution on total heterotrophic actinomycetes (THA) counts obtained, there was no significant difference ($P \geq .05$) in the count obtained after 8 weeks and 16 weeks of pollution.

TABLE 3
Effects of duration of pollution on microbial population

	8 weeks	16 weeks	LSD
THB (cfu/g ⁻¹)	$1.62 \pm 4.21 \times 10^7$ ^a	$7.84 \pm 0.98 \times 10^6$ ^b	2.69
THF (cfu /g ⁻¹)	$1.56 \pm 3.81 \times 10^5$ ^a	$1.07 \pm 2.06 \times 10^5$ ^b	3.38
THA (cfu/ g ⁻¹)	$9.4 \pm 1.28 \times 10^3$ ^b	$7.4 \pm 1.11 \times 10^3$ ^b	4.04

Key: means with the same letter along the horizontal arrays indicates significant difference ($P \leq .05$), and means with different letter along the column indicates no significance difference ($P \geq .05$)

THB = Total heterotrophic bacteria, THF= Total heterotrophic fungi, THA= Total heterotrophic actinomycetes, cfu/g = Colony forming unit/gram, LSD= Least significant difference

Characterization and Identification of Microbial Isolates

Microbial isolates were characterized and identified based on their morphological and biochemical characteristics. The following bacteria were isolated from POME polluted soil; *Pseudomonas spp.*, *Serratia spp.*, *Bacillus spp.*, *Micrococcus spp.*, *Proteus spp.*, while the fungal isolates were *Rhizopus spp.*, *Aspergillus spp.*, *Mucor spp.*, *Penicillium spp.*

Soil pH

The pH values of the control (0 ml) soils were in the range of 5.66 ± 0.50 to 5.7 ± 0.21 . The pH values in POME treated soils ranged from 6.28 ± 0.19 to 6.83 ± 0.24 . There was no significant difference ($p \leq .05$) in the pH value of the soils treated with 250 ml, 500 ml and 1000 ml POME and in the soil treated with 250 ml, 500 ml, 1000 ml and 2000 ml, but there was a significant difference ($p \geq .05$) in the pH value of the soil treated with 2000 ml of the effluent. There was significant difference ($p \geq 0.05$) in the pH values over the duration of pollution (Figure 1).




Figure 1: Effects of concentrations of palm oil mill effluent on the soil pH

Organic Matter Content

The values of organic matter in control (0 ml) soils were in the range of $2.41 \pm 0.23\%$ and $2.41 \pm 0.09\%$. The organic matter content of the soils treated with varying concentrations of POME ranged from $3.53 \pm 0.19\%$ to $4.97 \pm 0.26\%$ (Table 6 and Figure 2).




Figure 2: Effects of concentrations of palm oil mill effluents on soil organic matter

Total Nitrogen Content

The control soils (0 ml) had the total nitrogen content in the range of $0.06 \pm 0.2\%$ and $0.06 \pm 0.01\%$. In the soil treated with varying concentrations of POME, the total nitrogen content ranged from $0.08 \pm 0.01\%$ to $0.10 \pm 0.04\%$. Significant difference ($p \leq 0.05$) was not observed among concentration. There was no significant difference in total nitrogen content over the duration of study,




Figure 3: Effects of concentrations of palm oil mill effluent on total nitrogen content of the polluted soil

Effective Cation Exchange Capacity (ECEC)

The values of ECEC in the control (0 ml) soils were in the range of 9.45 ± 0.43 cmol/kg and 9.7 ± 0.06 cmol/kg (Table 5). The value of ECEC in palm oil mill effluent polluted soil increased to between 9.8 ± 0.14 cmol/kg and 11.55 ± 0.30 cmol/kg (Table 6 and Figure 3). There was significant increase ($p \leq 0.05$) in ECEC in 2000 ml polluted soil and significance difference ($p \geq 0.05$) did not exist in 250ml, 500 ml, 1000 ml and that of 250 ml, 500 ml 1000 ml.

TABLE 4
Physicochemical properties of the soil before the commencement

Parameters	Values
pH	5.60
Electrical conductivity (ds/m)	0.055
Organic matter (%)	2.41
Total nitrogen (%)	0.06
Available Phosphorus (mg/kg)	31.54
THC (mg/kg)	BDL
Ca (cmol/kg)	4.00
Mg (cmol/kg)	2.10
Na (cmol/kg)	0.06
K (cmol/kg)	0.21
EC (cmol/kg)	1.11
ECEC (cmol/kg)	8.82
Base saturation (%)	85.16
Sand (%)	75.80
Silt (%)	12.60
Clay (%)	11.60

Lead (mg/kg)	6.20
Nickel (mg/kg)	9.44
Cadmium (mg/kg)	4.32
Chromium (mg/kg)	5.60
Iron (mg/kg)	919.66

TABLE 5
Effects of concentrations of pollution on the physiochemical properties of the soil polluted with palm oil mill effluent

	0 ml	250 ml	500 ml	1000 ml	2000 ml	LSD
pH	5.65 ^c ±0.50	6.83 ^a ±0.24	6.72 ^a ±0.20	6.65 ^a ±0.18	6.28 ^b ±0.19	0.20
Organic matter	2.41 ^c ±0.23	3.53 ^c ±0.19	4.26 ^b ±0.26	4.16 ^b ±0.26	4.97 ^a ±0.26	0.18
Avail. P	31.54 ^a ±2.64	71.1 ^c ±1.24	14.49 ^b ±2.02	11.49 ^f ±1.05	15.42 ^d ±1.18	0.85
Total N	0.06 ^c ±0.02	0.08 ^b ±0.01	0.10 ^b ±0.03	0.09 ^b ±0.02	0.10 ^b ±0.04	0.02
Mg	2.37 ^d ±0.09	2.73 ^b ±0.11	2.13 ^c ±0.08	2.64 ^c ±0.14	3.25 ^a ±0.09	0.09
Sodium (Na)	0.09 ^b ±0.03	0.11 ^b ±0.01	0.11 ^b ±0.02	0.12 ^b ±0.01	0.12 ^b ±0.04	0.04
Calcium (Ca)	4.75 ^a ±0.10	5.18 ^a ±0.9	5.31 ^a ±0.08	5.40 ^a ±0.05	5.50 ^a ±0.08	0.61
Ec	0.05 ^b ±0.01	0.13 ^a ±0.02	0.14 ^a ±0.01	0.15 ^a ±0.03	0.17 ^a ±0.02	0.02
Base saturation	78.52 ^b ±2.01	78.32 ^b ±1.53	78.06 ^b ±0.69	78.27 ^b ±0.89	80.09 ^a ±1.14	1.35
K	0.21 ^b ±0.02	0.29 ^a ±0.03	0.23 ^b ±0.01	0.16 ^b ±0.09	0.29 ^a ±0.08	0.02
EFEC	9.45 ^b ±0.43	10.61 ^b ±0.21	9.8 ^b ±0.14	10.63 ^b ±0.08	11.55 ^a ±0.30	0.69
EA	2.03 ^a ±0.8	2.3 ^a ±0.05	2.15 ^a ±0.04	2.31 ^a ±0.08	2.3 ^a ±0.02	0.20

Key: means with the same letter along the horizontal arrays indicates significant difference ($P \leq 0.05$), and means with different letter along the column indicates no significance difference ($P \geq 0.05$)

TABLE 6
Effects of duration of pollution on physico- chemical properties of the soil polluted with palm oil effluent

	8 weeks	16 weeks	LSD
pH	6.6 ^b ±0.08	6.25 ^d ±0.04	0.13
Organic matter	4.02 ^a ±0.10	3.71 ^b ±0.09	0.11
Avail. P	18.31 ^b ±0.24	19.7 ^a ±0.36	0.54
Total N	0.09 ^b ±0.02	0.08 ^b ±0.02	0.01
Mg	2.87 ^a ±0.10	2.37 ^c ±0.05	0.06
Na	0.13 ^b ±0.02	0.09 ^c ±0.02	0.02
Ca	5.38 ^a ±0.11	5.00 ^a ±0.09	0.38
Ec	0.13 ^a ±0.02	0.13 ^a ±0.01	0.01
Bs	79.19 ^a ±2.10	78.25 ^a ±1.80	0.86
K	0.22 ^b ±0.02	0.25 ^a ±0.03	0.01
ECEC	10.86 ^a ±0.28	10.02 ^b ±0.20	0.44
EA	2.26 ^a ±0.09	2.18 ^a ±0.08	0.13

Key: means with the same letter along the horizontal arrays indicates significant difference ($P \leq 0.05$), and means with different letter along the column indicates no significance difference ($P \geq 0.05$).

Discussion

Soil is a potent system of terrestrial ecosystem. The direct discharge of untreated industrial effluents may have profound influence on the physicochemical and biological properties of the soil. The pollutant level of POME varies with the quality of the raw material and production process used to manufacture the palm oil.

In this study, following pollution of soil with varying concentrations of POME, the soil showed profound changes in the microbial populations and physicochemical properties. In assessing the effects of different concentrations of POME on microbial population, the result showed no significant difference ($p \geq .05$) in THB mean count in 250 ml and 500 ml of effluent polluted soil over the controls. This implies that the THB mean counts in 250 ml and 500 ml the polluted soil fall in the same range with that of the control. Also, there was a significant reduction ($p \leq .05$) in THB in 1000 ml and 2000 ml over the control. These indifferences in THB mean count possesses some similar characteristics and hence have the same effect on soil bacteria. Also, the result showed that there was a significant increase ($p \geq .05$) in THF and THA in 250 ml and 500 ml (Table 2). This increase in fungi population may be attributed to the acidity nature of the effluents, because fungi thrive best in acidic environments than bacteria which can only survive in a neutral to alkaline pH environment. Significant increase ($p \geq .05$) in THA count was also observed in 250 ml and 500 ml which had THA count higher than the controls. This may also be attributed to the acidic nature of the effluents and the soil since soil actinomycetes have been known to tolerate acidic environment than the true bacteria. Soil actinomycetes can tolerate a pH level of up to 6.80, and this was the pH level in most of the treated soils in this study. An increase in microbial population has earlier been reported [30].

In terms of concentrations of pollution, 250 ml of the effluents was found to be most favourable concentration for the proliferation of microorganism, followed by 500 ml in this study. Similar result had earlier been reported [29]. He observed that effluent impacted soil at low concentration encouraged proliferation of microorganisms but at high concentration, inhibits the same [29], this was also observed in this study and could be attributed to the harmful level of the effluent at higher concentrations (Table 2). The components of the POME used in this study revealed that the effluents contain some metabolizable nutrients for the soil microorganisms and excess of these nutrients with the high water content suppressed the microorganisms [29]. Excess water in soil restricts microorganisms and their activities by preventing oxygen movement into and through the soil in sufficient quantity to meet the oxygen demand of the microorganisms [18][33].

Also in assessing the effect of duration of pollution and different concentrations of pollution on microbial population, THB showed significant increase ($p \geq .01$) with decrease in concentrations of pollutions and significant reduction with increase in the duration of pollution (Table 3). This result implies that there was a steady reduction in THB with increase in concentrations and duration of pollution. Also, the THF mean counts at 250 ml showed significant increase ($p \geq .05$) after 8 weeks of pollution and a significant reduction was observed in the same concentration of pollution after 16 weeks of pollution and while there was no significant difference in THF count at 500 ml, 1000 ml of after 8 weeks and 16 weeks of pollution (Table 3). Similarly, there was significant increase ($p \geq .05$) in THA at 250 ml, 500 ml, after 8 weeks of pollution and 250 ml after 16 weeks of pollution, and no significant difference was also observed in THA in 1000 ml, 2000 ml after 8 weeks of pollution and 500 ml, 1000 ml and 2000 ml after 16 weeks of pollution. This result implies that, there was a significant increase ($p \geq .05$) in THB, THF and THA in all the soil samples,

416 followed by a reduction in THB and THF after 16 weeks of pollution except THA which did
417 not show reduction after 16 weeks of pollution (Table 3).

418
419 The result of this study also showed changes in physicochemical properties. This change has
420 earlier been reported that continual applications of effluents on the soil can change soil
421 properties, e.g. pH and nutrient concentrations [37].

422 The result also showed that there was a significant increase ($p \geq .05$) in the pH values in all
423 the soil polluted with POME over the control. Increase in pH in POME polluted soil is in
424 agreement with the report of Okwute and Isu [30], who reported that when raw POME is
425 discharged into the soil, the acidic pH seems to gradually increase to alkaline as
426 biodegradation takes place, and this was the case in this study as pH of the polluted soils
427 increases from 5.65 ± 0.05 in the control to between 6.28 ± 0.19 and 6.83 ± 0.24 . The
428 increase in pH of POME polluted soil has also been earlier reported [29][18].

429
430 The polluted soils were also characterised by increased in organic matter and total nitrogen.
431 Organic matter of a soil is usually determined and reported as a measure of organic carbon.
432 Organic matter content strongly affects soil fertility by defining the energy supply to
433 microorganisms, availability and quality of substrates and the biodiversity necessary to
434 sustain much soil functions[26][40][41]

435 436 **Conclusion**

437 This study on the effects of palm oil mill effluent on soil microflora and fertility showed that
438 the mean counts of total heterotrophic bacteria, total heterotrophic fungi and total
439 heterotrophic actinomycetes increased with decrease in concentration of pollution and
440 decreased with increase in the duration of pollution in the soils. Similarly, the POME also
441 caused profound changes on the soil physicochemical properties of the polluted soils.

442 The result revealed significant differences in soil pH, organic matter content, total nitrogen,
443 available phosphorus, magnesium, electrical conductivity, percentage base saturation,
444 potassium and effective cation exchange capacity between polluted and non-polluted soil
445 while no significant difference was observed in calcium, sodium and exchangeable acidity
446 between polluted and non-polluted soil.

447
448 In conclusion, this study revealed that POME could be harmful to the soil if not properly
449 managed and also useful to the soil if properly discharged since light application of the
450 effluents could enhance microbial proliferation which enhances soil fertility. Therefore,
451 government should create awareness to those involved in small and large scale palm oil
452 processing on the need for proper disposal of effluent because if not properly managed, it can
453 negatively affect soil fertility by inhibiting microbial proliferation.

454 455 **References**

- 456 1. Abdul, L. A., Suzylawati, I., Norliza, I., and Subhas, B. Removal of suspended solids
457 and residual oil from palm oil mill effluents. *J. Chem. Tech. Biotech.* 2003;
458 78(9):971-978.
- 459 2. Aggelis, G. C., Gavala, H. N. and lyberatos, G. Combined and separated Aerobic and
460 anaerobic bio-treatment of green olive debittering wastewater. *J. Agric Eng.*
461 *Res.*2001; 80 (3):283-292.
- 462 3. Ahmad, A., Ismail, S., and Bhatia, S. Water recycling from palm oil mill effluent
463 (POME) using membrane technology. *Desalination.* 2003; 157:87-95.
- 464
465

4. Antai, S.P., Unimke, A.A. & Agbor, R.B. Assessment of the Heterotrophic and Crude Oil Utilizing Microorganisms of Imo River Estuary of the Niger Delta Mangrove Ecosystem. *American International Journal of Biology*. 2014; 2(1): 29 – 42.
5. Arias, M. E. Jose, A.G., Gonzalez-Villa, F. J. and Ball, A. S. Soil Heath: A New Challenge to Microbiologist and Chemists. *Soil Heath Int. Microbial*. 2005; 8:13-21.
6. Bek-Nielsen, C., Singh, G., Toh, T. S. Bioremediation of palm oil mill effluent. In: Proceedings Porim International Palm Oil Congress. Kuala Lumpur, Malaysia; 1999.
7. Bohn. H.L., McNeal, B.L. and O'Connor, G.A. Soil Chemistry. John Wiley & Sons, New York; 1985
8. Bray, R. H. and Kurtz, L. T. Determination of Total, Organic and Available forms of Phosphorus in Soils. *Journal of Soil Science*. 1945; (5) : 35-45.
9. Bremmer, J. M. Total Nitrogen. In: D. L. Spark (eds). *Methods of Soil analysis. Part 3- chemical methods*, SSSA Book Series 5, Madison, Wisconsin; 1996.
10. Carter, M. R. Soil quality for sustainable land management: organic matter and aggregation interactions that maintain soil functions. *Agron*. 2002; *J*. 94: 38-47.
11. Chan, K. W., Watson, I., Lim, K. C. Use of oil palm waste material for increased production. Paper presented at the Conference on Soil Science and Agricultural development in Malaysia, Kuala Lumpur, Malaysia; 1980.
12. Day, P. R. Particle Fractionation and particle size analysis. In: C. A Black (eds). determination of phosphorus in Natural waters. *Analytical Chemistry*. 1965; *ACTA* 27:31-38.
13. Dilip K. D. Introductory Soil Science, Kalyani Publishers, Delhi.
Ecosystem. *American International Journal of Biology*. 2006; 2(1): 29 – 42.
14. Fakayode, S. O. Impact assessment of industrial effluent on water quality of the receiving Alaro river in Ibadan Nigeria. *AJEAM-RAGEE*. 2005; *10*: 1-13.
15. Holt, J. G., Krieg, N. R., Senath, P. H., Stanley, L. T. and Williams, S. T. Bergey's Manual of Determinative Bacteriology. (9th Ed). Baltimore, Willams and Wilkins; 1994.
16. Láng, I. Environmental Management and Sustainable Development. In Nath, B., Hens, L. and Devuyt, D (Eds.). *Environmental Management, Vol. 3. Instruments for Implementation Brussels*: VUB University Press, VUB University; 1993
17. Lim, C.H. and P'ng, T. C. Land application of digested palm oil mill effluent by sprinkler system. Proceeding of the Seminar on Land Application of Oil Palm and Rubber Factory Effluent, Serdang; 1983..
18. Lim, K. H. Trail on long term effects of application of palm oil mill effluent on soil properties, oil palm nutrition and yields. *Proc. cont. Agric*. 1987.

19. Lowe, H. T. (1994). Utilisation of short rotation forestry for on-site fuel wood grown as part of an effluent disposal scheme. A Thesis for Master of Agricultural Science at Massey University, Palmerston North, New Zealand; 1994..
20. Ma, A. N. Innovation in management of palm oil mill effluent. *The Planter Kuala Lumpur*. 1999;75(881): 381-389.
21. Ma, A.N. Environmental management for the palm oil industry. *Palm Oil Dev*. 2000; 30:1-9
22. Mill; A sustainable Waste Management Practice. Review in Environmental Science and Biotechnology, DOI: 10. 1007/s11157 – 010-9199-2.
23. Nagaraju, M., Narasimha, G. and Rangaswami, V. Influence of sugar industry effluents on soil enzyme activities. *Ecol. Environ. Cons.*2009; 15:89-94.
24. Narasimha, G. Sridevi, A., Venkata, S. R. and Rajasekhar R. B. Effects of cotton ginning mill effluents on soil enzymatic activities and nitrogen mineralization in soil. *J. Chem. Pharm. Res.* 2011; 3: 128-137.
25. Narasimha, G., Babu, A. K. and Rajasekhar, R. B. Physicochemical and biological properties of soil samples collected from soil contaminated with effluents of cotton ginning industry. *J. Environ. Biol.* 1999; 20: 235-239.
26. Nelson, D. W., Sommers, L. E. Total carbon, organic carbon and matter. In: Page, A. L., Miller, R. H. and Keeney, D. R. (eds.) *Methods of Soil Analysis, Part 2 Chemical and Microbiological Properties*. Madison, Wisconsin pp. 539-579; 1982.
27. Ngan, M.A., Zajima, Y., Asahi, M. and Junit, H. A novel treatment processes for palm oil mill effluent. *PROIM Technology*; 1996.
28. Nilima, W. and Madhuri, S. (2005). Fungal diversity of paper industry waste and soils irrigated by the effluents. *Nat. Environ. Pollut. Tech.* 2005;4: 44-52.
29. Nwaugo, V. O., Chinyere, G. C. and Inyang, C. U. Effects of palm oil mill Effluents (pome) on soil Bacterial flora and Enzyme activities in Egbama. *Plant Product Research Journal*. 2008; 12: 10 – 13.
30. Okwute, O. L. and Isu, R. N. Impact Analysis of palm oil mill effluent on the Aerobic Bacterial density and ammonium oxidizers in a dumpsite in Anyigba, Kogi State. *Afr. J. Biotech* 6(2):116-119; 2007.
31. Orhue, E. R., Uzu, O. F. and Osaigbovo, A, U. Effect of Combining Rubber Effluent with Rock Phosphate on Some Soil Chemical Properties and Early Growth of Maize (*Zea mays*. L). *International Journal of Soil Science*. 2007;2: 82-95.
32. Oviasogie, P. O, Aghimien, A. E. Macro nutrient status and speciation of Cu, Fe, Zn and Pb in soil containing palm oil mill effluent. *Global J. of Pure and Applied Science*. 2002; 9 (1): 71-80.

33. Paul, E. A., Clark, F. E. Soil Microbiology and Biochemistry. Academic Press Incorporated San Diego, California, pp.11-234; 1989.
34. Phang, S. M. The potential of integrating fish culture with an algal pond system treating palm oil mill effluent. In: Proceedings of the International Oil Palm/Palm Oil Conference PORIM and ISP, pp. 422 - 428. Phosphorus in Soils. *Journal of Soil Science*. 1987; (5) : 35-45.
35. Prescott, L.M., Klein, D, A. and Harley, J. P. Microbiology 5th edition. McGraw Hill, Education, United States; 2001.
36. Sonawane, D.V., Lawande, S. P., Gaiwad, V. B. and Kuchekar, S. R. (2010). Impact of Industrial Waste on Soil Quality and Organic Matter around Kurkumbh Industrial Area Daund, Pune District. *Int. J. Chem. Sci.* 2010; 8(1):97-102
37. Tubonimi, J. K., Patricia, U. A., Herbert, O. S. and Amonia, O. B. Impact of palm oil mill effluent on water quality. *Res. J. Appl. Sci.* 2007; 2(7):842 – 845.
38. Tugel, A.J. and Lewandowski, A. M. Soil Biology Primer. NRCS Soil Quality Institute. Ames, Iowa. pp.50; 1999.
39. Udo, E. J., Ibia, T. O., Ogunwale, J. A., Ano, A. O. and Esu, I. E. *Manual of Soil , Plant and Water Analysis*. Sibon Books Limited, Lagos; 2009.
40. Unimke, A.A., Antai, S.P. & Agbor, R.B. Influence of Seasonal Variation on the Microbiological and Physicochemical Parameters of Imo River Estuary of the Niger Delta Mangrove Ecosystem. *American International Journal of Biology*. 2014; 2 (1): 61-74.
41. Unimke, A.A., Antai S.P., Agbor, R.B., Nseabasi, NO & Agbo, BE. Evaluation of Seasonal Variation in the Microbial and Heavy Metal Concentrations of Imo River Estuary of The Niger Delta Mangrove Ecosystem. *Advanced Research in Agriculture and Veterinary Science*. 2014;1 (2): 88 – 94.
42. Vesiland, P.A Peirce, J J and Weiner, R.F. Environmental Pollution and Control Wastewater Discharges. *Water Science Technology*. 1990; 25 (11), 449–456.
43. Wood, B. J. A review of current methods for dealing with palm oil mill effluents. *Planter*. 1977; 53: 477-495.