

**Original Research Article**

**Preliminary Assessment and Status of Fauna Species  
Diversity in Ipinu Igede Community Range Forest in Oju  
Local Government of Benue State, Nigeria.**

**ABSTRACT**

Appraisal of fauna species which form an integral part of range ecology, in rangeland based protected areas is necessary before any meaningful conservation work can commence. It was aim at assessing number of fauna species and status in Ipinu Igede community forest, Oju.

Animal species were enumerated through direct on-site field observation and indirect indices to provide data that could enhance the management and sustainable utilization of the forest. Fifty-six species of wild animals from 40 families were found in the study area. There were 21 species of mammals from 15 families, 6 species of reptiles from 6 families and 29 species of birds from 19 families respectively. A total of 1,419 sightings were recorded. The most abundant animal species found in the area were *Epixerus ebii*, *Eidolon helvum*, *Chlorocebus tantalus*, *Papio anubis*, *Philothemus irregularis*, *Musophaga violacea*, *Numidia meleagris* and *Fringilla bicalcaratus*. The status of most mammals was least concern and some of the reptiles had data deficient. All the birds are known to breed in the reserve. Transect C had the highest diversity index (0.0130) and transect A had the lowest animal diversity index (0.0061) but highest species count of 36 as well as individual animal species sighted. transect C had the highest diversity index (0.0130) and the second highest species count of thirty four (34). Transect A on the other hand had the lowest animal diversity index (0.0061) and the highest species count of thirty six (36) as well as individual animal species sighted (454) It can be concluded from the result that Ipinu Igede community forest contains representative sample of fauna in the Guinea Savanna ecosystems of Nigeria and has to be protected through conservation awareness and community participation.

**Key words:** Range forest, fauna, diversity, Status, community forests.

**1. INTRODUCTION**

The survival and continuity of many endemic, rare and threatened species found in a given rangeland depend on sustainable conservation through its assessment to determine its

35 current status. The potential of range forest habitation of wild animals is grossly unexplored  
36 in many areas across Nigeria, especially local community forests [1]

37 According to Daniel *et al.* [2], appraisal of both flora and fauna species in protected  
38 areas is necessary before any meaningful conservation work can commence. Fauna resources  
39 are the entire wild animal of any particular region or ecosystem [3]. These wild animals can  
40 be found in all ecosystems including forests, grasslands, plains, wetlands and deserts [4]. The  
41 potential of range forest habitation to the management and sustainability of wild animals is  
42 grossly unexplored in many areas across Nigeria, especially local community range forests  
43 [1]. Fauna species assessment has more concentration to national parks and game/wildlife  
44 parks. However, many local rangeland communities support unique flora and fauna species  
45 making them important in terms of conservation and scientific interest.

46 Approach to species listing is an important initial stage in the collection of appropriate  
47 data necessary for effective management and conservation of animals and plants in a  
48 protected rangeland [5]. Therefore, knowledge of the species composition of a protected  
49 rangeland, their status and how they relate with other components of the habitat is highly  
50 essential and as well indicate the status of most fragile, threatened species. Insight to species  
51 list and status is becoming increasingly important as conservators and rangeland managers  
52 are tasked to assists conservation biologists to construct informed management plans for  
53 endangered species. This has become critical because most fauna species live in tropical  
54 forest which are increasingly been impacted by human modification and natural occurrences  
55 [6, 7, 8].

56 The status of the population of any individual species is crucial information to the  
57 wildlife ecologists, because this information determines individual fitness to its environment  
58 and also predicts their ultimate success or failure [9]. Idowu and Morenikeji, [10] report that,  
59 wildlife is increasingly being regarded as removable resources and man mostly is known for  
60 his high taste for exploiting its populations in the environment hence, their habitation to  
61 various rangeland and status ought to be monitored to ensure proper utilization of their  
62 habitat. In the same vein, Daniel *et al.* [2] stated that, appraisal of both flora and fauna  
63 species which form an integral part of animal ecology, in wildlife based protected areas is  
64 necessary before any meaningful conservation work can commence.

65 Ipinu Igede Community Forest in Oju Local Government of Benue State is one of the  
66 reserve that is rich in biodiversity, though had no appreciable ecological survey of the  
67 resources, hence, the dearth of information necessary for the development of the reserve. The  
68 area has suffered from activities of illegal logging operators and hunting thereby threatening

important flora and fauna species also the quest for a local fauna database and the alarming rate of species loss informed the need for wildlife based inventory in the study area. Thus there is need to appraise the composition of fauna species using diversity indices to ascertain the present status of fauna species of the community forest. The objective of the study is therefore to quantify fauna species composition and abundance using diversity indices and ascertain the status of the species. It is quiet obvious that baseline data generated from the study will promote effective conservation of biodiversity and management plan within the communal forest.

## 2. MATERIAL AND METHOD

### 2.1 The Study Area

The research was carried out at Ipinu Igede Community Forest Reserve in Oju Local Government Area of Benue State, Nigeria. The community forest is an ancestral heritage site for Igede people of Benue State stretching through three communities; Oyinyi, Andibilla and Uchenyim. The forest contains relicts of traditional worship practices in the area, although, the traditional religious worship practices are no longer strong and appreciated due to acceptance of Christianity. However, the laws and taboos governing the forest are still observed by the people of Igede.

The forest which is located in the Southern Guinea savanna belt comprise of both hilly and lowland part and lies between Longitude 8° 25' 0" E and 8° 41' 67"E and Latitude 6° 51' 0' N and 6° 85' 0' N [11]. It has an area of approximately 4 km<sup>2</sup> on a fairly flat land drained by four main seasonally flowing streams (Abadehe, Otuhukwu, Ekpaa and Ugbunwu) which are tributaries to River Ogbugwu. The mean annual rainfall is between 1200mm and 1500mm. The mean annual temperature is 30°C. Relative Humidity is between 60% and 80% wet but decreases in the early months of dry season.

It is a derived tropical rainforest characterized by luxuriant vegetation with high composition of riparian forest, of the large trees are *Cola gigantean*, *Elaeis guinensis*, *Ficus exasperate*, *Khaya spp*, *Afzelia africana* [11]. Dominant herbaceous species include *Sphenoclea zeylanica*, *Pentodon pentandrus*, *Ageratum conyzoides*, *Nymphaea lotus* and *asystasia gangetica*. The area has relatively abundant faunal resources; commonly sighted mammals are the primates (baboons and monkeys), bush buck, oribi, grass cutter, squirrel and common duiker. Reptiles were alligator and snakes. Birds include guinea fowl, francolin, village weavers, king fisher, Grey horn bill, Yellow billed kite and Abyssinian roller.

## **2.2 Data Collection Techniques**

### **2.2.1 Species Diversity and Status**

Species list and diversity was determined by direct observation along four transects of 2.0km by 10m broad (0.1ha) distributed randomly as described by Osunsina *et al.* [12] and indirect indices as well as through information from hunters and bush meat processing and selling centers.

Survey was carried out in the morning hours between 6:00 to 9:00am and early evening time between 4:00 to 7:00 pm. This was to ensure counting of even the shyest animal species as the period coincides with the time the animals are most likely to search for water and preys, or graze on land [13].

Status assessment of Mammals, Birds and Reptiles was based on the information from hunters and forest protected agent and follows Ezealor [14] and IUCN (International Union for Conservation of Nature) Red list.

## **2.3 Data Analysis**

Descriptive statistics (tables, chart and figures) were used to analyze species lists of mammals, reptiles and birds.

### **2.3.1 Status Categories of Mammals, Reptiles and Birds**

Categories outlined by Ezealor [14] were used to assign the status of mammals, reptiles and of birds. This is as follows;

Vu = Vulnerable (Likely to become endangered if the factor that is posing threat persists).

LR/ cd = Low risk-conservation dependent (Species in no immediate danger but survival will depend on implementation of effective conservation measures in the community forest).

NT = Near threatened (species is approaching the threshold of vulnerability)

EN = Endangered (species is unlikely to survive if the factor that is posing threat persists).

RB = Resident breeder

R {B} = Resident but breeding not approved.

PM = Palearctic migrant

129 AFM = Migrates within Nigeria

130 DD = Data deficient

### 131 2.3.2 Diversity Indices

132 Diversity indices were calculated for each transect using Simpson's diversity index, which is  
133 a measure of heterogeneity of a site taking into consideration the number of species and  
134 density of individual species [15, 16]. The index is expressed as;

$$135 \quad I = \frac{q \sum n(n-1)}{N(N-1)}$$

136 Where I = Simpson diversity index.

137 N = total number of individuals enumerated.

138 q = number different species enumerated.

139 n = number of individuals of species enumerated.

140

## 141 3. RESULTS

142 Fifty-six species of vertebrates (wild animals) belonging to 42 families were  
143 identified in the study area. They belong to three classes of Mammalia, Reptilia and Aves.  
144 Twelve species of mammals were identified through direct sighting while 9 species was  
145 through their signs and activities as well as interviews of hunters and bush meat processing  
146 and selling centers. Three species of reptiles were identified through direct sighting while 3  
147 were indirect assessment. All the Bird species were identified through direct sighting.

148 Majority of the identified mammal species were in the category of LR/cd, followed  
149 Vu and some NT approaching the threshold of vulnerability. Most of the Reptile species were  
150 of DD category (Data Deficient) while others fall within the LR/cd category. Almost all the  
151 identified birds' species are resident breeders in the forest (RB). Some of the species  
152 identified in the study area are presented in plate 1 to 6.

### 153 Wild Animal Species Distribution and Abundance Across the Transects

154 The total numbers of individual species recorded were 1,419. The class Aves had the  
155 highest frequency (974) 68.6% followed by Mammalia (429) 30.2% and Reptilia (16) 1.1%  
156 (Figure. 1). The total numbers of animals occurrence recorded for the various transects (A, B,  
157 C and D) were 454, 332, 294 and 339 respectively (Table 2 and Figure 2). The species with

the highest abundance to class mammalian was *Epixerus ebii* (67.60%) followed by *Eldolon helvum* (15.30 %) and the least was *Tragelaphus scriptus* (0.23%). For class reptilian; Green snake (*Philothamus irregularis*) (56.25%) followed by alligator (*Alligator mississippiensis*) (31.25%) and the least was Black cobra (*Naja melanoleuca*). Class aves was Violet plantain eaters (*Musophaga violacea*) (22.59%) followed by Guinea fowl (*Numidia meleagris*) (10.37%) and the lowest was Goshwak hawk (*Accipiter africana*) (0.10%) respectively. However, there were 18 constant species present in all the transects. This include *Arvicanthis niloticus*, *Epixerus ebii*, *Acrocephalus rufescens*, *Centropus sensgalensis*, *Colius striatus*, *Coracias abyssinicus*, *Crinifer piscator*, *Euplectes franciscannus*, *Francolinus bicalcaratus*, *Lamprotornis nitens*, *Lanchura cucullata*, *Musophaga violacea*, *Numidia meleagris*, *Phynonotus barbatus*, *Piocephalus senegalus*, *Streptopelia semitorquata*, *Streptopelia senegalensis* and *Vidua macroura*

The Simpson diversity indices of animal species showed that transect C had the highest diversity index (0.0130) and the second highest species count of thirty four (34). Transect A on the other hand had the lowest animal diversity index (0.0061) with the highest species count of thirty six (36) as well as individual animal species sighted (Table 3).

**Table 1: Species List, Mode of Identification and Status of Mammals, Reptiles and Birds in Ipinu Igede Community Forest**

176

Species		Mode of Identification				Status		
S/N	Common Names	Scientific Names	Family	DS	IND	INH	PC	Category
<b>Mammals</b>								
1	Common duiker	<i>Sylvicapra grimmia</i>	Bovidae	X	x	X	X	VU
2	Bush buck	<i>Tragelaphus scriptus</i>	Bovidae	X	-	X	X	LR/cd
3	Oribi	<i>Ourebia ourebi</i>	Bovidae	X	-	X	X	LR/cd
4	Water buck	<i>Kobus ellipsyprymnus</i>	Bovidae	-	-	X	-	LR/cd
5	Wild dog (pale fox)	<i>Vulpes pallida</i>	<i>Canidae</i>	-	x	X	-	LR/cd
6	Tantalus monkey	<i>Chlorocebus tantalus</i>	Cercopithecidae	X	x	X	-	LR/cd
7	Olive baboon	<i>Papio Anubis</i>	Cercopithecidae	X	-	X	-	LR/cd
8	African pigmy hedgehog	<i>Atelerix albiventris</i>	Erinaceidae	-	-	X	-	LR/cd
9	Bush baby	<i>Galago cameronensis</i>	Galagidae	-	-	X	-	LR/cd
10	Spotted hyena	<i>Crocuta crocuta</i>	Hyenidae	-	-	X	-	Vu
11	Porcupine	<i>Hystrix cristata</i>	Hystriidae	-	-	X	-	Vu
12	African bush rabbit	<i>Lepus microtis</i>	Leporidae	X	-	X	X	LR/cd
13	Pangolin	<i>Manis gigantean</i>	Manidae	-	-	X	-	Vu
14	African grass rat	<i>Arvicanthis niloticus</i>	Murinae	X	-	X	X	LR/cd
15	Giant rat	<i>Cricetomys gambianus</i>	Nesomyidae	-	x	X	X	LR/cd

16	Fruit bats	<i>Eidolon helvum</i>	Pteropodidae	X	-	X	-	NT
17	Tree squirrel	<i>Epixerus ebii</i>	Sciuridae	X	-	X	-	LR/cd
18	Ground squirrel	<i>Xerus erythropus</i>	Sciuridae	X	-	X	-	LR/cd
19	Grass cutter	<i>Thryonomys swinderianus</i>	Thryonomyidae	X	-	X	X	LR/cd
20	African civet	<i>Civettictis civetta</i>	Viverridae	X	-	X	X	LR/cd
21	Common Genet	<i>Genetta genetta</i>	Viverridae	-	-	X	-	LR/cd
<b>Reptiles</b>								
22	Alligator	<i>Alligator mississippiensis</i>	Alligatoridae	X	-	-	X	LR/cd
23	Green snake	<i>Philothemus iregularis</i>	Colubridae	X	-	X	-	LR/cd
24	Black cobra	<i>Naja melanoleuca</i>	Elapidae	X	-	X	-	DD
25	Ballpython	<i>Python regius</i>	Pythonidae	-	-	X	-	LR/cd
26	Monitor lizard	<i>Veranus niloticus</i>	Veranidae	-	-	X	X	DD
27	Puff adder	<i>Bitis arientans</i>	Viperidae	-	-	X	-	DD
<b>Aves (Birds)</b>								
28	Yellow billed kite	<i>Milvus aegyptius</i>	Accipitridae	X	-	-	-	RB
29	Black Fork tailed kite	<i>Milvus migrans</i>	Accipitridae	X	-	-	-	RB
30	Goshawk hawk	<i>Accipiter Africana</i>	Accipitridae	X	-	-	-	RB
31	King fisher	<i>Ispidina lecontei</i>	Alcedinidae	X	-	-	-	RB
32	Grey horn bill	<i>Tockus nasutus</i>	Bucerotidae	X	-	X	-	RB
33	Littleringed plover	<i>Charadrius dubius</i>	Charadriidae	X	-	-	-	RB
34	Ringed plover	<i>Charadrius hiaticula</i>	Charadriidae	X	-	-	-	RB
35	Mouse bird	<i>Colius striatus</i>	Coliidae	X	-	-	-	RB
36	Laughing dove	<i>Streptopelia senegalensis</i>	Columbidae	X	-	X	-	RB
37	Mourning dove	<i>Streptopelia decipiens</i>	Columbidae	X	-	-	-	RB
38	Redeyed pigeon	<i>Streptopelia semitorquata</i>	Columbidae	X	-	-	-	RB
39	Abyssinian roller	<i>Coracias abyssinicus</i>	Coraciidae	X	-	-	-	RB
40	Black magpie	<i>Ptilostomus afer</i>	Corvidae	X	-	-	-	RB
41	Senegal coucal	<i>Centropus sensgalensis</i>	Cuculidae	X	-	X	-	RB
42	Black throated coucal	<i>Centropus leucogaster</i>	Cuculidae	X	-	-	-	RB
43	Bronze manikin	<i>Lanchura cucullata</i>	Estrildidae	X	-	-	-	RB
44	Violet plantain eater	<i>Musophaga violacea</i>	Musophagidae	X	-	X	X	RB
45	Gray plantain eater	<i>Crinifer piscator</i>	Musophagidae	X	-	-	-	RB
46	Spurred francolin	<i>Francolinus bicalcaratus</i>	Phansianidae	X	x	X	X	RB
47	Guinea fowl	<i>Numidia meleagris</i>	Phansianidae	X	-	X	X	RB
48	Green wood hoopoe	<i>Phoeniculus purpureus</i>	Phoeniculidae	X	-	-	-	RB
49	Common garden bulbul	<i>Phynonotus barbatus</i>	Phynonotidae	X	x	-	-	RB
50	Red bishop	<i>Euplectes franciscannus</i>	Ploceidae	X	-	-	-	RB
51	Village weaver	<i>Ploceus cucullatus</i>	Ploceidae	X	-	-	-	RB
52	Senegal parrots	<i>Piocephalus senegalus</i>	Psittacidae	X	-	-	-	RB
53	Glossy starling	<i>Lamprotornis nitens</i>	Sturnidae	X	-	-	-	RB
54	Rufus cane warbler	<i>Acrocephatus rufescens</i>	Sylviidae	X	x	-	-	RB
55	Pintailed whydah	<i>Vidua macroura</i>	Viduidae	X	-	-	-	RB
56	Village indigo	<i>Vidua chalybeate</i>	Viduidae	X	-	-	-	RB



177 Field Survey, 2017  
 178 In the above table;  
 179 DS = Direct Sighting  
 180 IND = Indices (Animals sign and activities)  
 181 INH = Interview of hunters  
 182 PC = Bush meat processing and selling center  
 183 — = Absent  
 184 X = Present  
 185

186 **Table 2: Wild Animals Species Distribution and Abundance According to Transect**

S/N	Scientific Names	Common Names	Tran. A	Tran. B	Tran. C	Tran. D	Total	Abundance
<b>Mammals</b>								
1	<i>Arvicanthis niloticus</i>	African grass rat	4	3	5	1	13	3.03
2	<i>Chlorocebus tantalus</i>	Tantulus monkey	38	-	-	-	38	8.56
3	<i>Civettictis civetta</i>	African civet	2	-	1	-	3	0.79
4	<i>Eidolon helvum</i>	Fruit bats	38	2	-	26	66	15.38
5	<i>Epixerus ebii</i>	Tree squirrel	110	40	59	81	290	67.60
6	<i>Lepus microtis</i>	African bush rabbit	1	-	1	-	2	0.47
7	<i>Ourebia ourebi</i>	Oribi	1	1	-	-	2	0.47
8	<i>Papio Anubis</i>	Olive baboon	-	-	-	4	4	0.93
9	<i>Sylvicapra grimmia</i>	Common duiker	4	-	-	-	4	0.93
10	<i>Thryonomys swinderianus</i>	Grass cutter	1	1	1	-	3	0.79
11	<i>Tragelaphus scriptus</i>	Bush buck	-	-	-	1	1	0.23
12	<i>Xerus erythropus</i>	Ground squirrel	1	1	1	-	3	0.79
<b>Total</b>			<b>200</b>	<b>48</b>	<b>68</b>	<b>113</b>	<b>429</b>	<b>100%</b>
<b>Reptiles</b>								
13	<i>Alligator mississippiensis</i>	Alligator	3	-	-	2	5	31.25
14	<i>Naja melenaleuca</i>	Black cobra	1	-	1	-	2	12.50
15	<i>Philothemus irregularis</i>	Green snake	3	-	6	-	9	56.25
<b>Total</b>			<b>7</b>	<b>-</b>	<b>7</b>	<b>2</b>	<b>16</b>	<b>100%</b>
<b>Aves</b>								
16	<i>Accipiter Africana</i>	Goshawk hawk	-	-	1	-	1	0.10
17	<i>Acrocephatus rufescens</i>	Rufus cane warbler	13	8	7	11	39	4.00
18	<i>Centropus leucogaster</i>	Black throated coucal	-	2	1	2	5	0.51
19	<i>Centropus sensgalensis</i>	Senegal coucal	12	12	14	17	55	5.65
20	<i>Charadrius hiaticula</i>	Ringed plover	-	-	3	7	10	1.03
21	<i>Chardrius dubius</i>	Little ringed plover	-	-	2	-	2	0.20
22	<i>Chelictinia riocourii</i>	Forked tail kite	4	-	-	4	8	0.82
23	<i>Colius striatus</i>	Mouse bird	7	11	9	1	28	2.87
24	<i>Coracias abyssinicus</i>	Abyssinian roller	2	1	2	8	13	1.33
25	<i>Crinifer piscator</i>	Gray plantain eater	26	39	19	16	100	10.27
26	<i>Euplectes</i>	Red bishop	9	4	4	1	18	1.85




27	<i>franciscannus</i> <i>Francolinus</i>	Spurred francolin	14	13	12	19	58	5.95
28	<i>bicalcaratus</i> <i>Ispidina lecontei</i>	King fisher	-	-	4	13	17	1.74
29	<i>Lamprotornis nitens</i>	Glossy starling	4	6	3	2	15	1.54
30	<i>Lanchura cucullata</i>	Bronze manikin	11	9	7	6	33	3.39
31	<i>Milvus aegyptius</i>	Yellow billed kite	4	6	1	-	11	1.13
32	<i>Musophaga violacea</i>	Violet plantain eater	57	70	48	45	220	22.59
33	<i>Numidia meleagris</i>	Guinea fowl	20	32	26	23	101	10.37
34	<i>Phoeniculus purpureus</i>	Green wood hoopoe	2	-	11	-	13	1.33
35	<i>Phynonotus barbatus</i>	Common garden bulbul	16	8	18	11	53	5.44
36	<i>Piocephalus senegalus</i>	Senegal parrots	5	3	6	1	15	1.54
37	<i>Ploceus cucullatus</i>	Village weaver	4	5	4	-	13	1.33
38	<i>Ptilostomus afer</i>	Black magpie	-	2	-	2	4	0.41
39	<i>Streptopelia decipiens</i>	Mourning dove	12	-	2	3	17	1.75
40	<i>Streptopelia semitorquata</i>	Redeyed pigeon	8	11	5	4	28	2.87
41	<i>Streptopelia senegalensis</i>	Laughing dove	3	11	5	9	28	2.87
42	<i>Tockus nasutus</i>	Grey horn bill	11	27	-	13	51	5.24
43	<i>Vidua chalybeate</i>	Village indigo bird	2	-	2	-	4	0.41
44	<i>Vidua macroura</i>	Pintailed whydah	1	4	3	6	14	1.44
<b>Total :</b>			<b>247</b>	<b>284</b>	<b>219</b>	<b>224</b>	<b>974</b>	<b>100%</b>
<b>ΣTotal: Animals/Reptiles/Aves</b>			<b>454</b>	<b>332</b>	<b>294</b>	<b>339</b>	<b>1419</b>	<b>-</b>

187 Field Survey, 2017


188

189

190



**Fig. 1. Class Distribution of Wild Animals in Ipinu Igede Community Forest**



**Fig. 2. Distribution of Wild Animals According to Transect**

**Table 3: Simpson Diversity Index of Wild Animal Species in the Transect**

Transect	Individual Species (n)	Total of Species (N)	Diversity Index
A	36	454	0.0061
B	27	332	0.0064
C	34	294	0.0130
D	29	339	0.0071




Plate 1: *Francolinus bicalcaratus*



Plate 2: *Xerus erythropus*



Plate 3: *Thryonomys swinderianus*



Plate 4: *Veranus niloticus*



Plate 5: Feather of *Musophaga violacea*



plate6: dropping of *Sylvicapra grimmia*

#### 4. DISCUSSIONS

The relatively high population of vertebrates' species (wild animals) found in the area is typical of West African taxa [12]. The list of species surveyed is for the understanding of faunal dynamics in the conservancy of any protected area. This is in line with the observation by Yager *et al.* [17] at Makurdi zoological garden. More so, the high incidence of *Epixerus ebii*, *Eidolon helvum*, *Arvicanthis niloticus* and some primates in the community forest may not be unconnected to the fact that the species are not accepted as meat by the people in the surrounding communities. Similar observations have been made by Mbaya and Magwi [5] at

Sambisa game reserve. The relatively low status of some mammals and reptiles in the forest such as duiker, spotted hyena, porcupine and pangolin suggests high incidence of poaching for meat and traditional medicine because of the very little effort being made to protect the resources of the forest. Snake species are of least concern however the community tends to dislike them; hence the quest to eliminate them from their surrounding environment could have being the possible cause. Generally, some wild animals have higher tolerance of hunting pressure than other because of their home range and their reproductive potentials. Some may be subjected to less hunting pressure because the taste and acceptance of their meat or their ease of preparation. Local techniques used in capturing some species also put them under varying pressures. Some animal species also response to vegetation structure that allow a clear view of their surrounding and enable them to move with speed and agility through the under growth, like ground squirrel make use of the forest edge and strip vegetation because they are not able to survive an arboreal life unlike the tree squirrel that dominate the area. Duikers were also found in sparsely dense habitat.

Bird life in the study area is largely recorded in relation with trees ranging from the violet plantain eater, gray plantain eater, guinea fowl, grey horn bill, rufus cane warbler, king fisher, spurred francolin, village weaver and others which normally winter around the streams. A large number of birds live on seeds, fruits, buds, and nectar or insects that are found in the arboreal environment. These include, Grey plantain eater and Senegal kingfisher respectively. The high bird species diversity in the area could be due to the fact that the area acts as a sanctuary from the degraded habitats surrounding it and nesting materials and availability of edible fruits bearing tress. This observation is in line with the report by Egwumah *et al.* [18] at GRA and Ankpa quarters Benue State.

In a comparative form the total number of twenty-one mammalian species is just about 8.5% of 247 species reported for Nigeria by Happold [19]. The number is also lower than either of the 123 species reported for Guinea Savanna or 97 species for Sudan Savanna of Nigeria [19]. So, the species richness of the forest might not be unconnected with its size which is relatively small compared to the size of Guinea Savanna (473,904 km<sup>2</sup>) or Sudan Savanna (927,338km<sup>2</sup>). This observation agrees with Usher [20] report, that species diversity is often affected by the size of habitat and that diversity is positively correlated with habitat size. Biodiversity assessment and conservation management purposes, distribution or pattern of occupancy is very important and this has been found to vary with different environmental location and condition for a given species.

## 5. CONCLUSION

The study area showed a moderately high number of vertebrate species despite the human disturbance in the area through activities such as hunting and logging. Basic knowledge of species list and occurrence within a region is a necessary starting point to predict species extinction under habitat loss, to understand the potential impacts on biodiversity, as well as to prioritize conservation efforts and designing conservation areas. It is recommended that the nature reserve be protected from resource over exploitation which may lead to degradation, its management should be based on sound ecological principles, and conservation education be promoted.

## REFERENCES

1. Ogunjemite BG, Ajayi B, Agbelusi EA. Habitat structure of chimpanzee communities in Nigeria: a comparison of sites. *Acta Zoological Sinica*. 2007; 53 (4); 579-588.
2. Daniel IE, Henry MI, Augustine UO. Preliminary assessment of tree species diversity in Afi Mountain Wildlife Sanctuary, Southern Nigeria, *Agriculture and Biology Journal of North America*. 2012; 3.12.486.492 ScienceHuB, <http://www.scihub.org/ABJNA>
3. Wikipedia. <https://en.m.wikipedia.org/wiki/Fauna> Retrieved on 5th November, 2017 wildlife heritage of Africans. TRAFFIC Reports. [www.traffic.org](http://www.traffic.org).
4. Ijeomah HM, Augustine UO, Damilola O. Analysis of Poaching Activities in Kainji Lake National Park of Nigeria. *Environment and Natural Resources Research*. 2012; 3(1). <http://dx.doi.org/10.5539/enrr.v3n1p51>
5. Mbaya YP, Malgwi H. Species list and status of mammals and birds in Sambisa game reserve, Borno State, Nigeria, *Journal of Research in Forestry, Wildlife and Environment*. 2010; 2 (1): 135-140.
6. Alamu LO, Agbey BO. Deforestation and endangered indigenous tree species in Southwest Nigeria: *International Journal of Biodiversity and Conservation* 2011; 3.7 :291-310.
7. Babagana G, Mohammed MA, Garba M. Environmental Impact of Natural Resources Exploitation in Nigeria and the way forward. *Journal of Applied technology in Environmental sanitation* 2012; 2, (2): 95-102.
8. Ahmed RK. Biodiversity loss: A threat to urban landscape. *International Research Journal of Arts and Social Sciences* 2013; 2. 6:144-149
9. Martin C. *The Rainforest of West Africa*. Birkhauser, London. 1990.
10. Idowu MA, Morenikeji OA. Wild Fauna Conservation in Nigeria. *Environment and Natural Resources Research*, 2015; 5(3); 98-108.
11. Okwoche SO. Assessment of tree species composition and diversity of Ipinu-Igede community forest in Oju L.G.A, Benue State, Nigeria. Unpublished Undergraduate project submitted to the Department of Forestry, Wildlife and Range Management, University of Agriculture Makurdi, Nigeria 2017; pp 10-35.
12. Osunsina IOO, Inah EI, Ogunjinmi AA, Onadeko SA, Osunsina O. Distribution and diversity of flora and fauna in International Institute of Tropical Agriculture (IITA)

- 291 forest and nature reserve, Ibadan. Oyo State. Nigeria. *Journal of Agriculture, Forestry*  
292 *and the Social Sciences* (JOAFSS) 2012; 10 (2): 289-302.
- 293 13. Adamu IA. An assessment of floristic composition of Kwiambana Game Reserve, A  
294 Ph.D. thesis, presented to the Department of Geography, Usmanu Danfodiyo  
295 University, Sokoto (Unpublished). 2006; 122p.
- 296 14. Ezealor HU. Critical Sites for Biodiversity Conservation in Nigeria. *Nigerian*  
297 *Conservation Foundation*, Lagos, Nigeria. 2002; 110pp.
- 298 15. Simpson EH. Measurement of Diversity, *Nature*; 1949; 163: 688.
- 299 16. Ojo LO. Data collection and analysis for biodiversity conservation: *In Practical of the*  
300 *Inception Meeting and Training Workshop on BRAAF Assessment and Monitoring*  
301 *Techniques in Nigeria*. Eds. B.A. Ola-Adams and L.O. Ojo National Committee on  
302 Man and Biosphere 1996; Pg. 142-145
- 303 17. Yager GO, Enefolo JO, Tyowua BT. Comparative study of fauna species diversity of  
304 Makurdi Zoological Garden, Benue State, Nigeria, *International Journal of*  
305 *Development and Sustainability* ISSN: 2186-8662 – [www.isdsnet.com/ijds](http://www.isdsnet.com/ijds). 2017; 6  
306 (12): 2163-2172 ISDS Article ID: IJDS17112401
- 307 18. Egwumah PO, Iwar IM, Ogbonna L. A survey of the avi-fauna within Makurdi  
308 Metropolis of Benue State, Nigeria. *Journal of Research in Forestry, Wildlife and*  
309 *Environment*, 2009; 1 (1); 75-83.
- 310 19. Happold DCD. *The Mammals of Nigeria*; Oxford press, London. 1987.
- 311
- 312 20. Usher MB. *Conservation Biology; A Training Manual for Biological*  
313 *Diversity and Genetic Resources*. Rapoor- Vijay and White (Ed.) 1992; pp 80- 83.