

Biolog Identification of Fermenting Yeasts from Fermented Teff (*Eragrostis teff* (Zucc.)) Dough

Belay Tilahun¹, Anteneh Tesfaye², Diriba Muleta², Andualem Bahiru² and Zewdu Terefework³,

¹Wolkite University, Department of Biotechnology, Wolkite Ethiopia, belayt21@gmail.com; ²Institute of Biotechnology, Addis Ababa, University, Ethiopia and ³MRC-ET molecular Diagnostics, Addis Ababa, Ethiopia

ABSTRACT

Background: Injera is one of baked product, which is commonly prepared from teff (*Eragrostis teff* (Zucc.)) flour. It is a staple food and believed to be consumed on daily basis by two-thirds of Ethiopians, a population of nearly 100 million. As it is a product of naturally fermented dough, the course of fermentation is done by consortia of microorganisms. Therefore, this study was aimed to isolate and identify yeasts from fermenting Teff (*Eragrostis teff*) dough.

Place and Duration of Study: Samples were collected from different sources in Addis Ababa, Ethiopia. Laboratory isolation and identification of isolates were carried out at Holeta Biotechnology Institute, Microbial Biotechnology laboratory and Ethiopian Institute of Biodiversity, Microbial laboratory from December to April, 2015 to June, 2016 G. C.

Methodology: A total of 200 dough samples were collected from households with different fermentation stage. Twenty (20) yeast isolates with different cultural characteristics were selected and further identified by ominilog identification systems.

Results: The seven yeast isolates obtained from teff dough were identified as *Pichia fermentans*, *Pichia* spp., *Rhodotorula aurantiaca* B, *Pichia fluxuum*, *Candida humilis*, *Trichosporon beigelii* B and, *Cryptococcus albidus* Var *aerus*

Conclusion: This study has confirmed the presence of different yeast in the fermenting teff dough and also supports the involvement of consortia of various groups of microorganisms in the course of the fermentation.

Keywords: Teff dough, *Injera*, Microbial Identification, ominilog

1. INTRODUCTION

Yeasts are unicellular, eukaryotic and polyphyletic organisms classified in the kingdom fungi. They are ubiquitous, and commonly found on fruits, vegetables, insect and other plant materials. Some yeast is found in association with soil and water. Approximately 100 genera comprising more than 1500 species of yeast have been described (Kurtzman and Fell, 2006). The significance of yeasts in food technology in a world of low agricultural production and rapidly increasing population makes the production of food grade yeasts extremely important (Bekatorou *et al.*, 2006). In Ethiopia there are several fermented foods such as, kocho, bulla, tella, tej, milk product and injera, etc. A wide variety of fermented foods and beverages are consumed in Ethiopia being prepared from a wide range of raw materials using traditional techniques. These include: Injera, kocho, tella, awaze, borde and tejj (Askal and Kebede, 2013). Injera is one of the fermented foods that is made from different cereals, including sorghum, teff, corn, wheat, barley, or a combination of some of these cereals (Mogesse, 2006). Injera from teff (*Eragrostis teff*) is much more relished by most Ethiopians than that from any other source. It is a thin soft fermented baked food usually obtained after

the flour of cereals has been subjected to 24 to 96h of traditional fermentation depending on the ambient temperature (Askal and Kebede, 2013; Bultosa *et al.*, 2008)

The fermentation process uses natural inoculants from different sources in a mixed form (Stewart and Getachew, 1962). Teff injera is getting popular in the developed world because of its gluten free nature and being a whole grain product (Hiwot *et al.*, 2013). Teff is a cereal crop which is mainly cultivated in Ethiopia for the purpose of making injera (Bultosa *et al.*, 2008; Mogesse, 2006). For injera making, teff grain is considered by many as superior when compared to other cereal grains used in the country (Yetneberk *et al.*, 2005).

A lot of research was undertaken on microbial profile of these commodities through conventional methods. In most cases, strain of *Saccharomyces cerevisiae*, *Rhodotorula* spp, *pichia*, were found to dominate fermenter in tella, injera, milk product and other fermented foods. However, the yeast species involved in injera fermentation not studied well using standard Biolog MicroStation identification technology for shortening fermentation time and selecting the potential fermenter yeast in future. The aim of this study was isolation, identification and characterization of yeast species involved in injera fermentation by using biolog identification system.

2. MATERIALS AND METHODS

2.1. Sample collection and Description of the Study sites

A total of 200 teff dough samples (two hundred gram each) were collected from each of 14 sampling sites of fermenting dough samples with different time of fermentation (Brhanu, 1985) from around Addis Ababa, the capital of Ethiopia. The samples were transported aseptically to Holeta Biotechnology Institute, Microbial Biotechnology laboratory, for processing and microbial isolation. Sample processing and laboratory isolation of yeast were carried out in Holeta Biotechnology Institute, Microbial Biotechnology laboratory. Ominilog identification was carried out at microbial laboratory at Ethiopian Biodiversity Institute.

2.2. Isolation and selection of yeasts

After the samples were transported to the laboratory 10 g dough samples was transferred aseptically into separate flask with 90 mL sterile 0.1% peptone water and homogenized. Thereafter, a tenfold serial dilution was made. From appropriate dilution factor, 1ml of the suspension was streaked onto pre-solidified yeast extract glucose Chloramphenicol Bromophenol blue agar medium. The plates were incubated at 25°C for 24h. Then after 24h incubation, the selected colonies (10-20) were sub-cultured on yeast extract glucose Chloramphenicol Bromophenol blue agar medium three times to purify the isolates. Yeast isolates were selected according to their cultural characteristics (colony size, colony color, colony texture) on yeast extract glucose Chloramphenicol Bromophenol blue agar based on Bergey's Manual (Whitman, 2009).

2.3. Biolog Omnilog identification and characterization of isolates

Yeasts were sub cultured to Biolog Universal Yeast (BUY) agar and Biolog Universal and were incubated at 26°C for 24 to 48 h. Yeast suspension were prepared in 9ml sterile distilled water and adjusted to 47% T using Biolog YT turbidity standard. One hundred micro liters of inoculums were added to each well of the YT Micro Plate (Biolog Inc) and incubated at 26°C. (Kurtzman, and Fell, 2006). A YT Micro Plate was read by the Biolog Micro Station Reader (BiologInc) at 24 h, 48 h, and 72 h at a single wavelength of 590 nm. The Biolog

software micro log3 ver. 4.20.05 compared the results obtained with the test strain to the database and provided identification based on distance value of match and separation score produces similarity index value and probability. An acceptable species identification must have similarity index value >0.5 or probability >75% were Chosen only for species identification and characterization (Biolog, 1993). Since, some yeast species are inhibited by the tetrazolium violet redox dye used in Biolog Micro Plates, the YT Micro Plate was configured with both oxidation tests and assimilation tests. The first 1-36 column panel contained carbon source for oxidation tests using tetrazolium violet as a colorimetric indicator of oxidation. Because, the last column panel of 49-60 wells contained 2 carbon sources; these wells test was used for co-utilization of various carbon sources with D-xylose. The read is going to be positive on the Micro Plate, if <X>, and when the database result for that well was negative the printout showed <X- to indicate a mismatch where the database reaction was negative. If there were a negative read, X with no brackets and database value for that well were positive the well reads X+ indicating a positive reaction in the database. At the time of a read the data were compared to the database in order to determine the ID (Kurtzman and Fell, 2006).

2.4. Data Analysis

Omnilog clustering of yeasts based on the carbon utilization was done using past software.

3. RESULTS AND DISCUSSION

3.1. Isolation of yeast and gas production from glucose

The yeast isolates recovered were observed to have different features of colonial morphology. The yeast isolates showed diverse cultural characteristics with regard to colour (white, gray pigmentation or blue), shape (circular or irregular), edge (irregular or smooth) and elevation (flat or raised). Furthermore, the sizes of the colonies ranged from medium to large.

And a total of 20 yeast isolates were selected on the basis their potential in gas production (Bakheit, 2008). Yeast isolates with good potential of gas production were believed to be good fermenter and for the formation of many eyes on ready- to-eat *injera*, and these were selected for molecular identification. The percentage occurrence on culture media recorded as, 5% *Pichia fermentans*, 15% *Pichia spp.*, *Rhodotorula aurantiaca* B, 15% *Pichia fluxuum*, 10% *Candida humilis*, and, 10% *Cryptococcus albidus* Var *aerus* other 35% were not identified. Six yeast species were identified which is involved in dough fermentation by Omni Log identification system.

3.2. Biolog Omnilog identification and characterization of isolates

Selected yeasts isolates were aseptically transferred on BUY agar for Biolog identification. Seven yeast species were identified which may involve in dough fermentation by Omni Log identification system. Based on the result, Table 1 shows those species of yeasts: *Pichia fermentans*, *Pichia spp.*, *Rhodotorula aurantiaca* B, *Pichia fluxuum*, *Candida humilis*, *Trichosporon beigeli* B and, *Cryptococcus albidus* Var *aerus* with their identification statuses in the YT Biolog Micro Plate.

Table 1. Biolog Micro station identification result.

Species	Probability	Similarity	Distance	Remark
<i>Cryptococcus albidus</i> var. <i>aerus</i>	100%	0.73	5.98	Identified
<i>Pichia fermentans</i>	100%	0.653	5.33	Identified
<i>Rhodotorula acheniorum</i>	100%	0.623	5.78	Identified
<i>Trichosporon beigeli</i> B	100%	0.615	5.91	Identified
<i>Pichia</i> spp	99%	0.693	4.62	Identified
<i>Pichia fluxuum</i>	98%	0.668	4.87	Identified
<i>Candida humilis</i>	86%	0.553	5.47	Identified

135

136 Based on the result, Table 2 shows those species of yeasts: *Pichia fermentans*, *Pichia* spp.,
137 *Rhodotorula aurantiaca* B, *Pichia fluxuum*, *Candida humilis*, *Trichosporon beigeli* B and,
138 *Cryptococcus albidus* Var *aerus* which exhibited positive result for oxidation test and
139 assimilation potential of different carbon coated in the YT Biolog Micro Plate. Yeast Micro
140 Plate has two different reactions: assimilation of carbohydrates and oxidation. Six yeast
141 species which were identified by BioLog (*Pichia fermentans*, *Pichia* spp., *Rhodotorula*
142 *aurantiaca* B, *Pichia fluxuum*, *Candida humilis*, *Trichosporon beigeli* B and, *Cryptococcus*
143 *albidus* Var *aerus*) have positive oxidative test for Dextrin, stachyose, D-galactose, glucose,
144 L-aspartic acid, D-gluconic acid, D-melibiose, sucrose, turanose, D-mannitol, tween 80, D-
145 arabitol, D-meleitose, L-sorbose, Xylitol, Glycerol, Gentiobiose, Sucrose, Salicin, Maltotriose,
146 L-sorbose carbon sources.

147

148 **Table 2. Oxidation and assimilation result of yeast using YT Biolog MicroPlate**

Isolate code	Oxidation	Assimilations
AAUYT30B (<i>P. fermentans</i>)	Dextrin, stachyose, D- galactose, glucose,	Fumaric acid, L-malic, methyl succinate, bromo succinic acid, L-glutamic acid, a-keto- glutaric acid, D- gluconic acid, Dextrin, Cellobiose, Gentiobiose, Maltose, Maltotriose, Palatinose, D- raffinose, Stachyose, Sucrose, D-trehalose, Turanose, glucose, D-galactose, L-rhamnose, D-glucoside, Amygdalin, Arbutin, Salicin, D-mannitol, D-sorbitol, adonitol, xylitol, glycerol, tween 80, D- ribose, 1,2- propanediol+ D-xylose a
AAUYT26C AAUYT32A AAUYT34A (<i>Pichia</i> spp.)	Dextrin, stachyose, D- galactose, glucose,	Fumaric acid, L-malic, methyl succinate, bromo succinic acid, L-glutamic acid, a-keto- glutaric acid, D- gluconic acid, Dextrin, Inulin, Cellobiose, Gentiobiose, Maltose, Maltotriose, D-melibiose, Palatinose, D- raffinose, Stachyose, Sucrose, D- trehalose, Turanose, D-glucosamine, a-D-glucose, D-galactose, L-rhamnose, L-sorbose, Amygdalin, Arbutin, Salicin, D-maltitol, D-mannitol, D-sorbitol, adonitol, xylitol, glycerol, tween 80, D- ribose, D- xylose, D-galactose+ D-xylose, ,2-propanediol+ D- xylose
AAUYT21B AAUYT27A AAUYT25A (<i>P. fluxuum</i>)	L-aspartic acid, D-gluconic acid, dextrin, D-melibiose, sucrose, turanose, D- mannitol and tween 80	Gentiobiose, Maltotriose, Arbutin and D-mannitol, L-rhamnose, L-sorbose, Amygdalin, Salicin, D- sorbitol, adonitol, xylitol, glycerol, tween 80, D- ribose, D-xylose N-acetyl-L-glutamic acid, D-xylose, D-glucuronic acid+ D-xylose, dextrin, D-xylose, D- melibiose+ D-xylose, D-galactose+ D-xylose, 1,2- propanediol+ D-xylose
AAUYT31A	Dextrin, D- arabitol	Fumaric acid, L-malic, methyl succinate, bromo

(*C. albidus* Var *aerius*)

AAUYT23B
(*Candida humilis*)

L-aspartic acid, D-gluconic acid, Dextrin, D-melibiose, L-sorbose, Xylitol, Glycerol and tween 80

AAUYT22A
(*R. aurantiaca* B)

Gentiobiose, Sucrose, Salicin and tween 80

AAUYT28A
AAUYT24C
AAUYT35A
(Unidentified)

L-aspartic acid, D-gluconic acid, Dextrin, Maltotriose, L-sorbose, D- sorbitol, Glycerol and tween 80

succinic acid, Gentiobiose, Amygdalin, D-mannitol, D-sorbitol, Adonitol and D-arabitol


D- gluconic acid, Cellobiose, Palatinose, D- raffinose, Stachyose, D-trehalose, N-acetyl-D- glucosamine, Arbutin, Salicin, D-xylose and a-D- lactose+ D-xylose

Fumaric acid, methyl succinate, 2-keto-D- gluconic acid, D- gluconic acid, Dextrin, Cellobiose, Gentiobiose, Maltose, Maltotriose, D-melibiose, D- raffinose, Stachyose, Sucrose, D-trehalose, Turanose, N-acetyl-D-glucosamine, Arbutin and D- xylose

L-rhamnose, L-sorbose, Amygdalin, Arbutin, Salicin, D-maltitol, D-mannitol, D-sorbitol, adonitol, xylitol, l-erythritol, glycerol, tween 80, D- ribose, D- galactose+ D-xylose, 1,2-propanediol+ D-xylose

149

150 Based on Biolog reading result different yeast isolates were utilizing different carbon
151 sources. But all isolates utilized dextrin in common and L-aspartic acid, D-gluconic acid and
152 tween 80 were utilized more than seven isolates and the extant of utilizing were presented in
153 Figure 1.
154



155

156 **Figure 1. Number of yeast isolates utilizing the same carbon in the microplate.**

157 Based on the carbon utilization and assimilation the yeast was clustered using past analysis
158 software as indicated in Figure 2. The cluster shows six clusters which resembles similar
159 carbon utilization and assimilation characteristics.




Figure 2. Clustering of yeast isolates by carbon utilization using past

The percentage occurrence on culture media recorded as, 5% *Pichia fermentans*, 15% *Pichia* spp., *Rhodotorula aurantiaca* B, 15% *Pichia fluxuum*, 10% *Candida humilis*, and, 10% *Cryptococcus albidus Var aerus* other 35% were not identified. Six yeast species were identified which is involved in dough fermentation by Omni Log identification system. Those selected isolates were identified using Biolog OmniLog identification system using Biolog OmniLog Micro plates. Six yeast species were identified by OmniLog identification system. From those identified yeast species by OmniLog 54% were similar in genus, 15% were similar at species level and 8% were different with yeast species which identified by molecular identification system and 23% were unidentified. Yeast Micro Plate has two different reactions: assimilation of carbohydrates and oxidation. Six yeast species which were identified by BioLog (*Pichia fermentans*, *Pichia* spp., *Rhodotorula aurantiaca* B, *Pichia fluxuum*, *Candida humilis*, *Trichosporon beigeli* B and, *Cryptococcus albidus Var aerus* have positive oxidative test for Dextrin, stachyose, D-galactose, glucose, L-aspartic acid, D-gluconic acid, D-melibiose, sucrose, turanose, D-mannitol, tween 80, D- arabitol, D-meleitose, L-sorbose, Xylitol, Glycerol, Gentiobiose, Sucrose, Salicin, Maltotriose, L-sorbose carbon sources.

4. CONCLUSION

The study indicated that the fermenting teff dough is composed of different yeast groups. The identification of these isolates could possibly contribute pieces of information needed for understanding and verification of microbial consortia involved in the course of injera dough fermentation. And those microorganisms were appeared from 48h to 72h but some were appearing in early stage and others were continue up to the last stage (i.e. 72h).

188 **ACKNOWLEDGEMENTS**

189

190 I would like to acknowledge the following institutions: - Holeta Agricultural Biotechnology
191 Research Institute and microbial laboratory at Ethiopian Biodiversity Institute.

192

193

194 **COMPETING INTERESTS**

195

196 I declare that this research and the information presented in are my own and has been
197 generated by me as the result of my own original experimental research.

198

199

200 **5. REFERENCES**

201

- 202 1. Askal D.and Kebede A. Isolation, characterization and identification of lactic acid
203 bacteria and yeast involved in fermentation of Teff (*Eragrostis teff*) Batter. Intern.
204 J. 2013; 1:36-44.
- 205 2. Bakheit H. A. Technological Properties of Lactic Acid Bacteria Isolated from Different
206 Fermented Foods Faculty of Agriculture - University of Khartoum. Unpublished
207 Dissertation Submitted to the University of Khartoum in partial fulfillment of the
208 requirement of the degree of M.Sc. in Food Science and Technology. 2008.
- 209 3. Bekatorou, A., Food grade yeasts. Food. Technol. J. Biotechnol., 2006; 44(3), 407–
210 415.
- 211 4. Biolog, YT Microplate instruction for use. Biolog, Inc. 1993.
- 212 5. Birhanu G., Zerihun T. and Belay T. Isolation and characterization of yeast species
213 from ensete ventricosum product; Kocho and Bulla collected from Angacha
214 district. Interna. J. Adva. Biolo. Biomed. Rese. 2016; 5: 245-251.
- 215 6. Brhanu A.G. Involvement of lactic acid bacteria in the fermentation of teff (*Eragrostis*
216 *teff*), an Ethiopian fermented food. J. Food Sci. 1985;50: 800-801.
- 217 7. Bultosa, G., Hamaker B. R. and Be-Miller J. N. An SEC-MALLS study of molecular
218 features of water soluble amylopectin and amylose of teff [*Eragrostis teff* (Zucc.)
219 Trotter] starches. Starch/Stärke, 2008; 60: 8-22.
- 220 8. Hiwot, T. A., Ashore Z. W. and Gulelat D. H. Preparation of Injera From Pre-
221 Fermented Flour: Nutritional and Sensory Quality. Inter. J. of Science Innov.
222 Disco. 2013; 3:165-175.
- 223 9. Kurtzman, C.P. and Fell, J.W. "Yeast Systematics and Phylogeny—Implications of
224 Molecular Identification Methods for Studies in Ecology". Biodiversity and
225 Ecophysiology of Yeasts, Biodiversity and Ecophysiology of Yeasts (2006. p. 11-
226 30.
- 227 10. Mogessie A. A Review on the Microbiology of Indigenous Fermented Foods and
228 Beverages of Ethiopia. Ethiop. J. Biol. Sci. 2006; 5: 189-245.
- 229 11. Stewart, B.R. and Getachew A. Investigation of the nature of Injera. Econ. Bota.
230 1962;16: 127-30.
- 231 12. Whitman W.B. Bergey's Manual of Systematic Bacteriology. Second Edition Volume
232 Three the Firmicutes. University of Georgia Athens. 2009.
- 233 13. Yetneberk, S., Rooney, L.W. & Taylor, J.R.N. Improving the quality of sorghum
234 injera by decortication and compositing with teff. J. Sci. Food and Agri., 2005;
235 85:1252–1258.