

Preliminary Assessment of Fauna Species Diversity in Ipinu Igede Community Range Forest in Oju Local Government of Benue State, Nigeria.

G. O. Yager^{1*}, A. A. Alarape² and O. A. Onuwa

¹Department of Wildlife and Range Management, University of Agriculture, Makurdi, Benue State, Nigeria

²Department of Wildlife and Ecotourism Management, University of Ibadan, Ibadan, Nigeria.

Corresponding Author: gbrlyager@gmail.com.

Authors' contributions

This work was carried out in collaboration between all authors. Authors GOY and OAO designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors GOY and OAO managed the analyses of the study. Author AAA managed the literature searches. All authors read and approved the final manuscript.

ABSTRACT

Background and Objectives: Appraisal of fauna species which form an integral part of range ecology, in rangeland based protected areas is necessary before any meaningful conservation work can commence. It was aimed at assessing the composition of fauna species in the community forest.

Methodology: Animal species were enumerated through direct on-site using four transects line of 2.0km by 10m broad distributed randomly, field observation and indirect indices. Data were analyzed using descriptive statistics (table, figures and plates).

Results: Fifty-six species of wild animals from 40 families were found in the study area. There were 21 species of mammals from 15 families, 6 species of reptiles from 6 families and 29 species of birds from 19 families respectively. A total of 1,419 sightings were recorded. The most abundant animal species found in the area were *Epixerus ebii*, *Eidolon helvum*, *Chlorocebus tantalus*, *Papio anubis*, *Philothemus irregularis*, *Musophaga violacea*, *Numidia meleagris* and *Francolinus bicalcaratus*. The status of most mammals was Low risk-conservation dependent as applicable to reptiles. All the birds are known to breed in the reserve. Transect C had the highest diversity index (0.0130) and transect A had the lowest animal diversity index (0.0061) but with the highest species count of thirty six (36) and individual animal species sighted (454).

Conclusions: The community forest support unique fauna species making it significant in terms of conservation and scientific interest and has to be protected through conservation

awareness and community participation to conserve the current state and enhanced its range productivity.

Keywords: Range ecology, fauna, composition, status, Nigeria

1. INTRODUCTION

The survival and continuity of many endemic, rare and threatened species found in a given rangeland depend on sustainable conservation through its assessment to determine its current status. The potential of range forest habitation of wild animals is grossly unexplored in many areas across Nigeria, especially local community forests [1]

According [2], survey of both flora and fauna species which form an integral part of animal and forest ecology in wildlife based protected areas is necessary before any meaningful conservation work can commence. Fauna resources are the entire wild animal of any particular region or ecosystem [3]. These wild animals can be found in all ecosystems including forests, grasslands, plains, wetlands and deserts [4]. Fauna species assessment has more concentration to national parks and game/wildlife parks. However, many local rangeland communities support unique flora and fauna species making them important in terms of conservation and scientific interest.

Approach to species listing is an important initial stage in the collection of appropriate data necessary for effective management and conservation of animals and plants in a protected rangeland [5]. Therefore, knowledge of the species composition of a protected rangeland, their status and how they relate with other components of the habitat is highly essential and as well indicate the status of most fragile, threatened species. Insight into species list and status is becoming increasingly important as conservators and rangeland managers are tasked to assist conservation biologists to construct informed management plans for endangered species. This has become critical because most fauna species live in tropical forest which is increasingly been impacted by human modification and natural occurrences [6, 7, 8].

The status of the population of any individual species is crucial information to the wildlife ecologists, because this information determines individual fitness to its environment and also predicts their ultimate success or failure [9]. [10] report that, wildlife is increasingly being regarded as removable resources and man mostly is known for his high taste for exploiting its populations in the environment hence, their habitation to various rangeland and status ought to be monitored to ensure proper utilization of their habitat.

Ipinu Igede Community Forest in Oju Local Government of Benue State is one of the reserve that is rich in biodiversity, though had no appreciable ecological survey of the

resources, hence, the dearth of information necessary for the development of the reserve. The area has suffered from activities of illegal logging operators and hunting thereby threatening important flora and fauna species also the quest for a local fauna database and the alarming rate of species loss informed the need for wildlife based inventory in the study area. Thus there is need to appraise the composition of fauna species using diversity indices to ascertain the present status of fauna species of the community forest. The objective of the study is therefore to quantify fauna species composition and abundance using diversity indices and ascertain the status of the species. It is quiet obvious that baseline data generated from the study will promote effective conservation of biodiversity and management plan within the communal forest.

2. MATERIAL AND METHOD

2.1 The Study Area

The research was carried out at Ipinu Igede Community Forest Reserve in Oju Local Government Area of Benue State, Nigeria. The community forest is an ancestral heritage site for Igede people of Benue State stretching through three communities; Oyinyi, Andibilla and Uchenyim. The forest contains relicts of traditional worship practices in the area, although, the traditional religious worship practices are no longer strong and appreciated due to acceptance of Christianity. However, the laws and taboos governing the forest are still observed by the people of Igede.

The forest which is located in the Southern Guinea savanna belt comprise of both hilly and lowland part and lies between Longitude 8° 25' 0" E and 8° 41' 67" E and Latitude 6° 51' 0" N and 6° 85' 0" N [11]. It has an area of approximately 4 km² on a fairly flat land drained by four main seasonally flowing streams (Abadehe, Otuhukwu, Ekpaa and Ugbunwu) which are tributaries to River Ogbugwu. The mean annual rainfall is between 1200mm and 1500mm. The mean annual temperature is 30° C. Relative Humidity is between 60% and 80% wet but decreases in the early months of dry season.

It is a derived tropical rainforest characterized by luxuriant vegetation with high composition of riparian forest, of the large trees are *Cola gigantean*, *Elaeis guinensis*, *Ficus exasperate*, *Khaya spp*, *Afzelia africana* [11]. Dominant herbaceous species include *Sphenoclea zeylanica*, *Pentodon pentandrus*, *Ageratum conyzoides*, *Nymphaea lotus* and *asystasia gangetica*. The area has relatively abundant faunal resources; commonly sighted mammals are the primates (baboons and monkeys), bushbuck, oribi, grass cutter, squirrel and common duiker. Reptiles were alligator and snakes. Birds include guinea fowl, francolin,

village weavers, African dwarf-king fisher, African grey hornbill, Yellow billed kite and Abyssinian roller.

2.2 Data Collection Techniques

2.2.1 Species Diversity and Status

Species list and diversity was determined by direct observation along four transects of 2.0km by 10m broad (0.1ha) distributed randomly as described by [12] and indirect indices as well as through information from hunters and bush meat processing and selling centers.

Survey was carried out in the morning hours between 6:00 to 9:00am and early evening time between 4:00 to 7:00 pm. This was to ensure counting of even the shyest animal species as the period coincides with the time the animals are most likely to search for water and preys or graze on land [13].

Status assessment of Mammals, Birds and Reptiles was based on the information from hunters and forest protected agent and follows [14] and IUCN (International Union for Conservation of Nature) Red list.

2.3 Data Analysis

Descriptive statistics (tables, chart and figures) were used to analyze species lists of mammals, reptiles and birds.

2.3.1 Status Categories of Mammals, Reptiles and Birds

Categories outlined by [14] were used to assign the status of mammals, reptiles and of birds. This is as follows;

Vu = Vulnerable (Likely to become endangered if the factor that is posing threat persists).

LR/ cd = Low risk-conservation dependent (Species in no immediate danger but survival will depend on implementation of effective conservation measures in the community forest).

NT = Near threatened (species is approaching the threshold of vulnerability)

EN = Endangered (species is unlikely to survive if the factor that is posing threat persists).

RB = Resident breeder

R {B} = Resident but breeding not approved.

132 PM = Palearctic migrant

133 AFM = Migrates within Nigeria

134 DD = Data deficient

135 2.3.2 Diversity Indices

136 Diversity indices were calculated for each transect using Simpson's diversity index, which is
137 a measure of heterogeneity of a site taking into consideration the number of species and
138 density of individual species [15, 16]. The index is expressed as;

$$139 \quad I = \frac{q \sum n(n-1)}{N(N-1)}$$

140 Where I = Simpson diversity index.

141 N = total number of individuals enumerated.

142 q = number different species enumerated.

143 n = number of individuals of species enumerated.

144

145 3. RESULTS

146 Fifty-six species of vertebrates (wild animals) belonging to 42 families were
147 identified in the study area. They belong to three classes of Mammalia, Reptilia and Aves.
148 Twelve species of mammals were identified through direct sighting while 9 species was
149 through their signs and activities as well as interviews of hunters and bush meat processing
150 and selling centers. Three species of reptiles were identified through direct sighting while 3
151 were indirect assessment. All the Bird species were identified through direct sighting.

152 Majority of the identified mammal species were in the category of LR/cd, followed
153 Vu and some NT approaching the threshold of vulnerability. Most of the Reptile species were
154 fall within the LR/cd category. Almost all the identified birds' species are resident breeders in
155 the forest (RB). Some of the species identified and fecal droppings are presented in plate 1 to
156 6.

157 Wild Animal Species Distribution and Abundance Across the Transects

158 The total numbers of individual species recorded were 1,419. The class Aves had the
159 highest frequency (974) 68.6% followed by Mammalia (429) 30.2% and Reptilia (16) 1.1%

(Figure. 1). The total numbers of animals occurrence recorded for the various transects (A, B, C and D) were 454, 332, 294 and 339 respectively (Table 2 and Figure 2). The species with the highest abundance of class mammalian was *Epixerus ebii* (67.60%) followed by *Eldolon helvum* (15.30 %) and the least was *Tragelaphus scriptus* (0.23%). For class reptilian; the dominant species were *Philothamus irregularis* (56.25%) followed by *Elgaria coerulea* (31.25%) and the least was *Naja melanoleuca*. Class aves was *musophaga violacea* (22.59%) followed by *Numidia meleagris* (10.37%), *Crinifer piscator* (10.27%) and the lowest was *Accipiter africana* (0.10%) respectively. However, there were 18 constant species present in all the transects. This include *Arvicanthis niloticus*, *Epixerus ebii*, *Acrocephatus rufescens*, *Centropus sensgalensis*, *Colius striatus*, *Coracias abyssinicus*, *Crinifer piscator*, *Euplectes franciscannus*, *pternistis bicalcaratus*, *Lamprotornis nitens*, *Lanchura cucullata*, *Musophaga violacea*, *Numidia meleagris*, *Phynonotus barbatus*, *Poicephalus senegalus*, *Streptopelia semitorquata*, *spilopelia senegalensis* and *Vidua macroura*

The Simpson diversity indices of animal species showed that transect C had the highest diversity index (0.0130) and the second highest species count of thirty four (34). Transect A, on the other hand, had the lowest animal diversity index (0.0061) with the highest species count of thirty six (36) as well as individual animal species sighted (Table 3).

Table 1: Species List, Mode of Identification and Status of Mammals, Reptiles and Birds in Ipinu Igede Community Forest

Species				Mode of Identification				Status
S/N	Common Names	Scientific Names	Family	DS	IND	INH	PC	Category
Mammals								
1	Common duiker	<i>Sylvicapra grimmia</i>	Bovidae	X	x	X	X	VU
2	Bushbuck	<i>Tragelaphus scriptus</i>	Bovidae	X	-	X	X	LR/cd
3	Oribi	<i>Ourebia ourebi</i>	Bovidae	X	-	X	X	LR/cd
4	Waterbuck	<i>Kobus ellipsiprymnus</i>	Bovidae	-	-	X	-	LR/cd
5	Pale fox	<i>Vulpes pallida</i>	<i>Canidae</i>	-	x	X	-	LR/cd
6	Tantalus monkey	<i>Chlorocebus tantalus</i>	Cercopithecidae	X	x	X	-	LR/cd
7	Olive baboon	<i>Papio Anubis</i>	Cercopithecidae	X	-	X	-	LR/cd
8	Four-toed hedgehog	<i>Atelerix albiventris</i>	Erinaceidae	-	-	X	-	LR/cd
9	Allen's galago	<i>Sciurocheirus gabonensis</i>	Galagidae	-	-	X	-	LR/cd
10	Spotted hyena	<i>Crocuta crocuta</i>	Hyenidae	-	-	X	-	Vu
11	Crested Porcupine	<i>Hystrix cristata</i>	Hystriidae	-	-	X	-	Vu
12	Pygmy rabbit	<i>Brachylagus idahoensis</i>	Leporidae	X	-	X	X	LR/cd
13	Giant ground pangolin	<i>Smutsia gigantea</i>	Manidae	-	-	X	-	Vu
14	African grass rat	<i>Arvicanthis niloticus</i>	Murinae	X	-	X	X	LR/cd
15	Forest giant	<i>Cricetomys emini</i>	Nesomyidae	-	x	X	X	LR/cd

16	pouched rat African straw-coloured fruit bats	<i>Eidolon helvum</i>	Pteropodidae	X	-	X	-	NT
17	Western palm squirrel	<i>Epixerus ebii</i>	Sciuridae	X	-	X	-	LR/cd
18	Striped ground squirrel	<i>Xerus erythropus</i>	Sciuridae	X	-	X	-	LR/cd
19	Greater cane rat	<i>Thryonomys swinderianus</i>	Thryonomyidae	X	-	X	X	LR/cd
20	African civet	<i>Civettictis civetta</i>	Viverridae	X	-	X	X	LR/cd
21	Common genet	<i>Genetta genatta</i>	Viverridae	-	-	X	-	LR/cd
Reptiles								
22	Northern alligator lizard	<i>Elgaria coerulea</i>	Alligatoridae	X	-	-	X	LR/cd
23	Northern green bush snake	<i>Philothemus iregularis</i>	Colubridae	X	-	X	-	LR/cd
24	Black and white spitting cobra	<i>Naja siamensis</i>	Elapidae	X	-	X	-	Vu
25	Royal python	<i>Python regius</i>	Pythonidae	-	-	X	-	LR/cd
26	Savannah monitor lizard	<i>Veranus exanthematicus</i>	Veranidae	-	-	X	X	LR/cd
27	Red adder	<i>Bitis rubida</i>	Viperidae	-	-	X	-	LR/cd
Aves (Birds)								
28	Yellow billed kite	<i>Milvus aegyptius</i>	Accipitridae	X	-	-	-	RB
29	Black kite	<i>Milvus migrans</i>	Accipitridae	X	-	-	-	RB
30	Goshawk hawk	<i>Accipiter africana</i>	Accipitridae	X	-	-	-	RB
31	African dwarf-king fisher	<i>Ispidina lecontei</i>	Alcedinidae	X	-	-	-	RB
32	African grey hornbill	<i>Tockus nasutus</i>	Bucerotidae	X	-	X	-	RB
33	Little ringed plover	<i>Charadrius dubius</i>	Charadriidae	X	-	-	-	RB
34	Common ringed plover	<i>Charadrius hiaticula</i>	Charadriidae	X	-	-	-	RB
35	Speckled mousebird	<i>Colius striatus</i>	Coliidae	X	-	-	-	RB
36	Laughing dove	<i>Spilopelia senegalensis</i>	Columbidae	X	-	X	-	RB
37	Mourning collared dove	<i>Streptopelia decipiens</i>	Columbidae	X	-	-	-	RB
38	Yellow eyed-pigeon	<i>Columba eversmanni</i>	Columbidae	X	-	-	-	RB
39	Abyssinian roller	<i>Coracias abyssinicus</i>	Coraciidae	X	-	-	-	RB
40	Piapac	<i>Ptilostomus afer</i>	Corvidae	X	-	-	-	RB
41	Senegal coucal	<i>Centropus sensgalensis</i>	Cuculidae	X	-	X	-	RB
42	Black throated coucal	<i>Centropus leucogaster</i>	Cuculidae	X	-	-	-	RB
43	Bronze manikin	<i>Lanchura cucullata</i>	Estrildidae	X	-	-	-	RB
44	Violet turaco	<i>Musophaga violacea</i>	Musophagidae	X	-	X	X	RB
45	Western plantain eater	<i>Crinifer piscator</i>	Musophagidae	X	-	-	-	RB
46	Double-Spurred francolin	<i>Pternistis bicalcaratus</i>	Phasianidae	X	x	X	X	RB
47	Helmeted guineafowl	<i>Numida meleagris</i>	Phasianidae	X	-	X	X	RB

48	Green woodhoopoe	<i>Phoeniculus purpureus</i>	Phoeniculidae	X	-	-	-	RB
49	Common bulbul	<i>Phynonotus barbatus</i>	Phynonotidae	X	x	-	-	RB
50	Northern Red bishop	<i>Euplectes franciscannus</i>	Ploceidae	X	-	-	-	RB
51	Village weaver	<i>Ploceus cucullatus</i>	Ploceidae	X	-	-	-	RB
52	Senegal parrots	<i>Piocephalus senegalus</i>	Poicephalus	X	-	-	-	RB
53	Cape starling	<i>Lamprotornis nitens</i>	Sturnidae	X	-	-	-	RB
54	Rufus cane warbler	<i>Acrocephatus rufescens</i>	Sylviidae	X	x	-	-	RB
55	Pin-tailed whydah	<i>Vidua macroura</i>	Viduidae	X	-	-	-	RB
56	coastal indigo	<i>Indigofera miniata</i>	Viduidae	X	-	-	-	RB

179 Field Survey, 2017

180 In the above table;

181 DS = Direct Sighting

182 IND = Indices (Animals sign and activities)

183 INH = Interview of hunters

184 PC = Bush meat processing and selling center

185 — = Absent

186 X = Present

187

188 **Table 2: Wild Animals Species Distribution and Abundance According to Transect**

S/N	Scientific Names	Common Names	Tran. A	Tran. B	Tran. C	Tran. D	Total	Abundance
Mammals								
1	<i>Arvicanthis niloticus</i>	African grass rat	4	3	5	1	13	3.03
2	<i>Chlorocebus tantalus</i>	Tantulus monkey	38	-	-	-	38	8.56
3	<i>Civettictis civetta</i>	African civet	2	-	1	-	3	0.79
4	<i>Eidolon helvum</i>	African straw-coloured fruit bats	38	2	-	26	66	15.38
5	<i>Epixerus ebii</i>	Western palm squirrel	110	40	59	81	290	67.60
6	<i>Brachylagus idahoensis</i>	Pygmy rabbit	1	-	1	-	2	0.47
7	<i>Ourebia ourebi</i>	Oribi	1	1	-	-	2	0.47
8	<i>Papio Anubis</i>	Olive baboon	-	-	-	4	4	0.93
9	<i>Sylvicapra grimmia</i>	Common duiker	4	-	-	-	4	0.93
10	<i>Thryonomys swinderianus</i>	Greater cane rat	1	1	1	-	3	0.79
11	<i>Tragelaphus scriptus</i>	Bushbuck	-	-	-	1	1	0.23
12	<i>Xerus erythropus</i>	Striped ground squirrel	1	1	1	-	3	0.79
Total			200	48	68	113	429	100%
Reptiles								
13	<i>Elgaria coerulea</i>	Northern alligator lizard	3	-	-	2	5	31.25
14	<i>Naja melenaleuca</i>	Black spitting cobra	1	-	1	-	2	12.50
15	<i>Philothemus irregularis</i>	Northern green bush snake	3	-	6	-	9	56.25
Total			7	-	7	2	16	100%
Aves								

16	<i>Accipiter africana</i>	Goshawk hawk	-	-	1	-	1	0.10
17	<i>Acrocephatus rufescens</i>	Rufus cane warbler	13	8	7	11	39	4.00
18	<i>Centropus leucogaster</i>	Black throated coucal	-	2	1	2	5	0.51
19	<i>Centropus sensgalensis</i>	Senegal coucal	12	12	14	17	55	5.65
20	<i>Charadrius hiaticula</i>	Common ringed plover	-	-	3	7	10	1.03
21	<i>Charadrius dubius</i>	Little ringed plover	-	-	2	-	2	0.20
22	<i>Milvus migrans</i>	Black kite	4	-	-	4	8	0.82
23	<i>Colius striatus</i>	Speckled mousebird	7	11	9	1	28	2.87
24	<i>Coracias abyssinicus</i>	Abyssinian roller	2	1	2	8	13	1.33
25	<i>Crinifer piscator</i>	Westrn plantain eater	26	39	19	16	100	10.27
26	<i>Euplectes franciscannus</i>	Northern red bishop	9	4	4	1	18	1.85
27	<i>Pternistis bicalcaratus</i>	Double-Spurred francolin	14	13	12	19	58	5.95
28	<i>Ispidina lecontei</i>	African dwarf-fish king	-	-	4	13	17	1.74
29	<i>Lamprotornis nitens</i>	Cape starling	4	6	3	2	15	1.54
30	<i>Lanchura cucullata</i>	Bronze manikin	11	9	7	6	33	3.39
31	<i>Milvus aegyptius</i>	Yellow billed kite	4	6	1	-	11	1.13
32	<i>Musophaga violacea</i>	Violet plantain eater	57	70	48	45	220	22.59
33	<i>Numidia meleagris</i>	Helmeted guineafowl	20	32	26	23	101	10.37
34	<i>Phoeniculus purpureus</i>	Green woodhoopoe	2	-	11	-	13	1.33
35	<i>Phynonotus barbatus</i>	Common bulbul	16	8	18	11	53	5.44
36	<i>Piocephalus senegalus</i>	Senegal parrots	5	3	6	1	15	1.54
37	<i>Lamprotornis nitens</i>	Village weaver	4	5	4	-	13	1.33
38	<i>Ptilostomus afer</i>	Piapi	-	2	-	2	4	0.41
39	<i>Streptopelia decipiens</i>	Mourning collared dove	12	-	2	3	17	1.75
40	<i>Streptopelia semitorquata</i>	Red eyed pigeon	8	11	5	4	28	2.87
41	<i>Spilopelia senegalensis</i>	Laughing dove	3	11	5	9	28	2.87
42	<i>Tockus nasutus</i>	African grey hornbill	11	27	-	13	51	5.24
43	<i>Indigofera miniata</i>	Costal indigo	2	-	2	-	4	0.41
44	<i>Vidua macroura</i>	Pin-tailed whydah	1	4	3	6	14	1.44
Total :			247	284	219	224	974	100%
ΣTotal: Animals/Reptiles/Aves			454	332	294	339	1419	-

Field Survey, 2017

189
190




Fig. 1. Class Distribution of Wild Animals in Ipinu Igede Community Forest




Fig. 2. Distribution of Wild Animals According to Transect

Table 3: Simpson Diversity Index of Wild Animal Species in the Transect

Transect	Individual Species (n)	Total of Species (N)	Diversity Index
A	36	454	0.0061
B	27	332	0.0064
C	34	294	0.0130
D	29	339	0.0071



Plate1: *pternistis bicalcaratus*



Plate 2: *Xerus erythropus*



Plate 3: *Thryonomys swinderianus*



Plate 4: *Veranus niloticus*



Plate 5: Feather of *Musophaga violacea*



plate6: dropping of *Sylvicapra grimmia*

4. DISCUSSIONS

The relatively high population of vertebrates' species (wild animals) found in the area is typical of West African taxa [12]. The list of species surveyed is for the understanding of faunal dynamics in the conservancy of any protected area. This is in line with the observation by [17] at Makurdi zoological garden. More so, the high incidence of *Epixerus ebii*, *Eidolon helvum*, *Arvicanthus niloticus* and some primates in the community forest may not be unconnected to the fact that the species are not accepted as meat by the people in the surrounding communities. Similar observations have been made by [5] at Sambisa game reserve. The relatively low status of some mammals and reptiles in the forest such as duiker,

spotted hyena, porcupine and pangolin suggests high incidence of poaching for meat and traditional medicine because of the very little effort being made to protect the resources of the forest. Snake species are of least concern however the community tends to dislike them; hence the quest to eliminate them from their surrounding environment could have being the possible cause. Generally, some wild animals have higher tolerance of hunting pressure than other because of their home range and their reproductive potentials. Some may be subjected to less hunting pressure because the taste and acceptance of their meat or their ease of preparation. Local techniques used in capturing some species also put them under varying pressures. Some animal species also response to vegetation structure that allow a clear view of their surrounding and enable them to move with speed and agility through the under growth, like ground squirrel make use of the forest edge and strip vegetation because they are not able to survive an arboreal life unlike the tree squirrel that dominate the area. Common duikers were also found in sparsely dense habitat.

Bird life in the study area is largely recorded in relation with trees ranging from the violet plantain eater, western plantain eater, helmeted guineafowl, African grey hornbill, rufus cane warbler, African dwarfking-fisher, double-spurred francolin, village weaver which normally winter around the streams. A large number of birds live on seeds, fruits, buds, and nectar or insects that are found in the arboreal environment. These include western plantain eater and African dwarfking-fisher respectively. The high bird species diversity in the area could be due to the fact that the area acts as a sanctuary from the degraded habitats surrounding it and nesting materials and availability of edible fruits bearing tress. This observation is in line with the report by [18] at GRA and Ankpa quarters Benue State.

In a comparative form, the total number of twenty-one mammalian species is just about 8.5% of 247 species reported for Nigeria [19]. The number is also lower than either of the 123 species reported for Guinea Savanna or 97 species for Sudan Savanna of Nigeria [19]. So, the species richness of the forest might not be unconnected with its size which is relatively small compared to the size of Guinea Savanna (473,904 km²) or Sudan Savanna (927,338km²). This observation agrees with [20] report, that species diversity is often affected by the size of habitat and that diversity is positively correlated with habitat size. Biodiversity assessment and conservation management purposes, distribution or pattern of occupancy is very important and this has been found to vary with different environmental location and condition for a given species.

5. CONCLUSION

Different levels of disturbance have different effects on animal diversity in the study sites. Reliable information on the status and trends of forest fauna resources help give decision makers the prospective necessary for orienting wildlife policies and programs. Domestication of animal species should be advocated both for poverty alleviation in the communal lands of the area, and for a balance to be maintained in the ecosystem.

Ethical Approval:

As per international standard or university standard, ethical approval has been collected and preserved by the authors.

REFERENCES

1. Ogunjemite BG, Ajayi B, Agbelusi EA. Habitat structure of chimpanzee communities in Nigeria: a comparison of sites. *Acta Zoological Sinica*. 2007; 53 (4); 579-588.
2. Daniel IE, Henry MI, Augustine UO. Preliminary assessment of tree species diversity in Afi Mountain Wildlife Sanctuary, Southern Nigeria, *Agriculture and Biology Journal of North America*. 2012; 3.12.486.492 ScienceHub, <http://www.scihub.org/ABJNA>
3. Wikipedia. <https://en.m.wikipedia.org/wiki/Fauna> Retrieved on 5th November, 2017 wildlife heritage of Africans. TRAFFIC Reports. www.traffic.org.
4. Ijeomah HM, Augustine UO, Damilola O. Analysis of Poaching Activities in Kainji Lake National Park of Nigeria. *Environment and Natural Resources Research*. 2012; 3(1). <http://dx.doi.org/10.5539/enrr.v3n1p51>
5. Mbaya YP, Malgwi H. Species list and status of mammals and birds in Sambisa game reserve, Borno State, Nigeria, *Journal of Research in Forestry, Wildlife and Environment*. 2010; 2 (1): 135-140.
6. Alamu LO, Agbey BO. Deforestation and endangered indigenous tree species in Southwest Nigeria: *International Journal of Biodiversity and Conservation* 2011; 3.7 :291-310.
7. Babagana G, Mohammed MA, Garba M. Environmental Impact of Natural Resources Exploitation in Nigeria and the way forward. *Journal of Applied technology in Environmental sanitation* 2012; 2, (2): 95-102.
8. Ahmed RK. Biodiversity loss: A threat to urban landscape. *International Research Journal of Arts and Social Sciences* 2013; 2. 6:144-149
9. Martin C. *The Rainforest of West Africa*. Birkhauser, London. 1990.
10. Idowu MA, Morenikeji OA. Wild Fauna Conservation in Nigeria. *Environment and Natural Resources Research*, 2015; 5(3); 98-108.
11. Okwoche SO. Assessment of tree species composition and diversity of Ipinu-Igede community forest in Oju L.G.A, Benue State, Nigeria. Unpublished Undergraduate project submitted to the Department of Forestry, Wildlife and Range Management, University of Agriculture Makurdi, Nigeria 2017; pp 10-35.
12. Osunsina IOO, Inah EI, Ogunjinmi AA, Onadeko SA, Osunsina O. Distribution and diversity of flora and fauna in International Institute of Tropical Agriculture (IITA)

- 292 forest and nature reserve, Ibadan. Oyo State. Nigeria. *Journal of Agriculture, Forestry*
 293 *and the Social Sciences* (JOAFSS) 2012; 10 (2): 289-302.
- 294 13. Adamu IA. An assessment of floristic composition of Kwiambana Game Reserve, A
 295 Ph.D. thesis, presented to the Department of Geography, Usmanu Danfodiyo
 296 University, Sokoto (Unpublished). 2006; 122p.
- 297 14. Ezealor HU. Critical Sites for Biodiversity Conservation in Nigeria. *Nigerian*
 298 *Conservation Foundation*, Lagos, Nigeria. 2002; 110pp.
- 299 15. Simpson EH. Measurement of Diversity, *Nature*; 1949; 163: 688.
- 300 16. Ojo LO. Data collection and analysis for biodiversity conservation: *In Practical of the*
 301 *Inception Meeting and Training Workshop on BRAAF Assessment and Monitoring*
 302 *Techniques in Nigeria*. Eds. B.A. Ola-Adams and L.O. Ojo National Committee on
 303 Man and Biosphere 1996; Pg. 142-145
- 304 17. Yager GO, Enefolo JO, Tyowua BT. Comparative study of fauna species diversity of
 305 Makurdi Zoological Garden, Benue State, Nigeria, *International Journal of*
 306 *Development and Sustainability* ISSN: 2186-8662 – www.isdsnet.com/ijds. 2017; 6
 307 (12): 2163-2172 ISDS Article ID: IJDS17112401
- 308 18. Egwumah PO, Iwar IM, Ogbonna L. A survey of the avi-fauna within Makurdi
 309 Metropolis of Benue State, Nigeria. *Journal of Research in Forestry, Wildlife and*
 310 *Environment*, 2009; 1 (1); 75-83.
- 311 19. Happold DCD. *The Mammals of Nigeria*; Oxford press, London. 1987.
- 312
- 313 20. Usher MB. *Conservation Biology; A Training Manual for Biological*
 314 *Diversity and Genetic Resources*. Rapoor- Vijay and White (Ed.) 1992; pp 80- 83.