

RESPONSE OF CHEMOLITHOTROPHIC NITROBACTER, NITROSOMONAS TO TOXICITY OF ORGANOPHOSPHATE AND PYRETHROID PESTICIDES.

ABSTRACT

Aim: To investigate the response of *Nitrosomonas* and *Nitrobacter* species to organophosphate and pyrethroid pesticides.

Study design: This study employs experimental design and statistical analysis of data and interpretation.

Place and duration of studies: Soil samples were obtained from university farm, Rivers State University, Port Harcourt, Nigeria. Samples were transported to the microbiology laboratory of Rivers State University immediately for microbiological and toxicity testing. Pesticides was gotten No.4 Ignatius Ajuru University Road, St. John Campus, Aba Road Port Harcourt. The toxicity testing was done for the period of 28 days at room temperature.

Methodology: Standard microbiological techniques were used: toxicity testing procedures were carried out by preparing a stock culture of the pesticide based on manufactures directions (8ml into 1000ml of distilled water) from which the concentrations used for this research work were obtained 0%, 3.125%, 6.25%, 12.5%, 25% and 50% and tested on the soil samples for a period of 28 days. Samples were serially diluted and cultures were incubated at 35°C For 18 to 24 hours. LC₅₀ was determined using SPSS version 2.0

Results: The results indicate that logarithm mortality of *Nitrosomonas* and *Nitrobacter* species increases with increase toxicant concentration and exposure time for pyrethroid pesticide while decreases with increase toxicant concentration and exposure time for organophosphate pesticides. The median lethal concentration LC₅₀ of the pesticides increases in the following order: (Note: the higher the LC₅₀, the lower the toxic effect); pyrethroid pesticide on *Nitrosomonas* (53.1%) < organophosphate pesticide on *Nitrosomona* (47.9%), pyrethroid pesticide on *Nitrobacter* (53.5%) < organophosphate pesticide on *Nitrobacter* (47.5%).

Conclusion: The results revealed that different concentrations of the toxicants have both negative and positive effect on the survival rate of the test organisms which shows that the organophosphate pesticide can cause more harm to the environment affecting *Nitrosomonas* and *Nitrobacter* species that plays vital functions in nutrient fixation in the soil environment. While pyrethroid pesticides at appropriate concentrations can stimulate the growth of these organisms there by increasing the rate of nutrient fixation in the soil environment. But also, when these toxicants are misapplied they can cause harm to humans that would consume the crops.

Keywords: Pyrethroid pesticide, Organophosphate pesticide, Toxicity, *Nitrosomonas*, *Nitrobacter*, Nitrification.

Comment [L1]: Word of the first letter should be written in capital letters

Comment [L2]: words must be separated with each other

Comment [L3]: Word of the first letter should be written insmall letters

Comment [L4]: words must be separated with each other

1. INTRODUCTION

The tolerance abilities of chemolithotrophic bacteria (nitrifying bacteria) to some commonly used pesticides in the Rivers State University campus and environment cannot be under emphasized. Basically, the importance of chemolithotrophic bacteria in the biogeochemical cycle / environment has been noted beneficial [1]. Chemolithotrophic bacteria are known for their use of inorganic compounds as electron donor in their nutritional diet, their survival is dependent on the physicochemical condition of its immediate environment [1]. Pesticides are extensively used in agriculture as a part of pest control scheme. Due to the xenobiotic characteristics of pesticides, which may have led to adverse effects on the survival of beneficial soil microorganisms and their associated bio- transformation. Inactivation of nitrogen-fixing and phosphorus-solubilizing microorganisms have been observed in pesticide-contaminated soils. However, a few reports have revealed some positive effects of applied pesticides on soil health

2. MATERIAL AND METHODS

PLACE OF STUDY

Soil samples were collected from University farm, Faculty of Agriculture, Rivers State University, Port Harcourt Nigeria. With sterile trowel at the rhizosphere of legumous plant (mukuna beans) in a sterile polyethene bag and transported to the microbiology laboratory immediately. The used pesticides were gotten from No.4 Ignatius Ajuru University Road, St. John Campus, Aba Road Port Harcourt.

Microbiological Analysis

Isolation of *Nitrosomonas* species

Winogradsky Agar medium composition as modified by [2], was used: Agar Agar 15.0g, $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ 0.4g, NaCl 2.0g, K_2HPO_4 1.0g, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.5g, and $(\text{NH}_4)_2\text{SO}_4$ 2.0g were dissolved in 1000ml of distilled water and autoclaved at 121°C for 15 minutes (psi) after which was allowed to cool to about 40°C and the medium was poured into Petri dishes. Then, the medium was solidified before progress to the hot air oven to moisture. One (1) gms of soil was dissolved into 9ml of sterile distilled water and 10 fold serial dilution was done to 10^{-3} and an aliquot from each soil concentration were inoculated unto winogradsky agar and incubate aerobically for 2-3 days at room temperature ($30 \pm 2^\circ\text{C}$), greyish, mucoid, flat colonies revealed pear shaped, and Gram negative of *nitrosomonas*.

Confirmation of *Nitrosomonas* species

Suspected *Nitrosomonas* species were sub cultured on a fresh winogradsky agar medium and transferred into a broth containing Ammonium sulphate and Sodium nitrate and incubated at about ($30 \pm 2^\circ\text{C}$) for 2-3 days. 1ml of sulfanilic acid, dimethylnaphthalamine and zinc dust was added was added to the medium after 2days of incubation. Red coloration indicated by nitrate production from ammonia sulphate was a confirmation of *nitrosomonas* species.

Isolation of *Nitrobacter* species

Winogradsky Agar medium composition as modified by [2] was used: Agar Agar 15.0g, NaNO_2 0.05g, Na_2CO_3 1g, NaCl 0.3g, K_2HPO_4 0.5g, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.02g, ZnCl_2 0.03g and $\text{FeSO}_4 \cdot 6\text{H}_2\text{O}$ 0.02g were dissolved in 200ml of distilled water and autoclaved at 121°C for 15 minutes (psi) after which was allowed to cool to about 40°C and the medium was poured into Petri dishes. Then, the medium was solidified before progress to the hot air oven to moisture. One(1)gms of soil was dissolved into 9ml of sterile distilled water and 10 fold serial dilution was done to 10^{-3} and an aliquot from each soil concentration were inoculated unto winogradsky agar and incubate aerobically for 2-3 days at room temperature ($30 \pm 2^\circ\text{C}$), greyish, mucoid, flat colonies revealed pear shaped, and Gram negative of *Nitrobacter*.

Confirmation of *Nitrobacter* species

Suspected *Nitrobacter* species were sub cultured on a fresh winogradsky agar medium and transferred into a broth containing nitrite carbonate medium and incubated at about ($30 \pm 2^\circ\text{C}$) for 2-3 days. 5 drops of Griess illosvay's reagent was added to the medium after 2days of incubation. Absent of purplish colour indicate a positive result for *Nitrobacter* species, further confirmation was done by diphenylamine. Cherry red indicated presence of *Nitrobacter*.

Comment [L5]: Word of the first letter should be written in capital letters

Preparation stock toxicant

The stock toxicant was prepared based on manufacturer direction (800 ml of pesticides into 100 liters of water). The toxicant was prepared, with a volume of 8ml, pesticides transferred into 1litre of distilled water from which the concentrations were obtained from.

Toxicity test procedures

The toxicants were prepared aseptically by using different concentrations: as 3.125%, 6.25%, 12.5%, 25% and 50% respectively of the toxicant. These concentrations were obtained aseptically by transferring 3.125ml, 6.25ml, 12.5ml, 25ml, and 50ml of the different pesticides stock solution into 96.8ml, 93.75ml, 87.5ml, 75ml, 50ml, of sterile distilled water respectively. The toxicity test procedures was done by using 12 clay pots containing 1.5kg of oven sterilized soil, 10ml of bacteria (*Nitrobacter* and *Nitrosomonas spp*) was added separately and each toxicant concentration were added separately into different clay pots and a control experiment was done without inoculation of pesticides. One grams of soil sample from all concentrations was serially diluted and an aliquot from 10^{-3} dilution was used for inoculation using spread plate techniques on winogradsky media after 1, 7, 14, 21 and 28 days respectively and was incubated for 2-3 days at room temperature ($37\pm 2^{\circ}\text{C}$).

Toxicity test of bacteria (*Nitrobacter* and *Nitrosomonas spp*) in pesticides

The percentage log survival of *Nitrobacter* and *Nitrosomonas* species isolated in the pesticides polluted soils were calculated according to formula used by [3]. The percentage log survival of the bacteria isolates in the soil was calculated by obtaining the log of the count in toxicant concentration, divided by the log of the count in the zero toxicant concentration and multiplying by 100. Thus:

$$\text{Percentage (\%)} \log \text{ survival} = \frac{\text{Log } C}{\text{Log } c} \times 100$$

Where Log C = Logarithm count in each toxicant concentration, log c = Logarithm count in the control (zero toxicant concentration).

$$\text{Percentage (\%)} \log \text{ mortality} = 100 - \% \log \text{ survival}$$

3. RESULTS AND DISCUSSION

The logarithm counts of *Nitrosomonas* and *Nitrobacter* species revealed the response of these bacteria to organophosphate and pyrethroid pesticides are revealed in Tables 1 and 2 respectively. Percentage logarithm mortality of the counts are presented in the figures below.

The result obtained from this study revealed that pesticides can inhibit nitrification process as well as encourage it by *Nitrobacter* and *Nitrosomonas* species. Similar observation have been reported [4,5,6]. An increase in the percentage logarithm of mortality of *Nitrosomonas* and *Nitrobacter* in in treated with organophosphate pesticide after 28days of exposure to the toxicant concentrations, and an increase in percentage logarithm of survival rate of *Nitrosomonas* and *Nitrobacter* species in soil treated with pyrethroid pesticide after 28days of exposure to the toxicant concentrations were observed. (figs. 1to5) respectively. This study also revealed that the toxicant (organophosphate pesticide) is more toxic to the organisms than pyrethroid pesticide. This may be as a result of its chemical composition and at same time its degradability by this organisms. The site of action of any toxicant depends on the nature of the toxicant.

Table 1. log count of *Nitrobacter* species with organophosphate and pyrethroid pesticides

Pyrethroid pesticide + <i>Nitrobacter</i>						Organophosphate pesticide + <i>Nitrobacter</i>					
CONC/DURATION	1	7	14	21	28	(Days)	1	7	14	21	28
0%	5.68	5.75	5.77	5.81	5.84		6.0	5.91	5.83	5.74	5.72
3.125%	5.72	5.78	5.82	5.85	5.87		5.97	5.86	5.80	5.72	5.68
6.25%	5.74	5.82	5.84	5.88	5.90		5.92	5.79	5.66	5.60	5.57
12.5%	5.76	5.85	5.87	5.89	5.91		5.87	5.70	5.60	5.51	5.43
25%	5.79	5.87	5.89	5.92	5.94		5.83	5.65	5.52	5.50	5.41
50%	5.81	5.94	5.95	5.96	5.97		5.81	5.61	5.48	5.45	5.38

Comment [L6]: which pesticides used in the research should be written

Comment [L7]: words must be separated with each other

Comment [L8]: Word of the first letter should be written in capital letters

113

114 Table 2. log count of *Nitrosomonas* species with organophosphate and pyrethroid pesticides

Pyrethroid pesticide + <i>Nitrosomonas</i>						Organophosphate pesticide + <i>Nitrosomonas</i>					
CONC/DURATION	1	7	14	21	28	(Days)	1	7	14	21	28
0%	5.74	5.79	5.81	5.87	5.90		6.10	6.0	5.93	5.86	5.83
3.125%	5.77	5.81	5.86	5.89	5.93		6.08	5.98	5.88	5.85	5.79
6.25%	5.79	5.85	5.87	5.91	5.96		5.99	5.88	5.82	5.81	5.76
12.5%	5.86	5.88	5.90	5.92	5.97		5.88	5.81	5.76	5.74	5.71
25%	5.88	5.90	5.93	5.95	5.98		5.80	5.75	5.71	5.66	5.62
50%	5.93	5.97	5.98	5.99	6.01		5.72	5.65	5.63	5.60	5.57

116

117 Table 3. Median lethal conc. (LC₅₀) from percentage log mortality of pyrethroid on *Nitrosomonas*

CONC.	%mortality	Mean of mortality	Conc. Diff	Sum conc.diff. x mean mortality
0	-	-	-	-
3.125%	-6	-1.2	3.125	-3.75
6.25%	-15	-3	3.125	-9.38
12.5%	-23	-4.6	6.25	-28.75
25%	-28	-5.6	12.5	-70
50%	-40	-8	25	-200 =-311.9

118

119 LC₅₀ = 50 - (-) $\frac{311.9}{100}$

120

121 LC₅₀ = 50+3.1122 LC₅₀ = 53.1

123

124

125 Table 4. Median lethal conc. (LC₅₀) from percentage log mortality of pyrethroid on *Nitrobacter*

CONC.	%MORTALITY	MEAN OF MORTALITY	CONC. DIFF.	SUM OF CONC.DIFF* MEAN MORTALITY
0%	-	-	-	-
3.125%	-10	-2	3.125	-6.25
6.25%	-21	-4.2	3.125	-13.13
12.5%	-24	-4.8	6.25	-30
25%	-31	-6.2	12.5	-77.5
50%	-44	-8.8	25	-220 = -346.9

126

127 Median lethal conc (LC₅₀) from percentage log mortality of pyrethroid on *Nitrobacter*128 LC₅₀ = 50-(-) $\frac{346.9}{100}$

129 50-(-) 3.5 = 50 + 3.5

130 LC₅₀ = 53.5

131

132

133

134

135 Table 5. Median lethal concentration (LC₅₀) from percentage % log mortality of *Nitrobacter*

CONCENTRATION	%MORTALITY	MEAN%MORTALITY	CONC. DIFF.	SUM OF CON DIFF x MEAN%MORTALITY
0%	-	-	-	-
3.125%	3	0.6	3.125	1.88
6.25%	11.7	2.3	3.125	7.2
12.5%	21.1	4.2	6.25	26.37
25%	26.5	5.3	12.5	66.25
50%	30.1	6.02	25	150.5 = 252.2

136
137
138
139
140
141


$$\begin{aligned} \text{LC50} &= 50 - \frac{252.2}{100} \\ &= 50 - 2.5 = 47.5 \end{aligned}$$

Table 6. Median lethal concentration (LC50) from %percentage log mortality OF *Nitrosomonas*

CONCENTRATION	%MORTALITY	MEAN OF %MORTALITY	CONC. DIFF.	SUMOF CON.DIF. X MEAN%MORTALITY
0%	-	-	-	-
3.125%	2.6	0.52	3.125	1.63
6.25%	8	1.6	3.125	5
12.5%	14	2.8	6.25	17.5
25%	20.2	4.04	12.5	50.5
50%	26.3	5.3	25	132.5
				= 207.13

142
143
144
145

$$\begin{aligned} \text{LC50} &= 50 - \frac{207.13}{100} \\ &= 50 - 2.07 = 47.9 \end{aligned}$$



146
147
148
149

Fig.1. summary of median lethal concentration of pesticides (organophosphate and pyrethoid) on *Nitrosomonas* and *Nitrobacter* sp.




Fig2. % mortality rate of *Nitrobacter* when exposed to pyrethroid pesticide.




Fig3. % mortality rate of *Nitrosomonas* when exposed to pyrethroid pesticide




Fig4. % mortality rate of *Nitrobacter* when exposed to organophosphate pesticide




Fig 5: % mortality rate of *Nitrosomonas* when exposed to organophosphate pesticide

The percentage log survival of *Nitrosomonas* and *Nitrobacter* species during 28 days exposure period to soil treated with Organophosphate and Pyrethroid pesticides. (Tables 1 and 2) respectively shows that organophosphate pesticide exhibited little effect on the test organisms than pyrethroid pesticide. This may be due to the chemical composition of the toxicant. The percentage log mortality of *Nitrosomonas* and *Nitrobacter* species during 1, 7, 14, 21 and 28 days exposure periods to the different concentrations of the toxicants reveals that the mortality rate of organophosphate pesticide is higher than that of pyrethroid. Hence the results of this study suggest that organophosphate pesticide causes cell death which resulted in reduction of viable cell counts while pyrethroid pesticide was able to serve as carbon source for these organisms and thereby they were able to utilize them and proliferate which lead to increase in viable cell counts. The reduction of viable cell count of organophosphate pesticide may lead to inhibition of nitrification process during the 28days exposure period. Similar result was observed by [9], while pyrethroid pesticide lead to the increase in nitrification process because the organisms were able to utilize it as their sole carbon source, similar observation was detected by [8].

Nitrosomonas and *Nitrobacter* sp. Mortality expressed as median lethal concentration (LC_{50}) was used as indices to monitor toxicity [2], the sensitivity of these bacteria to the toxicity of the different concentration of the pesticides. (Tables 3 to 5) shows the median lethal concentrations (LC_{50}) of the pesticides used, which increased in the following order: pyrethroid on *Nitrosomonas* (53.5%) < organophosphate on *Nitrosomonas* (47.5%), pyrethroid on *Nitrobacter* (53.1%) < organophosphate on *Nitrobacter* (47.9%) (Note: the higher the LC_{50} , the lower the toxic effect and vise-versa). Conclusively organophosphate pesticide was more toxic to the organisms, even though most toxic to *nitrosomonas* (LC_{50} = 47.5%) having the most toxic effect while pyrethroid on *nitrosomonas* (LC_{50} =53.5%) having the lowest toxicity effect.

Conclusion and Recommendation

The results revealed that different concentrations of the toxicants have both negative and positive effect on the survival rate of the test organisms which shows that the organophosphate pesticide can cause more harm to the environment affecting *Nitrosomonas* and *Nitrobacter* species that plays vital functions in nutrient fixation in the soil environment. While pyrethroid pesticides at appropriate concentrations can stimulate the growth of these organisms there by increasing the rate of nutrient fixation in the soil environment. But also, when these toxicants are misapplied they can cause harm to humans that would consume the crops.

Therefore, it is recommended that pesticides should be applied according to manufacturer prescription and not misapplied, pyrethroid pesticides should be encouraged.

Reference

1. Tolli, J. and King, G.M. (2005). Diversity and structure of **Bacterial** chemolithotrophic communities in pine forest and Agro ecosystem soils. *Applied and Environmental Microbiology*.71 (12): 8411-8418.
2. Williams, J. O. and Ogolo, C. N. (2018). Response of *Aspergillus* species to the toxicity of domestic detergent's. *Nigeria journal of Microbiology*. 32(1):4334-4343.
3. Williams, J. O (2018). Assessment of the physicochemical and microbiological quality of palm oil mill effluent and soil in Aluu. *Asian Journal of Biology*.6 (3):1-11.
4. Monkiedje, A., Ilori, M. O., and Spiteller, M. (2002). Soil quality changes resulting from the application of the fungicides mefenoxam and metalaxyl to a sandy loam soil. *Soil Biology. Biochemistry*. 34, 1939–1948.
5. Tyess, D. L., Shea, P. J., and Parkhurst, A. M., (2006). Mineralization potential of atrazine and degradation intermediates from clustered characteristics in inoculated soils. *Soil Sediment Contamination*. 15, 87–102.
6. Das, A. C., and Mukherjee, D. (2000a). Soil application of insecticides influences micro- organisms and plant nutrients. *Applied Soil Ecology*. 14, 55–62.
7. Das, A. C., and Mukherjee, D. (2000b). Influence of insecticides on microbial transforma- tion of nitrogen and phosphorus in Typic Orchraguall soil. *Journal of Agricultural Food Chemistry*. 48, 3728–3732.

Comment [L9]: Word of the first letter should be written in capital letters

Comment [L10]: 1-Word of the first letter should be written in small letters

- 213 8. Sarnaik, S. S., Kanekar, P. P., Raut, V. M., Taware, S. P., Chavan, D. S., and Bhadbhade, B. J., (2006).
214 Effect of application of different pesticides to soybean on the soil microflora. *Journal of Environmental*
215 *Biology*. 27, 423–426.
- 216 9. Martinez-Toledo, M. V., Salermon, V., Rodelas, B., Pozo, C., and Gonzalez-Lopez, J. (1998). Effects of
217 the fungicide captan on some functional groups of soil microflora. *Applied Soil Ecology*. 7, 245–255.