

Original Research Article

HELMINTH PARASITES OF GOBIES FROM TWO CREEKLETS OF THE NEW CALABAR RIVER, RIVERS STATE, NIGERIA

Abstract

Every animal species is a definitive, paratenic or intermediate host to helminth parasites which may be species or site specific. This study was to investigate the intestinal helminth parasite prevalence of gobies (*Bostrychus africanus* and *Periophthalmus papilio*) collected from two creeklets of the New Calabar River. Fifteen samples of each species were collected from the two stations fortnightly between October 2016 and May 2017, measured and weighed. All fish was cut open and intestinal parasites were identified and counted. Physico-chemical parameters (temperature, dissolved oxygen (DO) and salinity) of the study areas were also measured in dry and rainy seasons. A total of 240 fish samples were examined of *B. africanus* (SL of 3 - 12 cm, weight of 30 - 36.1g), and *P. papilio* (SL of 5.9 - 15 cm, weight of 4 - 28.2g). 389 Ascarids were isolated from 25.4 % of the sampled fish. *B. africanus* had higher percentage prevalence and mean intensity (91.6 % and 23 in station 1, 21.8 % and 20.5 in station 2). *P. papilio* had a lower percentage prevalence of 0.7 % and mean intensity of 4 in station 1 and 5.7 %, 8 in station 2. The Ascarid was isolated from the stomach, small and large intestine of infested fish. The total length of infected and non-infected *B. africanus* from station 1 was different at $P < 0.05$ revealing that size affects the prevalence of the parasite. There was no significant difference in the relative condition factor (Kn) of infected and non-infected *B. africanus* and *P. papilio* from both stations indicating that the parasite did not affect the condition of the fish. Sex appeared to play a role in parasite infestation in *B. africanus*. The physico-chemical parameters of both stations were all within range for a tropical estuarine environment. The presence of ascarids is of interest as an ascarid (*Anisakis*) is a parasite of fish that is zoonotic.

Key words: *Ascaris*, *Periophthalmus*, *Bostrychus*, New Calabar River, Helminth, Gobies.

Introduction

Helminths are the most diversified group of parasites that infest freshwater and saltwater fishes. They have very high complex ecological inter-relationship with their hosts. Endoparasitic helminths possess an indirect life cycle, which usually involves one or more hosts. Fishes are known to be definitive, paratenic (transport) or intermediate host to these helminthic parasites [1]. Nematodes which are usually called roundworms are common in marine fishes [2]. Fish ingestion of larval helminth parasites occur regularly because of the abundance and diversity of these parasites in the aquatic ecosystem [3]. The health of fish is affected by parasites which make them susceptible to secondary infection by disease causing agents (e.g. bacteria, fungi and viruses). Par-

asites also compete for food, thereby depriving fish of essential nutrients and inhibiting growth leading to morbidity and mortality with consequent economic losses [4].

Gobies are fishes of the family Eleotridae, Gobiinae, Periophthalmidae and Gobiidae, which make up one of the largest fish families, comprising of more than 2,000 species in more than 200 genera. Most of them are relatively small, typically less than 10 cm in length. Some large gobies such as *Gobiodes* or *Periophthalmus* can reach over 30cm in length in exceptional cases [5]. The Eleotridae are usually known as sleeper gobies and they are small to medium-sized fishes, which are closely related to gobies in the family Gobiidae. Eleotrids are found worldwide in fresh water, estuarine and marine environments in tropical and warm temperate regions. A few small-sized species occur on rocky or coral reef environments. They are small to medium sized, ranging from 2 cm to 50 cm. Most species live on the bottom where they feed on benthic invertebrates especially crustaceans, a few species swim in the water column and feed on zooplankton. Periophthalmidae is the family of mudskippers. They are also small to medium-sized fishes which have cylindrical bodies and reach 27cm in length. They possess large heads and bulging eyes, which are very mobile and are adapted for easy vision in water and on land [6].

There are reports of helminthic parasite infestation in freshwater fishes from a few localities in Nigeria, but there appears to be a dearth of information on parasites of fish from the brackish water ecosystem. A few reports on brackish water fisheries are those on Lagos lagoon [7] and Warri river [8], the study of the gastrointestinal helminth parasites of Sciaenid species [9] and the threadfin *Polydactylus quadrifilis* [10] from Buguma Creek, Niger Delta. The aim of this study was to investigate the different types of helminth parasites found in gobiid fish from both creeks of the New Calabar River.

Materials and Methods

Description of the Study Area

The New Calabar River is in Rivers State and is a part of the Niger Delta basin which covers all the land between latitude 4°14'N and 5°35'N and the longitude 5°26'E and 7°37'E with a total area of 20,000 km² [11]. The vegetation is mangrove *Rhizophora mangle*, *R. racemosa*, *Avicennia* and *Nypa fruticans* (Nipa palm) which grows from a muddy substrate characterized by its foul smell. There are many anthropogenic activities that has led to environmental stress on this river [12]. This river is also connected to the Bonny River [13]. The present study is on the Rumuolumeni and the Bakana creeks (Fig 1) of the New Calabar

river and will provide information on the helminth parasites of two gobiid species belonging to two families.




Fig. 1: The Map of the Study Area, Rumuolumeni and Bakana Communities in Rivers State

Collection of samples

Sampling was for a period of eight months, from October 2016 to May 2017. Samples were collected from both creeks fortnightly between 6.00 am and 8.00 am using cone basket traps. Two hundred and forty (240) samples of each fish species (*B. africanus* and *P. papilio*) were collected from the intertidal region of Rumuolumeni and Bakana creeks. Fish samples were conveyed to the laboratory in estuarine water collected from the creeks. Physico-chemical parameters (temperature, dissolved oxygen (DO), salinity, turbidity and pH) were measured at both creeks using standard methods.

In the laboratory fish samples were weighed to the nearest g and length measured in cm to the nearest mm. They were examined for both ecto and endoparasites. For ectoparasites, the external surface of the fish was examined using a hand lens. For endoparasites, the fish was dissected laterally, the entire gastrointestinal tract (GIT) was removed and placed in a petri dish with normal saline. The GIT was then divided into sections: stomach, small intestine and large intestine in separate petri dishes. Helminths were collected using a pipette, killed in warm alcohol and preserved in 70 % alcohol. The parasites were cleared in lactophenol, examined under the compound microscope and identified using standard . The length of the parasites was measured to the nearest mm. The number of parasites from each fish was counted and preserved in 70 % alcohol. The mean intensity and parasite prevalence was calculated from:

$$\text{Mean Intensity of parasite} = \frac{\text{Number of individual parasites}}{\text{Infected Individual host species}}$$

$$\text{Percentage (\%) parasite prevalence} = \frac{\text{Number of infected fish}}{\text{Total number of all fish}} \times 100$$

The relative condition (Kn) of fish was calculated from the equation

$$Kn = W / a * L^b = \text{Observed weight} / \text{Expected weight}$$

where

W = weight of fish (g)

L = Length of fish (cm)

a = a constant

b = Growth factor

The statistical test for parasite prevalence, intensity and Kn was analyzed using the student's T- test of JMP statistical software.

107 **RESULTS**

108 The descriptive statistics of length and weight of fish samples are presented in Table 1. The
 109 range of the standard length of *B. africanus* was 3 - 12 cm, and 5.9 - 15 cm for *P. papilio*
 110 samples. The *P. papilio* samples were generally bigger than the *B. africanus* samples col-
 111 lected.

112 **Table 1 Descriptive Statistics of Measured Morphometric Parameters of *B. africanus* and**
 113 ***P. papilio* from Station 1 and Station 2**

Descrip- tive Sta- tistics	<i>B. africanus</i>		<i>P. papilio</i>	
	Rumuolumeni	Bakana	Rumuolu meni	Bak ana
Mean standard length (cm) ± S.D	7.9 2 ± 1.2 8	6 . 9 7 ± 0 . 9 3	8.48 ± 1.54	
Range of standard length (cm)	5 – 12	3 - 8 . 5	5.9 - 15	
Mean weight (g) ± S.D	11. 07 ± 5.9 1	7 . 6 6 ± 2 . 3 8	9.53 ± 5.15	
Range of Weight (g)	3 - 36. 1	3 . 2 - 1 3 . 7	4 - 28.2	

114

The Kn (Relative condition factor) of *P. papilio* from Bakana was between 0.98 - 1.05 (mean of 1.0) and between 1.20 - 8.37 (mean of 3.5) for Rumuolumeni. *B. africanus* samples from both had a Kn of 1.0. The length - weight relationship gave b values of < 3 for both species at both sample stations (Table 2).

Table 2 : The Length-Weight Relationship (b) and Relative Condition Factor (Kn) of *B. africanus* and *P. papilio*.

Species-Station	Length-Weight Relationship (b)	Mean Condition Factor $\pm$ S.D
<i>B.africanus</i> Rumuolumeni	2.94	1.0 $\pm$ 0.08
<i>B.africanus</i> – Bakana	1.51	1.0 $\pm$ 0.16
<i>P. papilio</i> – Bakana	2.95	1.0 $\pm$ 0.01
<i>P.papilio</i> Rumuolumeni	2.52	3.5 $\pm$ 0.08

The Kn of infected against non-infected fish were not different at $P \leq 0.05$ for both *B. africanus* and *P. Papilio*. The length of infected *B. africanus* was different from non-infected at $P \leq 0.05$. The length of infected *P. papilio* against non-infected were not significantly different at $P \leq 0.05$.

Identification of Parasites


134 The parasites from the GIT of both fishes were ascarid nematodes presented in Plates 1 - 3. The
 135 male and female worms could be distinguished externally and internally by size and the presence
 136 or absence of some bodily structures, while internally they had different reproductive organs
 137 Plates (2 and 3). The female worms were wider and grew up to 2 – 4 cm long, while males were
 138 thinner and grew up to 1.5 – 3 cm long. Examination of posterior opening showed that male
 139 ascarid had pineal spicules or spine – like extensions near its opening. It also had papillae or bump
 140 – like protrusions, in front and behind its posterior opening (Plate 3) while the females lacked the-
 141 se structures (Plate 2). The female worm had a reproductive opening on the posterior third of its
 142 body, while the males lacked such opening. Examining the posterior region of the body cavity un-
 143 der a light microscope, a tube – shaped reproductive organ was seen. The females had two tubes
 144 that joined together to form a “Y”, while the males had one straight tube. The plates show that
 145 there are 3 different types of the nematode but due to difficulties in identifying to species were
 146 pooled as ascarids.

147

148


149

150



151

152



153

154

Plates 1 A– E: Five variations in the anterior regions of ascarid from gobies

155

Scale: 0.25mm

156




157

158

Plates 2 A – C: Three variations in the posterior regions of female ascarid from gobies. Scale: 0.25mm

159



160

161 Plates 3 A – C: Three variations in the posterior regions of male ascarid from gobies. Scale: 0.25mm

162

163 **Percentage (%) Occurrence of *Ascaris* in Fish Species**

164 In Bakana, the percentage of occurrence of the ascarids in *B. africanus* and *P. papilio* was
 165 91.6% and 0.7% respectively, while in Rumuolumeni, the percentage occurrence in *B.*
 166 *africanus* and *P. papilio* were 21.8% and 5.7% respectively (Table 3). *B. africanus* had the
 167 highest percentage occurrence from both stations. The mean intensity of the worms in *B.*
 168 *africanus* and *P. papilio* were 23.0 and 4.0 respectively in Bakana, and 20.5 and 8.0 respec-
 169 tively in Rumuolumeni. The mean intensity was higher in *B. africanus* at both stations. In
 170 both gobies, these ascarid nematodes were found in their large intestine, small intestine
 171 and stomach. The cummulative parasite burden for both fish species was 25.4 %.

172 **Table 3: Incidence of *Ascaris* in *B. africanus* and *P. papilio***

STATION	SPECIES	% (Percentage) Occurrence of Ascarid	Mean In- tensity of ascarids	Location of Ascarids in Fish Host
Rumuolumeni	<i>B. africanus</i>	91.6%	23.0	Small intestine, large intes- tine, Stomach.
	<i>P. papilio</i>	0.7%	4.0	Small intestine, Stomach
Bakana	<i>B. africanus</i>	21.8%	20.5	Small intestine, Stomach.
	<i>P. papilio</i>	5.7%	8.0	Small intestine, Large In- testine

173

174

175

176

177

178 **Incidence of Parasite in *B. africanus* and *P. papilio* at Rumuolumeni and Bakana Rivers in Rivers**
 179 **State.**

180

181 In Rumuolumeni river, the percentage occurrence of helminth parasite in *B.africanus* and *P.*
 182 *papilio* were 91.6% and 0.7% respectively. While in Bakana river, the percentage occurrence of
 183 helminth parasite in *B.africanus* and *P. papilio* were 21.8% and 5.7% respectively, as shown in ta-
 184 ble 4. In Rumuolumeni river, the parasite mean intensity in *B.africanus* and *P. papilio* were 23.0
 185 and 4.0 respectively. In Bakana river, the mean intensity of parasite infection in *B.africanus* and *P.*

186 *papilio* were 20.5 and 8.0 respectively, as shown in Table 4. The *B.africanus* species collected from
 187 Rumuolumeni river had some helminth parasites located in their small intestine, large intestine
 188 and stomach. *P. papilio* species had some of the parasite located in the small intestine and stom-
 189 ach.

190 Table 4: **Incidence of Parasite in *B.africanus* and *P. papilio* from Rumuolumeni and Bakana Riv-**
 191 **ers.**

STATION	SPECIES	% (Percentage) Occurrence of Parasite	Parasite of Mean Intensi-ty	Location of Parasite in Fish Host
Rumuolumeni	<i>B.africanus</i>	91.6%	23.0	Small intestine, large intestine, Stomach.
	<i>P. papilio</i>	0.7%	4.0	Small intestine, Stomach
Bakana	<i>B.africanus</i>	21.8%	20.5	Small intestine, Stomach.
	<i>P. papilio</i>	5.7%	8.0	Small intestine, Large Intestine

192

193

194 The students T- test analysis of parasite prevalence of *B. africanus* from Bakana and Rumuolumeni
 195 revealed significant difference @ P = 0.05 Fig 2, while the parasite prevalence of *P. papilio* from
 196 Bakana and Rumuolumeni were not significantly different @ P = 0.05 (Fig 3). The

197

198




Fig. 2: Student's T- Test analysis of parasite prevalence of *B. africanus* from Bakana and Rumuolumeni




Fig 3: Student's T- Test analysis of parasite prevalence of *P. papilio* from Bakana and Rumuolumeni

DISCUSSION

Approximately 25.4 % of 240 fish investigated from this study were infested with ascarids indicating a low level of parasite burden in the fish population from both sampling stations. This does not conform to the report by [14] on the parasite load of *Ascaris* in *Battygobius soporator* from Lagos lagoon which had a high level (47 %) of parasite burden in the fishes investigated. The parasite burden of the individual species however revealed a higher burden on *B. africanus*. The presence of only one family of nematode from this study, on both species of gobies from both sampling stations indicated low parasite species diversity

which is not in agreement with the report by [15] on the Niger Delta tidal creek in Nigeria where they recorded no infestation in the gobies species collected from Buguma creek. The parasite *Ascaris* and *Anisakis* has been recorded from Atlantic cod, *Gadus morhua* which was sampled in the Barents, Baltic and North Seas [16; 17]. [18] and [19] recorded *Camallanus* and *Procamallanus* species from another Nigerian creek. This difference might be because parasites are focal and their prevalence are determined by prevailing environmental parameters. In fishes, host and pathogen interaction is modulated by environmental factors and this is a key determinant of susceptibility to disease [20]. Ascaroid nematodes are important helminths which have the potential of being zoonotic as was reported by [21] during a study, when they isolated the eggs of some helminth parasites (nematodes; *Capillaria sp.*, *Ascaris sp.*, *Enterobius sp.*, and hookworm) which had zoonotic potency, from sediments and samples of Carp in India. The presence of ascarid nematode in the sampled fish species is also in conformity with the study by [22] who identified Ascaridoid nematode larvae (*Anisakis*, *Hysterothylacium*, *Raphidascaris* and *Terranova*) from the intestinal lumen and abdominal organs of marine fishes from New Caledonia. Their research reported those nematodes as being important and potential zoonotic parasites in that region. The ratio of female to male Ascarid nematodes isolated were 2:1 and the presence of matured females (gravid) and male nematode parasite in the fish samples indicates that the species of Gobies investigated were the definitive host of the parasites.

From this study, *B. africanus* species from Rumuolumeni were larger than those in Bakana and recorded more parasite intensity and prevalence (91.6 %). This result is in agreement with the study of [14] who recorded higher parasite prevalence (61.5 %) in *B. soporator* in bigger fish of 15 – 17 cm than the smaller 9 – 11 cm fish (19.4 %). This would mean that older fish may have higher parasite prevalence probably due to longer exposure to the parasite. The record of *P. papilio* samples collected from both sampling stations does not agree with the findings of [14] as it appears that size was not a determining factor on parasite prevalence. There was also no relationship between parasite load (intensity) and fish length and weight, which is in agreement with the findings of [23] on the helminth parasites of *Tilapia zilli* from River Oshun. The intensity should depend on prevailing environmental conditions that make the host susceptible to the parasite.

The gobies (*B. africanus* and *P. papilio*) from this study inhabit the intertidal regions of the creeks and feed on crustaceans and planktons. Children are often seen on the tidal flats

picking periwrinkles (*Tympanotonus*) and mudskippers (*P. papilio*). The possibility of infection when they ingest the eggs of these parasites can not be ruled out. [14] has suggested that the presence of *Ascaris* species in *B. saporator* was because of fecal waste discharged into the Lagos lagoon. The significantly higher prevalence of ascarids in *B. africanus* during the wet season does not agree with previous work by [15], who recorded the highest dominance of parasites in *Polydactylus quadrifilis* during the dry season. This shows some variability in parasite prevalence for different seasons.

Many factors such as the change in physico-chemical properties of the water body, fish species, food, sex and stage of maturity [24] can be responsible for the change of b values for the length – weight relationships of gobies. The length- weight relationship analysis of both species of gobies from both sampling stations, indicated negative allometric growth ($b < 3$). This implies more increase in length as growth proceeds, giving a slimmer fish as the increase in length does not give a corresponding increase in weight. These findings are similar with the findings of [25] and [26] that observed negative allometric growth for different species of cichlid from the Anambra River.

The condition factor of a fish depends on a number of factors, availability of food, ideal temperature and salinity for growth. The *P. papilio* from Rumuolumeni appeared to be in a better condition with a Kn of 3.5 while *B. africanus* was 1. [25] also recorded mean condition factors from cichlids which showed variations due to seasonal and environmental changes. Fishes thrive better in water bodies that have less anthropogenic activities and are less polluted. The condition factor of fish describes the physiological state of the fish with respect to its welfare and nutritional status[27]. *P. papilio* from Rumuolumeni with a mean relative condition factor of 3.5 with low parasite prevalence were in better condition. This agrees with the records from the study by [28] on the cat fish, *Synodontis filamentosus* and *Calamoichthys calabariscus* sampled from Lekki lagoon, which were in good condition due to the absence of parasites. The condition factor of infected and non – infected fish was not significantly different implying that the infection with these parasites did not affect the length – weight relationship of the fish. However, it is possible that there could be pathological effects on the gastrointestinal tract (GIT) of the fish which may not result in weight loss. When nematodes are numerous, they can block the lumen of the GIT and affect digestion and absorption of food.

In conclusion, *B. africanus* and *P. papilio* from Rumuolumeni and Bakana creeks in the Niger Delta were infected with Ascarid nematodes in the GIT. The prevalence and intensity of infection was higher in *B. africanus* and at Rumuolumeni. Size appeared to be a factor in determining percentage prevalence for *B. africanus*. The mean Kn was greater than 1 for both species but *P. papilio* with lower parasite prevalence was in better condition.

REFERENCES

1. Kaur P. Histopathological Effect of *Senga* Species (Cestode: Pseudophyllidae) in the Intestine of *Piscia* Host. *World Journal of Pharmacy and Pharmaceutical Sciences*. (2014). 3(2): 1506 – 1513.
2. Hilderband K, Price R, and Olson R. Parasites in Marine Fishes, Questions And Answers for Seafood Retailers, Oregon State University Extension Service, Oregon, USA: JG Publication. (1985).
3. Lagrue C, Kelly D, Hicks A and Poulin R. Factors Influencing Infection Patterns of Tightly Transmitted Parasites among a Fish Community: Host Diet, Host – Parasite Compatibility or Both. *Journal of Fish Biology*, (2011). 79 (4): 466- 485.
4. Khalil L and Polling L. Checklist of Helminth Parasite of African Freshwater Fishes. University of North Republic of South Africa. (1997). 161 pp.
5. Patzner R, Van J and Kapoor B. Biology of Gobies. United Kingdom: CRC Press (2011).
6. Hernaman V, Probert P and Robbins W. Trophic Ecology of Coral Reef Gobies: Interspecific, Ontogenetic and Seasonal Comparison of Diet and Feeding Intensity. *Marine Biology*, (2009). 156 (4): 317-330.
7. Akinsanya B, Otubanjo A and Ibidapo C. Helminth Biload of *Chrysichthys Nigrodigitatus* from Lekki Lagoon, Lagos, Nigeria, *Turkish Journal of Fisheries and Aquatic Sciences*, (2007). 4(7): 83 – 87.
8. Worgu M and Okaka E. A comparative of Study Gastrointestinal Helminth Parasites Infection of Fresh and Brackish Water Fishes from Warri River, Southern Nigeria. *African Research Review*, (2012). 6(2): 13 – 23.

- 302 9. Oribhabor J, Ogbeibu E and Okaka E. The Parasites Gastrointestinal Helminth of some
 303 *Scianid* Species (Croakers) in a Niger Delta Mangrove Creek, Nigeria. *Tropical*
 304 *Freshwater Biology*, (2010), 91 (7):15 – 23.

- 305 10. Oribhabor J, Ogbeibu E and Okaka E. The gastrointestinal Helminth Parasites of the
 306 Threadfin Fish, *Polydactylus quadrifilis* (Family: Polynemidae) in a Niger Delta man-
 307 grove creek, Nigeria. *International Journal of Animal and Veterinary Advances*,
 308 (2012), 4: 240 -243.

- 309 11. Powell C. Effect of Fresh Water Oil Spillage on Fish and Fisheries. In: Proceedings of In-
 310 ternational Seminar on the Petroleum Industry and Nigerian Environment, Rivers
 311 State, Nigeria, July, 1987, (1987)

- 312 12. Dienye H and Woke G. Physicochemical parameters of the upper and lower reach of
 313 the New Calabar Rivers, Niger Delta. *J Fisheries Livest. Prod.* (2015) 3: 154.

- 314 13. Dublin -Green C. Seasonal Variation in some Physico-chemical Parameters of the Bonny
 315 Estuary, Niger Delta. *NIOME Technical*, (1990). 59(3): 21 – 25.

- 316 14. Bamidele J, Amaeze N, and Adeoye G. Prevalence and Intensity of *Ascaris*
 317 Infection and Heavy Metal Accumulation in *Battygobius soporator* (Frilfin gobby) of
 318 the Lagos Lagoon, Nigeria. *The Zoologist*, (2014). 12(3): 8 – 15.

- 319 15. Ogbeibu A, Okaka C and Oribhabor B. Gastrointestinal Helminth Parasites Community
 320 of Fish Species in a Niger Delta Tidal Creek, Nigeria. *Journal of Ecosystems*, (2013).
 321 20(5): 1 - 10.

- 322 16. Gay M, Bao M, Mackenzie K, Pascual S, Buchmann K, Bourgau O, Couvreur C, Matiucci
 323 S, Paoletti M, Hastie LC, Levse A and Pierce GJ. Infection levels and species diversity
 324 of ascaridoid nematodes in Atlantic cod, *Gadus morhua*, are correlated with geo-
 325 graphic are and fish size. *Fisheries Research*, (2018), 202: 90 - 102.

- 326 17. Pierce GJ, Bao M, Mackenzie K, Dunser A, Giuletti L, Cipriani P, Matiucci S, Hastie LC.
 327 Ascaridoid nematode infection in haddock (*Melanogrammus aeglefinus*) and whit-
 328 ing (*Merlangius merlangus*) in Northeastern Atlantic waters. *Fisheries Research*,
 329 (2018), 202: 122 - 133.

- 330 18. Okaka E and Akhigbe E. Helminth Parasites of Some Tropical Freshwater Fish from Osse
331 River in Benin, Southern Nigeria. *Tropical Freshwater Biology*, (1999). 8(1): 41 – 48.
- 332 19. Okaka E. A Survey into the Helminth Parasites of Fishes of Asa River and the impound-
333 ment at Asa Dam, Ilorin, Nigeria, *Rivista di parassitologia*, (2010). 22(5): 207 - 210.
- 334 20. Arkoosh M, Casillas E, Clemons E and Stein J. Effect of Pollution on Fish Disease: Po-
335 tential Impacts on Salmonid Populations. *Journal of Aquatic Animal Health*, (1998).
336 10(3): 182 - 190.
- 337 21. Soham P and Gadahar D. Isolation and Identification of Helminth Parasite Eggs Having
338 Zoonotic Potency from Sewage-Fed Pond Sediment, West Bengal, India. *Interna-
339 tional Journal of Advance Research and Biological Science*, (2016). 3(1): 283 - 291.
- 340 22. Shokoofet S, Poupa A and Jean-Lou J. Characterisation of Ascaridoid Larval from Ma-
341 rine Fish Off New Caledonia with Description of New Hysterothylacium larval types
342 XIII and XIV. *Parasitology International*, (2015). 64(4): 399 - 404.
- 343 23. Kayode O, Okafor J, Alade A and Oronaye O. Helminth Parasites of *Sarotherodon*
344 *galilaeus* and *Tilapia zilli*, (Pisces; Cichlidae) from River Oshun, South West Nigeria.
345 *International Journal of Aquatic Science*, (2012). 3(3): 49 - 55.
- 346 24. Obasohan E, Imasuen J and Isidahome C. Preliminary Studies of the Length – Weight
347 Relationship and Condition Factor of Five Fish Species from Ibiekuma Stream,
348 Ekpoma, Edo state, Nigeria. *Journal of Agricultural Research and Development*,
349 (2012). 2(6): 61 – 69.
- 350 25. Atama C, Okeke O, Ekeh F and Ezenwaji N. Length – Weight Relationship and Condition
351 Factor of Six Cichlid Species of Anambra River. *Journal of Fisheries and Aquaculture*,
352 (2013). 4(5): 82 - 86.
- 353 26. Soyinka OO, Ayoola SO, Onyema IC and Ayeni MO. Aspects of the Biology and Culture
354 Trials of Cichlids from Lagos Lagoon. *J. Sci. Resource Dev.* (2013). 14: 47-54.
- 355 27. Le Cren E. The Length-Weight Relationship and Seasonal Cycle in Gonad Weight and
356 Condition in Perch, *Perca fluviatilis*. *Journal of Animal Ecology*, (1951). 20 (6): 201 –
357 219.

- 358 28. Akinsanya B. A Two Fish Species Study of the Parasitic Helminth Fauna of *Synodontis*
359 *filamentus* and *Calamoichthys calabaricus* from Lekki Lagoon, Lagos, Nigeria. *Ife*
360 *Journal of Science*, (2015). 17(1): 97 – 111.

361

362

363

364