

Charge Exchange of Proton-Potassium Atom Collision

ABSTRACT

The coupled static approximation is modified for the first time to make it applicable to multi-channels problem of the collision of the proton by alkali atom. The possibility of producing more hydrogen during the proton-alkali atom collision is investigated. The formation of hydrogen H(1s) and excited hydrogen (in 2s- and 2p-states) of p-K collision is treated to test the convergence of our method. The modified method is used to calculate the total cross-sections of seven partial waves($0 \leq \ell \leq 6$, where ℓ is the total angular momentum) at a range of energy between 50 and 1000 keV. Our p-K results are compared with previous ones.

Keywords: proton-alkali, proton-potassium, hydrogen formation, excited hydrogen formation, cross-sections.

PACS Nos.: 34.80.Dp . 34.80.Gs . 34.80.Ht

1.INTRODUCTION

The most interesting phenomenon in quantum mechanics is the intermediate states that appear in a nuclear reaction . Most theoretical and experimental studies of proton-atom interactions are discussed in the last decade by calculating differential and total cross-sections as functions of incident energies. Choudhury and Sural [1] have studied p-alkali atom (Na, K, Rb, Cs) scattering in the wave formation of impulse approximation at a range of energy from 50 to 500 keV. . Daniele et al. [2] have been reported the total cross-sections for high energy proton scattering by alkali atom using eikonl - approximations. Ferrante and Fiordilino [3] have been discussed the

eikonal-approximation to study high-energy proton collision with alkali atom. Ferrante et al. [4] have also studied the total H-formation cross-sections in p-alkali atom scattering using OBK approximation. Tiwari [5] have been investigated the differential and total cross-sections of H-formation of the collision of p-Li and p-Na atom using the Coulomb-projected Born approximation.

The present work is to explore the possibility of producing more hydrogen during the proton-potassium atom collision. For this reason, it is important to discuss the scattering of p-K atom . In the present paper, the CSA method used by Elkilany ([6]-[8]) will be modified to make it applicable to discuss the MCSA problem ($n=4$) of the collision of p-K atom at intermediate energies of the projectile. A numerical procedure will be generalized to solve the obtained multi-coupled equations. Throughout this paper Rydberg units have been used and the total cross-sections are expressed in units of πa_0^2 ($=8.8 \times 10^{-17} \text{ cm}^2$) and energy units of keV.

2. THEORETICAL FORMALISM

The MCSA of protons scattered by alkali atoms can be written by (see Fig. 1)

$$p + A = \begin{cases} p + A & \text{Elastic channel (first channel)} \\ H(n\ell) + A^+ & H(n\ell) \text{ formation channels } ((n-1) - \text{channels}) \end{cases} \quad (1)$$

where p is the proton, A is an alkali target atom, $H(n\ell)$ is hydrogen formation of $n\ell$ -states and n is the number of open channels.




Fig. 1 Configuration space of p-atom collision: $\bar{x}_i$ and $\bar{r}_i$ are the vectors of the proton and the valence electron of the target with respect to the center of mass of the target, $\bar{\rho}_i$ is the vector of the proton with respect to the valence electron of the target, $\bar{\sigma}_i$ is the vector of the center of mass of H from the target, M_T = mass of the nucleus of the target.

The Hamiltonian of the elastic channel is given by:

$$H = H^{(1)} = H_T - \frac{1}{2\mu_1} \nabla_{x_1}^2 + V_{\text{int}}^{(1)}(x_1) = -\frac{1}{2\mu_T} \nabla_{r_1}^2 - \frac{2}{r_1} + V_c(r_1) - \frac{1}{2\mu_1} \nabla_{x_1}^2 + V_{\text{int}}^{(1)}(x_1), \quad (2)$$

where H_T is the Hamiltonian of the target atom. μ_T is the reduced mass of the target atom.

The Hamiltonian of the (n-1)-rearrangement channels are expressed by:

$$H = H^{(i)} = H_i - \frac{1}{2\mu_i} \nabla_{\sigma_i}^2 + V_{\text{int}}^{(i)}(\sigma_i) = -\frac{1}{2\mu_i} \nabla_{\rho_i}^2 - \frac{2}{\rho_i} - \frac{1}{2\mu_i} \nabla_{\sigma_i}^2 + V_{\text{int}}^{(i)}(\sigma_i), \quad i = 2, 3, 4, \dots, n$$

(3)

66 where H_i , $i = 2, 3, 4, \dots, n$ are the Hamiltonians of the hydrogen formation atoms, $H(n\ell)$,
 67 respectively. μ_i , $i = 2, 3, 4, \dots, n$ are the reduced masses of (n-1)- channels, respectively.

68 $V_c(r_1)$ is a screened potential and $V_{\text{int}}^{(1)}(x_1)$ is the interaction potential of the first channel and
 69 are given by:

$$70 \quad V_c(r_1) = V_{c\text{Coul}}(r_1) + V_{cex}(r_1)$$

71 (4)

72 where $V_{c\text{Coul}}(r_1)$ and $V_{cex}(r_1)$ are the Coulomb and exchange parts of the core potential,
 73 respectively(see ref. [8]), and

$$74 \quad V_{\text{int}}^{(1)}(x_1) = \frac{2}{x_1} - \frac{2}{\rho_1} + V_{c\text{Coul}}(x_1) \quad \text{where} \quad V_{c\text{Coul}}(x_1) = -V_{c\text{Coul}}(r_1) \quad (5)$$

75 and $V_{\text{int}}^{(i)}(\sigma_i)$, $i = 2, 3, 4, \dots, n$, are the interaction potentials of the (n-1)-hydrogen formation
 76 channels, respectively and are given by:

$$77 \quad V_{\text{int}}^{(i)}(\sigma_i) = \frac{2}{x_i} - \frac{2}{r_i} + V_{c\text{Coul}}(x_i) + V_{c\text{Coul}}(r_i) + V_{cex}(r_i), \quad i = 2, 3, 4, \dots, n \quad (6)$$

78 The total energies E of the n-channels are defined by:

$$79 \quad E = E_i + \frac{1}{2\mu_i} k_i^2, \quad i = 1, 2, 3, \dots, n \quad (7)$$

80 where $\frac{1}{2\mu_1} k_1^2$ is the kinetic energy of the incident proton relative to the target and $\frac{1}{2\mu_i} k_i^2$
 81 , $i = 2, 3, 4, \dots, n$ are the kinetic energy of the center-of-mass of the hydrogen formation atoms,
 82 $H(n\ell)$, respectively, with respect to the nucleus of the target. E_1 is the binding energy of the

83 target atom, and E_i , $i=2,3,4,\dots,n$ refer to the binding energies of the hydrogen formation
84 atoms, respectively.

85 In the multi-channels coupled-static approximation (MCSA), it is assumed that the
86 projections of the vector $(H-E)|\Psi\rangle$ onto the bound state of the n-channels are zero. Thus, the
87 following conditions:

$$88 \quad \langle \Phi_i | (H-E) | \Psi \rangle = 0, \quad i = 1, 2, 3, \dots, n \quad (8)$$

89 are satisfied. The total wavefunction $|\Psi\rangle$ is expressed by:

$$90 \quad \Psi = \sum_{i=1}^n |\Phi_i \psi_i\rangle, \quad (9)$$

$$91 \quad \psi_1 = \sum_{\ell} \ell(\ell+1) f_{\ell}^{(1)}(x_1) Y_{\ell}^0(\hat{x}_1), \quad (10)$$

$$92 \quad \psi_i = \sum_{\ell} \ell(\ell+1) g_{\ell}^{(i)}(\sigma_i) Y_{\ell}^0(\hat{\sigma}_i), \quad i = 2, 3, \dots, n \quad (11)$$

93 where $f_{\ell}^{(1)}(x_1)$ and $g_{\ell}^{(i)}(\sigma_i)$, $i = 2, 3, \dots, n$ are the radial wavefunctions of the elastic and the
94 hydrogen formation atoms, respectively, corresponding to the total angular momentum ℓ .

95 $Y_{\ell}^0(x_1)$ and $Y_{\ell}^0(\hat{\sigma}_i)$ $i = 2, 3, \dots, n$ are the related spherical harmonics.

96 $\hat{x}_1$ and $\hat{\sigma}_i$, $i = 1, 2, 3, \dots, n$ are the solid angles between the vectors $\hat{x}_1, \hat{\sigma}_i, i = 2, 3, \dots, n$ and the z-
97 axis, respectively. ψ_i , $i = 1, 2, 3, \dots, n$ are the corresponding scattering wavefunction of the n-

98 channels, respectively. Φ_1 is the wavefunction for the valence electron of the target atom which
99 is calculated using ref. [9]. Φ_i , $i = 2, 3, 4, \dots, n$ are the wavefunctions of the hydrogen

100 formation atoms, $H(n\ell)$, respectively, which are defined using hydrogen like wavefunction.

101 The multi-channels coupled static approximation (MCSA) (eq. (8)) can be solved by considering
 102 the n- integro-differential equations

$$103 \quad \left[\frac{d^2}{dx_1^2} - \frac{\ell(\ell+1)}{x_1^2} + k_1^2 \right] f_\ell^{(1)}(x_1) = 2\mu_1 U_{st}^{(1)}(x_1) f_\ell^{(1)}(x_1) + \sum_{\alpha=2}^n Q_{1\alpha}(x_1), \quad (12)$$

$$104 \quad \left[\frac{d^2}{d\sigma_i^2} - \frac{\ell(\ell+1)}{\sigma_i^2} + k_i^2 \right] g_\ell^{(i)}(\sigma_i) = 2\mu_i U_{st}^{(i)}(\sigma_i) g_\ell^{(i)}(\sigma_i) + \sum'_{\alpha=1}^n Q_{i\alpha}(\sigma_i), \quad i=2,3,\dots,n, \quad (13)$$

105 where the prime on the summation sign means that $i \neq \alpha$, and

$$106 \quad Q_{1\alpha}(x_1) = \int_0^\infty K_{1\alpha}(x_1, \sigma_\alpha) g_\ell^{(\alpha)}(\sigma_\alpha) d\sigma_\alpha, \quad \alpha=2,3,\dots,n \quad (14)$$

$$107 \quad Q_{i1}(\sigma_i) = \int_0^\infty K_{i1}(\sigma_i, x_1) f_\ell^{(1)}(x_1) dx_1, \quad i=2,3,\dots,n \quad (15)$$

$$108 \quad Q_{i\alpha}(\sigma_i) = \int_0^\infty K_{i\alpha}(\sigma_i, \sigma_\alpha) g_\ell^{(\alpha)}(\sigma_\alpha) d\sigma_\alpha, \quad i, \alpha=2,3,\dots,n, \quad i \neq \alpha \quad (16)$$

109 the Kernels $K_{i\alpha}$, $i=1,2,3,\dots,n, i \neq \alpha$ are expanded by:

$$110 \quad K_{1\alpha}(x_1, \sigma_\alpha) = 2\mu_1 (8x_1 \sigma_\alpha) \left[\Phi_1(r_1) \Phi_\alpha(\rho_\alpha) \left[-\frac{1}{2\mu_\alpha} (\nabla_{\sigma_\alpha}^2 + k_\alpha^2) + V_{\text{int}}^{(\alpha)} \right] Y_\ell^o(\hat{x}_1) Y_\ell^o(\hat{\sigma}_\alpha) d\hat{x}_1 d\hat{\sigma}_\alpha, \quad \alpha=2,3,\dots,n, \right. \\ 111 \quad \left. (17) \right]$$

$$112 \quad K_{i1}(\sigma_i, x_1) = 2\mu_i (8\sigma_i x_1) \left[\Phi_i(\rho_i) \Phi_1(r_1) \left[-\frac{1}{2\mu_1} (\nabla_{x_1}^2 + k_1^2) + V_{\text{int}}^{(1)} \right] Y_\ell^o(\hat{\sigma}_i) Y_\ell^o(\hat{x}_1) d\hat{\sigma}_i d\hat{x}_1, \quad i=2,3,\dots,n, \right. \\ 113 \quad \left. (18) \right]$$

$$114 \quad K_{i\alpha}(\sigma_i, \sigma_\alpha) = 2\mu_i (8\sigma_i \sigma_\alpha) \left[\Phi_i(\rho_i) \Phi_\alpha(\rho_\alpha) \left[-\frac{1}{2\mu_\alpha} (\nabla_{\sigma_\alpha}^2 + k_\alpha^2) + V_{\text{int}}^{(\alpha)} \right] Y_\ell^o(\hat{\sigma}_i) Y_\ell^o(\hat{\sigma}_\alpha) d\hat{\sigma}_i d\hat{\sigma}_\alpha, \quad i, \alpha=2,3,\dots,n, \quad i \neq \alpha. \right. \\ 115 \quad \left. (19) \right]$$

116 The static potentials $U_{st}^{(1)}(x_1)$ and $U_{st}^{(i)}(\sigma_i)$, $i=2,3,\dots,n$ are defined by

$$U_{st}^{(1)}(x_1) = \langle \Phi_1(r_1) | V_{\text{int}}^{(1)} | \Phi_1(r_1) \rangle, \quad U_{st}^{(i)}(\sigma_i) = \langle \Phi_i(\rho_i) | V_{\text{int}}^{(i)} | \Phi_i(\rho_i) \rangle \quad (20)$$

The equations (12,13) are inhomogeneous equations in x_i , and σ_i , $i=1,2,3,\dots,n$ and are possessing the general form

$$(\mathcal{E} - H_0) |\chi\rangle = |\eta\rangle \quad (21)$$

where $\mathcal{E}$ is k_i^2 ($i=1,2,\dots,n$). H_0 is $-\frac{d^2}{dx_1^2} + \frac{\ell(\ell+1)}{x_1^2}$ or $-\frac{d^2}{d\sigma_i^2} + \frac{\ell(\ell+1)}{\sigma_i^2}$, $i=2,3,\dots,n$. $|\chi\rangle$

is $|f_\ell^{(1)}(x_1)\rangle$ or $|g_\ell^{(i)}(\sigma_i)\rangle$, $i=2,3,\dots,n$. $|\eta\rangle$ is the right-hand side of the equations, respectively.

The solution of eqs. (12,13) are given (formally) by Lippmann-Schwinger equation in the form

$$|\chi\rangle = |\chi_0\rangle + G_0 |\eta\rangle \quad (22)$$

where G_0 is Green operator $(\mathcal{E} - H_0)^{-1}$ and $|\chi_0\rangle$ is the solution of the homogeneous equation

$$(\mathcal{E} - H_0) |\chi_0\rangle = |0\rangle, \quad (23)$$

Using Green operator G_0 , the solutions of (12,13) are given formally by

$$f_\ell^{(1,j)}(x_1) = \{\delta_{j1} + \frac{1}{k_1} \int_0^\infty \tilde{g}_\ell(k_1 x_1) [2\mu_1 U_{st}^{(1)}(x_1) f_\ell^{(1,j)}(x_1) + \sum_{\alpha=2}^n Q_{1\alpha}^{(j)}(x_1)] dx_1\} \tilde{f}_\ell(k_1 x_1) \\ + \{-\frac{1}{k_1} \int_0^\infty \tilde{f}_\ell(k_1 x_1) [2\mu_1 U_{st}^{(1)}(x_1) f_\ell^{(1,j)}(x_1) + \sum_{\alpha=2}^n Q_{1\alpha}^{(j)}(x_1)] dx_1\} \tilde{g}_\ell(k_1 x_1), \quad j=1,2,3,\dots,n \quad (24)$$

$$g_\ell^{(i,j)}(\sigma_i) = \{\delta_{ji} + \frac{1}{k_i} \int_0^\infty \tilde{g}_\ell(k_i \sigma_i) [2\mu_i U_{st}^{(i)}(\sigma_i) g_\ell^{(i,j)}(\sigma_i) + \sum_{\alpha=1}^n Q_{i\alpha}^{(j)}(\sigma_i)] d\sigma_i\} \tilde{f}_\ell(k_i \sigma_i) \\ + \{-\frac{1}{k_i} \int_0^\infty \tilde{f}_\ell(k_i \sigma_i) [2\mu_i U_{st}^{(i)}(\sigma_i) g_\ell^{(i,j)}(\sigma_i) + \sum_{\alpha=1}^n Q_{i\alpha}^{(j)}(\sigma_i)] d\sigma_i\} \tilde{g}_\ell(k_i \sigma_i), \quad i=2,3,\dots,n \quad j=1,2,3,\dots,n \quad (25)$$

133 where the delta functions δ_{ji} , $i, j = 1, 2, 3, \dots, n$, specify two independent forms of solutions for each
 134 of $f_\ell^{(1, j)}(x_1)$ and $g_\ell^{(i, j)}(\sigma_i)$, $i = 2, 3, \dots, n$. The functions $\tilde{f}_\ell(\eta)$ and $\tilde{g}_\ell(\eta)$,
 135 $\eta = k_1 x_1$, or $\eta = k_i \sigma_i$, $i = 2, 2, 3, \dots, n$ are related to the Bessel functions of the first and second
 136 kinds, i.e. $j_\ell(\eta)$ and $y_\ell(\eta)$, respectively, by the relations $\tilde{f}_\ell(\eta) = \eta j_\ell(\eta)$ and $\tilde{g}_\ell(\eta) = -\eta y_\ell(\eta)$.

137 The iterative solutions of Eqs.(24, 25) are calculated by:

$$138 \quad f_\ell^{(1, j, \nu)}(x_1) = \left\{ \delta_{j1} + \frac{1}{k_1} \int_0^{X_1} \tilde{g}_\ell(k_1 x_1) [2\mu_1 U_{st}^{(1)}(x_1)] f_\ell^{(1, j, \nu-1)}(x_1) + \sum_{\alpha=2}^n Q_{1\alpha}^{(j, \nu-1)}(x_1) dx_1 \right\} \tilde{f}_\ell(k_1 x_1) \\
 139 \quad + \left\{ -\frac{1}{k_1} \int_0^{X_1} \tilde{f}_\ell(k_1 x_1) [2\mu_1 U_{st}^{(1)}(x_1)] f_\ell^{(1, j, \nu-1)}(x_1) + \sum_{\alpha=2}^n Q_{1\alpha}^{(j, \nu-1)}(x_1) dx_1 \right\} \tilde{g}_\ell(k_1 x_1), \quad j = 1, 2, 3, \dots, n, \nu \geq 1. \\
 140 \quad (26)$$

$$141 \quad g_\ell^{(i, j, \nu)}(\sigma_i) = \left\{ \delta_{ji} + \frac{1}{k_i} \int_0^{\sum_i} \tilde{g}_\ell(k_i \sigma_i) [2\mu_i U_{st}^{(i)}(\sigma_i)] g_\ell^{(i, j, \nu)}(\sigma_i) + \sum_{\alpha=1}^n Q_{i\alpha}^{(j, \nu)}(\sigma_i) d\sigma_i \right\} \tilde{f}_\ell(k_i \sigma_i) \\
 142 \quad + \left\{ -\frac{1}{k_i} \int_0^{\sum_i} \tilde{f}_\ell(k_i \sigma_i) [2\mu_i U_{st}^{(i)}(\sigma_i)] g_\ell^{(i, j, \nu)}(\sigma_i) + \sum_{\alpha=1}^n Q_{i\alpha}^{(j, \nu)}(\sigma_i) d\sigma_i \right\} \tilde{g}_\ell(k_i \sigma_i), \quad i = 2, 3, \dots, n, j = 1, 2, 3, \dots, n, \nu \geq 0. \\
 143 \quad (27)$$

143 where $X_1, \sum_i$, $i = 2, \dots, n$ specify the integration range away from the nucleus over which the
 144 integrals at equations (26, 27) are calculated using Simpson's expansions.

145 Taylor expansion of $U_{st}^{(1)}(x_1)$, $\tilde{f}_\ell(k_1 x_1)$ and $\tilde{g}_\ell(k_1 x_1)$ are used to obtain starting value of

146 $f_\ell^{(1, j, 0)}(x_1)$ (see ref. [8]).

147 Equations (26, 27) can be abbreviated to

$$f_{\ell}^{(1,j,\nu)}(x_1) = a_1^{(j,\nu)} \tilde{f}_{\ell}(k_1 x_1) + b_1^{(j,\nu)} \tilde{g}_{\ell}(k_1 x_1), \quad j=1,2,3,\dots,n; \quad \nu > 0 \quad (28)$$

$$g_{\ell}^{(i,j,\nu)}(\sigma_i) = a_i^{(j,\nu)} \tilde{f}_{\ell}(k_i \sigma_i) + b_i^{(j,\nu)} \tilde{g}_{\ell}(k_i \sigma_i), \quad i=2,\dots,n, j=1,2,3,\dots,n; \quad \nu > 0 \quad (29)$$

The preceding coefficients of eqs (28,29) are elements of the matrices a^{ν} and b^{ν} which are given by:

$$\left. \begin{aligned} (a^{\nu})_{ij} &= \sqrt{2\mu m_i / k_i} a_i^{(j,\nu)} \\ (b^{\nu})_{ij} &= \sqrt{2\mu m_i / k_i} b_i^{(j,\nu)}, \quad i, j=1,2,\dots,n, \quad \nu > 0 \end{aligned} \right] \quad (30)$$

and we can obtain the reactance matrix, R^{ν} , using the relation:

$$\{R^{\nu}\}_{\beta\gamma} = \left\{ b^{\nu} (a^{\nu})^{-1} \right\}_{\beta\gamma}, \quad \beta, \gamma=1,2,3,\dots,n \quad \nu > 0. \quad (31)$$

The partial and total cross sections in the present work are determined (in πa_0^2) by :

$$\sigma_{\beta\gamma}^{(\ell,\nu)} = \frac{4(2\ell+1)}{k_1^2} |T_{\beta\gamma}^{\nu}|^2, \quad \beta, \gamma=1,2,3,\dots,n \quad \nu > 0 \quad (32)$$

where k_1 is the momentum of the incident protons, ν is the number of iterations and $T_{\beta\gamma}^{\nu}$ is the

elements of the $n \times n$ transition matrix T^{ν} which is given by:

$$T^{\nu} = R^{\nu} \left(I - \tilde{t} R^{\nu} \right)^{-1}, \quad \nu > 0, \quad (33)$$

where R^{ν} is the reactance matrix and I is a $n \times n$ unit matrix and $\tilde{t} = \sqrt{-1}$.

The total cross sections (in πa_0^2 units) can be obtained (in ν^{th} iteration) by:

$$\sigma_{ij}^{\nu} = \sum_{\ell=0}^{\infty} \sigma_{ij}^{(\ell,\nu)}, \quad i, j=1,2,3, \dots, n \quad \nu > 0 \quad (34)$$

3-PROTON POTASSIUM COLLISION

We are going to apply our MCSA in the case of $n=4$ (four channels CSA) to the scattering of p-K atoms. Our problem can be written in the form:

$$p + K(4s) = \begin{cases} p + K(4s) & \text{Elastic channel (first channel)} \\ H(1s) + K^+ & H(1s) \text{ formation channel (second channel)} \\ H(2s) + K^+ & H(2s) \text{ formation channel (third channel)} \\ H(2p) + K^+ & H(2p) \text{ formation channel (fourth channel)} \end{cases} \quad (35)$$

$\Phi_1(r_1)$ is the valence electron wavefunction of the target (potassium) atom which is calculated using (Clementi's tables [9]), and $\Phi_i(\rho_i)$, $i = 2,3,4$ are the wavefunctions of the hydrogen formation which are given by:

$$\Phi_2 = \frac{1}{\sqrt{\pi}} \exp(-\rho_2), \quad \Phi_3 = \frac{1}{\sqrt{32\pi}} (2 - \rho_3) \exp(-\rho_3/2) \quad \text{and} \quad \Phi_4 = \frac{1}{\sqrt{32\pi}} \rho_4 \cos\theta_{\rho_4, \sigma_4} \exp(-\rho_4/2) \quad (36)$$

4. RESULTS AND DISCUSSION

We start our calculations of p-K scattering by testing the variation of the static potentials $U_{st}^{(1)}(x_1)$ and $U_{st}^{(i)}(\sigma_i)$, $i = 2,3,4$ of the considered channels with the increase of x_1, σ_i ($i = 2,3,4$). It is found the excellent convergence of the calculated integrals can be obtained with Simpson's interval $h=0.0625$, and number of points 512 which give integration range $IR = 32a_0$ and with iterations, $\nu = 50$. We have calculated the total cross sections of p-K corresponding to $0 \leq \ell \leq 6$ at incident energies between 50 and 1000 keV.

Table 1 shows the present total cross-sections of p-K scattering with those of Choudhury and Sural [1], Daniele et al. [2], Ferrante et al. [4] in the energy range (50-1000 keV). Our results and those of comparing results in the range of energy (500-1000 keV.) are also displayed in Figs. (2-

4). In Fig.5 we also show the present results of the total cross-sections of the four channels (elastic and the hydrogen formation (H(1s), H(2s), H(2p)) in the range of energy (50-1000 keV.). The present values of the total cross-sections of the four channels have similar trends of the comparison results. Our values of the total cross-sections of the four channels decrease with the increase of the incident energies. The calculated total cross-sections σ_{12} of H(1s) are about (9.6%-11.8%) lower than the Choudhury and Sural results[1] and about (2.9%-13.2%) higher than those of Elkilany [8]. The total cross-sections σ_{13} of H(2s) are slightly about (7.4%-9.7%) lower than those of Choudhury and Sural [1] and about (3.2%-22.6%) higher than those of Elkilany [8]. Our values of the total cross-sections σ_{14} of H(2p) are about (7.1%-29.6%) lower than the available values of Choudhury and Sural [1]. The present calculations show that, we have more H-formed if we are open more excited channels of H-formed in the collision of p-K atoms. The present calculated total cross sections have the same trend of the comparison results and give good agreement with the available previous results of Choudhury and Sural [1].

5. CONCLUSION

p-K scattering is studied using MCSA as a four channels problem (elastic, H(1s), H(2s) and H(2p)). Our interest is focused on the formation of ground, $H(1s)$, and excited hydrogen, $H(2s)$, and $H(2p)$ in p-K inelastic scattering. The difference between the four channels problem and the three or two channels problems (which are used by Elkilany ([6]-[8]) is in improving the total cross sections of the considered channel by adding the effect of more kernels of the other three channels (in two channels problem, we have only one kernel and in three channels, we have two kernels).

REFERENCES

- [1] Choudhury, K. B. and Sural, D. B. Electron Capture in Ground and Excited States in Proton-Alkali-Metal-Atom Collisions. 1992 ; J. Phys. B25, 853-867.
- [2] Daniele, R., Ferrante, G. and Fiordilino, E. Cross-Section of High-Energy Proton-Alkali Atom Charge Exchange in the Eikonal Approximation . 1979; IL Nuovo Cimento B54, 185-196.
- [3] Ferrante,G. and Fiordilino, E. Proton-Alkali Atom Charge Exchange in the Eikonal Approximation. 1980; IL Nuovo Cimento B57, 1-10.
- [4] Ferrante,G., Fiordilino, E. and Zarcone, M. Valence and Core Electron Capture in Proton-Alkali Atom Collisions. 1979; IL nuovo Cimento B52, 151-164 .
- [5] Tiwari, Y. N. Formation of H-Atom in 2s Excited State of Proton-Lithium and Proton-Sodium Scattering. 2008; Pranama-journal of physics., 70, 753-758.
- [6] Elkilany, S. A. Effect of Polarization Potential on Proton–Sodium Inelastic Scattering. 2014; Brz. J. Phys., 44(6), 629-637 .
- [7] Elkilany, S. A. Formation of hydrogen atom in 2s state in proton–sodium inelastic scattering. 2015; Chin. Phys. B 24(3), 033402-033407 .
- [8] Elkilany, S. A. Formation of Ground and Excited Hydrogen Atoms in Proton–Potassium Inelastic Scattering . 2016; Pramana-J Phys., 5,78-87 .
- [9] Clementi, E. and Roetti, C. Roothaan-Hartree -Fock Atomic Wave Functions. 1974; Atomic Data Nuclear Data Tables, 14, 177-478 .

232 **Table 1.** Present σ_{12} , σ_{13} , and σ_{14} (in πa_0^2) of p-K scattering with the results of [1], [2], [4] and [8].

k^2 keV.	Present σ_{11}	Present σ_{12}	Choudhury [1]	Daniele [2]	Ferrante et al. [4]	Elkilany [8]	Present σ_{13}	Choudhury [1]	Elkilany [8]	Present σ_{14}	Choudhury [1]
	Elastic	1s	1s				2s	2s		2p	2p
50	3.9912E-03	2.6012E-03	2.91E-03			2.5110E-03	5.8342E-04	6.38E-04	4.6134E-04	3.4652E-04	3.73E-04
100	1.7304E-03	4.9638E-04	5.51E-04	1.0862E-03	3.4992E-3	4.6494E-04	4.0021E-05	4.34E-05	3.0972E-05	2.2881E-05	3.19E-05
150	5.9509E-04	1.3581E-04	1.52E-04			1.3045E-04	9.6954E-06	1.06E-05	7.6303E-06	5.0585E-06	6.52E-06
200	9.8816E-05	4.6745E-05	5.30E-05			4.5405E-05	3.3239E-06	3.68E-06	2.6438E-06	1.3163E-06	1.87E-06
250	7.4749E-05	4.0914E-05				3.7385E-05	2.2665E-06		2.1942E-06	1.0957E-06	
300	5.7265E-05	3.7593E-05				3.0467E-05	2.0208E-06		1.7726E-06	4.9739E-07	
350	4.3379E-05	2.6831E-05				2.3289E-05	1.4868E-06		1.3497E-06	2.2509E-07	
400	3.0505E-05	1.6885E-05				1.5708E-05	9.7862E-07		9.2164E-07	1.2308E-07	
450	9.9314E-06	9.0918E-06				8.2370E-06	6.2442E-07		4.9136E-07	3.8615E-08	
500	3.1596E-06	8.6363E-07	9.55E-07			8.2923E-07	7.4507E-08	8.05E-08	5.8735E-08	1.7367E-08	2.06E-08
550	5.7985E-07	4.1476E-07					4.4936E-08			1.4209E-08	
600	2.8166E-07	1.9702E-07					2.2458E-08			8.7431E-09	
650	1.9051E-07	1.3089E-07					1.3457E-08			6.7504E-09	
700	1.5603E-07	1.0745E-07					7.2316E-09			3.1171E-09	
750	1.0531E-07	7.7671E-08					4.0742E-09			1.7424E-09	
800	6.0302E-08	4.3181E-08					2.8408E-09			1.2134E-09	
850	3.4993E-08	2.3842E-08					1.5941E-09			9.1426E-10	
900	1.3314E-08	9.1924E-09					1.3325E-09			6.3453E-10	
950	8.4385E-09	6.1925E-09					9.7984E-10			3.0241E-10	
1000	5.2655E-09	3.685E-09					5.9991E-10			1.3694E-10	

233




Fig. 2 σ_{12} (in πa_0^2) of p-K scattering with those of Choudhury and Sural [1].




Fig. 3 σ_{13} (in πa_0^2) of p-K scattering with those of Choudhury and Sural [1].




Fig. 4 σ_{14} (in πa_0^2) of p-K scattering with those of Choudhury and Sural [1].




Fig. 5 Elastic, $H(1s)$, $H(2s)$ and $H(2p)$ cross sections (in πa_0^2) of p-K scattering.