

# Theoretical Study of High-nitrogen Salts Composed of Amine Cations and Pyrazole-based Anions

**Abstract:** A series of energetic salts, which consist of simple amine cations ( $\text{NH}_4^+$ ,  $\text{N}_2\text{H}_5^+$  and  $\text{NH}_2\text{OH}^+$ ) and pyrazole-based anions were designed. The anions were attached by substituents of  $-\text{NH}_2$ ,  $-\text{NO}_2$ ,  $-\text{N}_3$ ,  $-\text{NF}_2$  or  $-\text{CN}$ . The densities and heats of formation (HOFs) of the salts were predicted by using DFT-B3LYP and volume-based thermodynamics methods. The detonation performance and impact sensitivity of the ionic salts were also studied by several empirical formulas. The results show that for the promotion of energetic performance of the salts,  $\text{NH}_3\text{OH}^+$  is the best cation among the three cations. When one substituent is  $-\text{NO}_2$ , another substitution of the  $-\text{NO}_2$ ,  $-\text{N}_3$  or  $-\text{NF}_2$  group is helpful for enhancing the densities of the salts and the substitution of  $-\text{NF}_2$  group presents the best density. The combination of the  $-\text{NO}_2$  group with the  $-\text{NO}_2$ ,  $-\text{N}_3$ ,  $-\text{NF}_2$  or  $-\text{CN}$  group is favorable for improving the HOFs of the salts and the incorporation of  $-\text{N}_3$  shows the greatest HOFs. The synergy of  $-\text{N}_3$  and  $-\text{NF}_2$  group also exhibits the potential to improve the detonation performance of the salts. By comparing with the commonly used explosives RDX and HMX, we screened out the salts with high energy and acceptable sensitivity.

**Keywords:** Crystal density, Density functional theory, Detonation properties, Heats of formation

## 1. Introduction


Along with the growing demand of a new generation of advanced energetic materials to replace some conventional explosives, extensive studies have been focused on finding N-heterocycles with lower impact sensitivity and better detonation performance as well as environmentally friendly<sup>1-4</sup>. This is because N-heterocycles have higher nitrogen content, densities and heats of formation than carbocyclic energetic compounds and are compatible with many energetic groups. By incorporating different energetic functionalized group with heterocyclic backbone, we can get advanced energetic materials to meet ever-changing criteria, such as reducing pollution for the environment, low cost and safe storage.

Over the past years, energetic ionic salts have received a substantial amount of interest as a new class of energetic materials<sup>5-8</sup>. As the cations and anions in the salts can be modified independently, we can produce a large amount of different ionic salts by combining different potential cations and anions<sup>9</sup>. Usually, ionic compounds have regular molecular sorting and compact crystal structure, thus have higher densities than atomically similar nonionic molecules<sup>10</sup>. Furthermore, forming ionic salts is a promising strategy in modifying the sensitivity for energetic materials with strong acidity, such as nitroazoles<sup>11</sup>. Therefore, when the neutral molecules are converted to ionic salts, the detonation performance can be improved significantly.

In the design of high energy density molecules, combining various energetic substituents with a variety of backbone is the most popular strategy<sup>12-14</sup>. Among explosives groups, nitro<sup>15</sup>, nitramino<sup>16</sup>, dinitromethyl<sup>17</sup> and trinitromethyl<sup>18</sup> groups have been proved excellent to enhance the detonation performance of the backbone since they are favorable for increasing the density and the oxygen balance value. Recently, the polynitromethyl energetic derivatives based on N-heterocycles have aroused a large amount of interest. And in the N-heterocyclic structures, the three linked carbon atoms of the pyrazole provide a unique opportunity for the skeleton to be diversified functionalized<sup>3</sup>. For example, Shreeve and co-workers synthesized a new

family of pyrazole with three induced hydrogen bonds by incorporating various energetic functional groups<sup>19</sup>. Therefore, different substituents of pyrazole can produce a variety of properties and finally enhance the detonation performance. What's more, the N-H of the pyrazole can be a reaction site for functionalization and some N-polynitromethyl-substituted pyrazoles have been synthesized<sup>20-21</sup>. Considering the advantages of explosophore groups and pyrazole in improving the detonation properties, their combination is certainly an ideal strategy for designing HEDMs. However, there is still lacking a good understanding of energetic substituents of pyrazole and various counter ions on their property.

On the account of above situations, the salts composed of pyrazole-based anions containing different substituents (-NH<sub>2</sub>, -NO<sub>2</sub>, -N<sub>3</sub>, -NF<sub>2</sub>, -CN) with some simple energetic cations (NH<sub>4</sub><sup>+</sup>, N<sub>2</sub>H<sub>5</sub><sup>+</sup>, NH<sub>3</sub>OH<sup>+</sup>) were designed in Figure 1. We performed density functional theory (DFT)<sup>22</sup> and volume-based thermodynamics calculations<sup>23</sup> to search potential salts with high energy and good stability. Our main purpose is investigating the important role of simple amine cations, different substituents and their synergetic effect in the design of efficient pyrazole-based salts.



**Figure 1.** Frameworks of cations and pyrazole anions.

## 2. Computational method

The Gaussian G09 program package were used to perform the calculations<sup>24</sup>. The geometries of all pyrazole-based salts were fully optimized at the DFT (B3LYP) level with 6-31++G\* basis set which was successfully applied to calculate the properties of energetic ionic salts<sup>25-26</sup>. Since density is the key parameter to predict energetic performance, some approaches have been evolved in order to predict the crystal density more accurately<sup>27-29</sup>. According to these methods, we need to take intermolecular interactions within the crystal into account since the traditional  $M/V$  procedure often leads to large errors. Therefore, the formula for calculating the crystal density of ionic salts has improved as follows<sup>30</sup>:

$$\rho(\text{g/cm}^3) = 1.0260(M/V) + 0.0514(V_s^+/A_s^+) + 0.0419(V_s^-/A_s^-) + 0.0277 \quad (1)$$

where,  $M$  is the molecular mass of the ionic compound and  $V$  is the volume of that compound.  $A_s^+$  is the positive electrostatic potential of the cation and  $V_s^+$  is the average value of the potential, while  $A_s^-$  and  $V_s^-$  are the analogous quantities for an anion. The electrostatic potential was calculated by the Multiwfn program<sup>31</sup>.

When the ionic crystal has the formula unit  $M_pX_q$ , its volume is simply the sum of the volumes of the ions and can be calculated in the following equation<sup>27</sup>:

$$V = pV_M^+ + qV_X^- \quad (2)$$

where  $M$  denotes the cation and  $X$  denotes the anion.

Based on Born-Haber cycle (Scheme 1), the heats of formation of energetic salts were predicted using Eq. (3):

$$\Delta H_f^0(\text{ionic salt, 298K}) =$$

$$\Delta H_f^\circ (\text{cation, 298K}) + \Delta H_f^\circ (\text{anion, 298K}) - \Delta H_L \quad (3)$$

in above equation,  $\Delta H_L$  is the lattice energy of the salts which can be predicted by the formula proposed by Jenkins et al<sup>32</sup> as:

$$\Delta H_L = U_{\text{POT}} + [p(n_M/2 - 2) + q(n_X/2 - 2)]RT \quad (4)$$

the  $n_M$  and  $n_X$  values depend on the nature of the cation  $M_p^+$  and anion  $X_q^-$ , respectively.  $U_{\text{POT}}$  (kJ/mol) is the lattice potential energy which can be calculated by the following expression:

$$U_{\text{POT}} = \gamma(\rho/M)^{1/3} + \delta \quad (5)$$

where  $\rho$  is the predicted density (g/cm<sup>3</sup>),  $M$  is the chemical formula mass (g/mol) of salts and the coefficients  $\gamma$  and  $\delta$  for our designed 1:1(charge ratio) salts are 1981.2kJ·mol<sup>-1</sup>·cm and 103.8 kJ/mol, respectively.

The detonation velocity and detonation pressure of the salts were computed using the empirical Kamlet–Jacobs as shown in the following equations<sup>33</sup>:

$$D (\text{km/s}) = 1.01(NM^{1/2}Q^{1/2})^{1/2}(1+1.30\rho) \quad (6)$$

$$P (\text{GPa}) = 1.558\rho^2NM^{1/2}Q^{1/2} \quad (7)$$


in which,  $N$  stands for the moles of detonation gases which is produced by per gram explosive,  $M$  represents the average molecular weight of these gases (g/mol),  $Q$  denotes the heat of detonation (cal/g) which is related to the HOF difference between products and explosives according to the principle of exothermic reactions, and  $\rho$  stands for the predicted density of salts (g/cm<sup>3</sup>) here.

As the impact sensitivity become increasingly important for energetic materials, there is a simple method developed by Keshavarz<sup>34</sup> to estimate the  $h_{50}$  (cm) as follows:

$$(\log h_{50})_{\text{core}} = -0.584 + 61.62a + 21.53b + 27.96c \quad (8)$$

$$(\log h_{50}) = (\log h_{50})_{\text{core}} + 84.47F^+/MW - 147.1F^-/MW \quad (9)$$

where  $(\log h_{50})_{\text{core}}$  is the core function for prediction of impact sensitivity based on the elemental composition;  $MW$  is the molecular mass of the energetic compound;  $a$ ,  $b$ , and  $c$  are the number of carbon, hydrogen, and nitrogen atoms divided by molecular mass of the energetic compound, respectively. The data of  $F^+$  and  $F^-$  were selected from Ref[34].



**Scheme 1.** Born–Haber cycle for the formation of energetic salts.

### 3. Results and Discussion

#### 3.1 Crystal Density

Density is one of the most significant factor for energetic materials as higher density means that more energy will be packed per unit volume of these materials. Specifically, crystal density can directly influence the detonation performance which is shown in the K-J equation. Here, we investigate a series of energetic salts and study the impact of different combinations of substituents and simple cations on the densities of pyrazole-based salts. The volumes, densities and other related data were listed in Table 1, while Figure 2 displayed a comparison of the densities of different pyrazole-based salts.

As shown in Table 1, some salts possess high densities over 1.9 g/cm<sup>3</sup> and 2.08 g/cm<sup>3</sup>. When the pyrazole-based anions coupled with **A1** or **A3** cation, the densities of the salts are all higher than RDX (1.80


g/cm<sup>3</sup>)<sup>35</sup>. Meanwhile, the salts of **A2** cation also have comparable densities. For the same cation, **B12** anion shows the largest potential for enhancing the densities of the pyrazole-based salts as it has the highest density when incorporated with the cation **A1**. And the salts of **B7**, **B10** or **B16** anion have the lowest densities in the salts of the same cation. For the **B1~B6** anions, all have same -NO<sub>2</sub> group and **B2~B6** anions have another different substitution groups. When the other substituent is -NO<sub>2</sub>, -N<sub>3</sub> or -NF<sub>2</sub> group, the salt contained **B3**, **B4** or **B5** anion has high density as compared to the salts including **B1** anion. This is because the  $\pi$  conjugation between pyrazole and substituents can not only stabilize the structure of energetic materials, but also increase the crystal density. Among **B3**, **B4** and **B5** anions, **B5** anion plays the biggest role in enhancing the crystal density due to -NF<sub>2</sub> group can significantly increase the mass of the salts but affect the molecular volume relatively little. However, when the H atom of the pyrazole is replaced by -NH<sub>2</sub> or -CN group, a decrease of the density can be observed in the salts of **B2** and **B2** anions in comparison to the salts of **B1** anion. This suggests that these two groups are not beneficial for improving the densities of the pyrazole-based salts. For the **B7~B16** anions, the influences of substituents on the densities of the pyrazole anions are further elaborated. When the -NF<sub>2</sub> group is associated with other groups, their densities greatly increase. It is noteworthy that the combination of -N<sub>3</sub> and -NF<sub>2</sub> group has the greatest effect for the promotion of the crystal densities as the salts of **B12** anion possess the largest density when incorporated with the same cation. This proves that the -NF<sub>2</sub> group is the most effective group among the five groups for increasing the densities of the pyrazole-based salts once again.

Figure 2 exhibits a comparison of the densities of three series of energetic salts with anions of different combinations of substituents and cations. By comparing the **A1**, **A2**, **A3** series, it is obvious that their change trends are similar. Among three cations, **A1** is the best cation to increase the densities of the pyrazole-based salts, while **A2** is not so good for promoting the densities of these salts. This is because the ammonium cation has smaller size, resulting in better packing of the salts and less molecular volume than the hydrazine cation.

**Table 1.** Volumes (cm<sup>3</sup>/mol) and densities (g/cm) of the pyrazole salts

Cation	Anion	V	A <sub>s</sub> <sup>+</sup>	V <sub>s</sub> <sup>+</sup>	A <sub>s</sub> <sup>+</sup>	V <sub>s</sub> <sup>+</sup>	$\rho$
A1	B1	182.00	47.50	171.70	240.43	-72.88	1.95
A1	B2	154.58	47.50	171.70	246.67	-72.54	1.94
A1	B3	172.90	47.50	171.70	260.69	-70.75	2.02
A1	B4	175.47	47.50	171.70	256.96	-71.54	2.01
A1	B5	163.32	47.50	171.70	259.26	-71.40	2.05
A1	B6	169.91	47.50	171.70	260.62	-70.28	1.94
A1	B7	180.60	47.50	171.70	225.04	-77.21	1.85
A1	B8	182.87	47.50	171.70	248.48	-73.83	1.89
A1	B9	178.68	47.50	171.70	244.45	-74.94	1.99
A1	B10	179.94	47.50	171.70	246.93	-73.30	1.85
A1	B11	173.35	47.50	171.70	246.26	-72.15	1.86
A1	B12	171.34	47.50	171.70	263.57	-71.05	2.08
A1	B13	170.18	47.50	171.70	265.17	-70.05	1.91
A1	B14	178.10	47.50	171.70	258.75	-70.96	2.03
A1	B15	163.30	47.50	171.70	236.95	-74.15	1.99
A1	B16	160.69	47.50	171.70	235.95	-74.52	1.85
A2	B1	192.53	61.68	152.51	240.43	-72.88	1.86
A2	B2	165.11	61.68	152.51	246.67	-72.54	1.85
A2	B3	183.43	61.68	152.51	260.69	-70.75	1.93
A2	B4	186.00	61.68	152.51	256.96	-71.54	1.92
A2	B5	173.85	61.68	152.51	259.26	-71.40	1.95
A2	B6	180.44	61.68	152.51	260.62	-70.28	1.85

A2	B7	191.13	61.68	152.51	225.04	-77.21	1.76
A2	B8	193.40	61.68	152.51	248.48	-73.83	1.80
A2	B9	189.21	61.68	152.51	244.45	-74.94	1.90
A2	B10	190.47	61.68	152.51	246.93	-73.30	1.76
A2	B11	183.88	61.68	152.51	246.26	-72.15	1.77
A2	B12	181.87	61.68	152.51	263.57	-71.05	1.98
A2	B13	180.71	61.68	152.51	265.17	-70.05	1.82
A2	B14	188.63	61.68	152.51	258.75	-70.96	1.94
A2	B15	173.83	61.68	152.51	236.95	-74.15	1.89
A2	B16	171.22	61.68	152.51	235.95	-74.52	1.76
A3	B1	188.51	67.18	146.42	240.43	-72.88	1.92
A3	B2	161.09	67.18	146.42	246.67	-72.54	1.91
A3	B3	179.41	67.18	146.42	260.69	-70.75	1.99
A3	B4	181.98	67.18	146.42	256.96	-71.54	1.98
A3	B5	169.83	67.18	146.42	259.26	-71.40	2.01
A3	B6	176.42	67.18	146.42	260.62	-70.28	1.91
A3	B7	187.11	67.18	146.42	225.04	-77.21	1.82
A3	B8	189.38	67.18	146.42	248.48	-73.83	1.86
A3	B9	185.19	67.18	146.42	244.45	-74.94	1.96
A3	B10	186.45	67.18	146.42	246.93	-73.30	1.82
A3	B11	179.86	67.18	146.42	246.26	-72.15	1.83
A3	B12	177.85	67.18	146.42	263.57	-71.05	2.04
A3	B13	176.69	67.18	146.42	265.17	-70.05	1.88
A3	B14	184.61	67.18	146.42	258.75	-70.96	2.00
A3	B15	169.81	67.18	146.42	236.95	-74.15	1.96
A3	B16	167.20	67.18	146.42	235.95	-74.52	1.82



**Figure 2.** Comparison of the densities of the pyrazole salts.


### 3.2 Heats of Formation (HOFs)

The heats of formation are usually important for evaluating energetic material because higher HOFs means greater energy content<sup>36</sup>. The designed isodesmic reactions for the anions are given in Scheme 2. Table 2 lists the calculated and experimental gas-phase HOFs of small molecules and ions at 298 K and the calculated values are very close to the literature values<sup>9,37</sup>. The G2 method was used to calculate the HOFs of small molecules and ions<sup>38</sup>.

It is seen in Table 3 that the HOFs of cations are all large, while the anions are not so beneficial for improving the HOFs of the pyrazole-based salts except for **B4**, **B8**, **B11**, **B12** or **B13** anion. Especially, for the same cation, the salt containing **B4** anion has the greatest HOF. This indicates that the -N<sub>3</sub> group is the

best substituent to enhance the HOFs of pyrazole-based anions among the five groups. When the other substituent is  $-\text{NO}_2$ ,  $-\text{NF}_2$  or  $-\text{CN}$ , its substituted pyrazole-based salts have slightly higher HOFs than its unsubstituted one, while for the substituent  $-\text{NH}_2$ , the HOFs of the salts are lower. However, when the  $-\text{NH}_2$  group is coupled with the  $-\text{N}_3$  group, it can promote the HOFs of the salts significantly. By comparing the **B11**, **B12** and **B13** anions, **B11** anion has less substituents but possess higher HOFs than **B12** anion when they form salts with the same cation. It is interesting that the combination of  $-\text{N}_3$  group and  $-\text{NF}_2$  group is not so good for the HOFs of the pyrazole-based salts.

Figure 3 presents a comparison of the HOFs for the salts composed of pyrazole-based anions and ammonium, hydrazine and hydroxylamine cations. It is seen that the variation trends of the salts of the three cations are unanimous. This shows that incorporating different cations hardly affects the change trends of the HOFs of the salts even if they have anions with different substituents. When the cation is  $\text{N}_2\text{H}_5^+$ , its corresponding salts with the same anion have the largest HOFs among the salts. This is because this cation has higher nitrogen content than the other two.




**Scheme 2.** Isodesmic reactions for the anions.

**Table 2.** Calculated and experimental gas-phase heats of formation (kJ/mol) for small molecules and ions at 298 K

Molecules/ions	$\Delta H_f^\circ$ (calculated)	$\Delta H_f^\circ$ (literature)
$\text{CH}_4$	-73.6 <sup>a</sup>	-74.4 <sup>c</sup>
$\text{NH}_3$	-45.3 <sup>a</sup>	-46.1 <sup>c</sup>
$\text{CH}_3\text{NO}_2$	-85.9 <sup>a</sup>	-80.8 <sup>c</sup>
$\text{CH}_3\text{NH}_2$	-23.2 <sup>a</sup>	-22.5 <sup>c</sup>
$\text{CH}_3\text{N}_3$	295.8 <sup>a</sup>	296.5 <sup>c</sup>
$\text{CH}_3\text{NF}_2$	-109.2 <sup>a</sup>	
$\text{CH}_3\text{CN}$	75.2 <sup>a</sup>	74 <sup>c</sup>
$\text{H}^+$		1536.2 <sup>c</sup>
$\text{CH}_2\text{NH}_2^-$	196.11 <sup>b</sup>	
$\text{NH}_4^+$	630.51 <sup>b</sup>	626.4 <sup>f</sup>
$\text{N}_2\text{H}_5^+$	760.37 <sup>b</sup>	759.7 <sup>f</sup>
$\text{NH}_3\text{OH}^+$	668.58 <sup>b</sup>	664.4 <sup>f</sup>
pyrazole	183.01 <sup>a</sup>	

<sup>a</sup>The values were calculated at G2 level. <sup>b</sup>The values were calculated by protonation reactions:  $\text{CH}_2\text{NH}_2^- + \text{H}^+ \rightarrow \text{CH}_3\text{NH}_2$ ,  $\text{NH}_3 + \text{H}^+ \rightarrow \text{NH}_4^+$ ,  $\text{N}_2\text{H}_4 + \text{H}^+ \rightarrow \text{N}_2\text{H}_5^+$ ,  $\text{NH}_2\text{OH} + \text{H}^+ \rightarrow \text{NH}_3\text{OH}^+$ . <sup>c</sup>The values were taken from Ref[36]. <sup>f</sup>The values were taken from Ref [9]

182



183

184

**Figure 3.** Comparison of the HOFs of the pyrazole salts.

185

186

187

**Table 3.** Heats of formation ((kJ/mol) for the pyrazole anions, amine cations, and their salts and lattice energies of these salts

188

Cation	Anion	$\Delta H_f^0(\text{cation})$	$\Delta H_f^0(\text{anion})$	Lattice energy	$\Delta H_f^0(\text{salt})$
A1	B1	630.51	-197.54	487.45	-54.49
A1	B2	630.51	-233.43	480.41	-83.33
A1	B3	630.51	-168.88	473.61	-11.98
A1	B4	630.51	528.80	474.59	684.72
A1	B5	630.51	-191.14	472.86	-33.49
A1	B6	630.51	-42.28	476.41	111.81
A1	B7	630.51	-179.13	495.12	-43.75
A1	B8	630.51	294.75	478.93	446.33
A1	B9	630.51	-220.46	481.14	-71.09
A1	B10	630.51	-64.15	483.07	83.29
A1	B11	630.51	159.83	483.21	307.13
A1	B12	630.51	94.38	475.93	248.96
A1	B13	630.51	297.24	475.82	451.93
A1	B14	630.51	-67.63	479.52	83.36
A1	B15	630.51	-216.12	487.17	-72.79
A1	B16	630.51	-55.93	490.23	84.34
A2	B1	760.37	-197.54	477.67	85.16
A2	B2	760.37	-233.43	471.05	55.90
A2	B3	760.37	-168.88	464.85	126.64
A2	B4	760.37	528.80	465.75	823.42
A2	B5	760.37	-191.14	464.20	105.03
A2	B6	760.37	-42.28	467.30	250.79
A2	B7	760.37	-179.13	484.56	96.68
A2	B8	760.37	294.75	469.54	585.58
A2	B9	760.37	-220.46	471.85	68.05
A2	B10	760.37	-64.15	473.30	222.92
A2	B11	760.37	159.83	473.46	446.75
A2	B12	760.37	94.38	467.15	387.60
A2	B13	760.37	297.24	466.66	590.95
A2	B14	760.37	-67.63	470.43	222.31
A2	B15	760.37	-216.12	477.50	66.75
A2	B16	760.37	-55.93	480.00	224.43
A3	B1	668.58	-197.54	475.95	-4.91



A3	B2	668.58	-233.43	469.50	-34.35
A3	B3	668.58	-168.88	463.41	36.30
A3	B4	668.58	528.80	464.29	733.09
A3	B5	668.58	-191.14	462.76	14.69
A3	B6	668.58	-42.28	465.85	160.45
A3	B7	668.58	-179.13	482.72	6.73
A3	B8	668.58	294.75	468.06	495.27
A3	B9	668.58	-220.46	470.25	-22.14
A3	B10	668.58	-64.15	471.76	132.67
A3	B11	668.58	159.83	471.91	356.51
A3	B12	668.58	94.38	465.62	297.34
A3	B13	668.58	297.24	465.24	500.58
A3	B14	668.58	-67.63	468.84	132.11
A3	B15	668.58	-216.12	475.75	-23.30
A3	B16	668.58	-55.93	478.29	134.36

### 3.3 Energetic properties and sensitivity

According to our predicted densities and HOFs, we calculated the corresponding energetic parameters which were summarized in Table 4. Meanwhile, commonly used explosives RDX and HMX were also listed for comparison<sup>39</sup>.

The calculated heat of detonation (Q) in Table 4 show that for the **B1~B6** anions, the substitution of the -NO<sub>2</sub>, -N<sub>3</sub> and -NF<sub>2</sub> group increases the Q value of its salts as compared to the salts incorporated with **B1** anion(who don't have the other substituent), whereas for the substituent -NH<sub>2</sub> and -CN, the case is quite the contrary. Especially, **B4** anion presents the best potential for enhancing the Q value among the six anions and the salts of **B4** anion even has larger Q than HMX. By comparing the **B7~B16** anions, the influences of different substituents on the Q value are further presented. When the two substituents are -N<sub>3</sub> and -NF<sub>2</sub> group (anion **B12**), the Q of its salt is the best among the ten anions but is lower than **B3**, **B4** and **B5** anions. When the two substituents are -NH<sub>2</sub> and -CN (anion **B10**), the Q value of its salt is the lowest. This indicates that the -NO<sub>2</sub>, -N<sub>3</sub> and -NF<sub>2</sub> groups are effective substituents for enhancing the Q of the pyrazole-based salts.

As is shown in Table 4, for the **B1~B6** anions, the salts of **B3~B5** have larger D and P than the salts of **B1**. On the contrary, the salts of **B2** and **B6** anions have smaller ones. This indicates that the substitution of group -NO<sub>2</sub>, -N<sub>3</sub> and -NF<sub>2</sub> can enhance the D and P of the pyrazole-based salts. Among the six anions, the salts of **B4** has relatively larger D and P values which means that the substituent -N<sub>3</sub> is the best group for increasing the D and P. For the **B7~B16** anions, the combination of different substituents slightly enhances D and P of their salts except **B9**, **B12** and **B15** anions, indicating that the -NO<sub>2</sub> group is important for increasing the energetic properties. When the two substituents are -N<sub>3</sub> and -NF<sub>2</sub> group (anion **B12**), the D and P of its salts is the largest among **B7~B16** anions and only slightly lower than the **B4** and **B5** anions. It is concluded that the -NO<sub>2</sub>, -N<sub>3</sub> or -NF<sub>2</sub> group is favorable for increasing the detonation properties of its pyrazole-based salts and the combination of -N<sub>3</sub> and -NF<sub>2</sub> group is also beneficial for enhancing the energetic properties.

Figure 4 exhibits a comparison of the calculated Q, D and P values of the pyrazole-based salts. The **A1**, **A2** and **A3** cations hardly influence the detonation properties. Among **A1~A3** cations, **A3** cation has the largest potential to enhance the detonation properties. By comparing the detonation properties of these salts with RDX and HMX, we can find that the salts of **B3**, **B4**, **B5** and **B12** anion have better energetic properties than RDX and HMX.

Impact sensitivity has been more and more important in the field of high energy density materials. The impact sensitivity is generally reported as the height in cm, designated  $h_{50}$ <sup>40</sup>. The higher the  $h_{50}$  is, the less




sensitive the explosive is. Figure 5 presents the influence of different combination of substituents and amine cations on the  $h_{50}$  of the pyrazole-based salts. When the anion is same, the salts of **A2** series have the largest  $h_{50}$  among the three series salts, the middle is the salts of **A1** series and the salts of **A3** series have the lowest  $h_{50}$ . This indicates that the **A2** cation is good for reducing the impact sensitivity of the pyrazole-based salts. For the salts of **B1~B6** anions and the same cation, the  $h_{50}$  of salts containing **B6** anion are larger than the salts of **B1** anion, while the  $h_{50}$  of the salts of **B3**, **B4** and **B5** anion are smaller. Among the **B7~B16** anions, when the cation is same, the salts of **B12** anion have the smallest  $h_{50}$ . As a whole, incorporating -N<sub>3</sub>, -NO<sub>2</sub> or -NF<sub>2</sub> group into pyrazole-based salts is not helpful for reducing the impact sensitivity while incorporating -NH<sub>2</sub> or -CN group can reduce the impact sensitivity. When the two substituents are -N<sub>3</sub> and -NF<sub>2</sub>, it is also not so good for reducing the impact sensitivity of the pyrazole-based salts. Furthermore, it can be found that the better the detonation properties are, the larger the sensitivity is.

**Table 4.** Predicted heats of detonation (Q), detonation velocities (D) and pressures (P), and sensitivity for the pyrazole salts


Cation	Anion	Q(cal/g)	D(km/s)	P(GPa)	$h_{50}$ (cm)
A1	B1	1397.79	9.06	38.10	45.62
A1	B2	1321.41	8.95	37.16	45.23
A1	B3	1508.45	9.58	43.53	23.31
A1	B4	1753.67	9.85	45.94	17.57
A1	B5	1558.54	9.77	45.62	21.48
A1	B6	1279.46	8.65	34.74	49.95
A1	B7	1242.37	8.40	31.76	108.20
A1	B8	1451.81	8.93	36.42	33.42
A1	B9	1407.12	9.29	40.62	40.80
A1	B10	1091.02	7.95	28.49	110.58
A1	B11	1390.44	8.67	33.94	33.16
A1	B12	1496.30	9.70	45.28	16.26
A1	B13	1254.02	8.43	32.64	37.33
A1	B14	1331.86	9.06	39.06	45.11
A1	B15	1460.81	9.31	40.73	40.92
A1	B16	1134.34	7.97	28.61	117.52
A2	B1	1458.31	8.89	35.75	51.63
A2	B2	1382.37	8.80	34.90	50.91
A2	B3	1555.32	9.37	40.57	26.74
A2	B4	1790.18	9.62	42.61	20.44
A2	B5	1602.35	9.55	42.41	24.67
A2	B6	1340.30	8.52	32.77	55.76
A2	B7	1319.30	8.29	30.12	118.25
A2	B8	1507.18	8.76	34.11	38.24
A2	B9	1462.79	9.11	37.96	46.05
A2	B10	1168.50	7.88	27.22	119.80
A2	B11	1452.15	8.53	31.94	38.22
A2	B12	1543.45	9.49	42.15	18.93
A2	B13	1316.88	8.31	30.86	42.31
A2	B14	1389.15	8.90	36.67	50.49
A2	B15	1516.89	9.13	38.06	46.45
A2	B16	1214.08	7.90	27.37	127.45
A3	B1	1521.45	9.21	39.04	34.47
A3	B2	1442.56	9.10	38.03	34.66
A3	B3	1609.62	9.66	43.84	18.87
A3	B4	1844.43	9.90	45.99	14.38
A3	B5	1655.58	9.84	45.79	17.52

A3	B6	1398.67	8.81	35.63	38.41
A3	B7	1390.22	8.63	33.26	75.19
A3	B8	1567.70	9.06	37.15	25.93
A3	B9	1521.62	9.42	41.33	31.59
A3	B10	1233.58	8.19	29.93	79.20
A3	B11	1516.04	8.83	34.89	25.40
A3	B12	1597.52	9.79	45.62	13.40
A3	B13	1376.04	8.60	33.61	29.03
A3	B14	1446.39	9.20	39.88	35.03
A3	B15	1578.59	9.45	41.53	31.27
A3	B16	1282.70	8.21	30.17	82.36
RDX		1597.40	8.9(8.8 <sup>a</sup> )	34.8(34.7 <sup>a</sup> )	29.17(26.33 <sup>a</sup> )
HMX		1633.90	9.3(9.1 <sup>a</sup> )	39.2(39.0 <sup>a</sup> )	31.28(29.36 <sup>a</sup> )

<sup>a</sup> The values were taken from Ref[38]



**Figure 4.** Heats of detonation, detonation velocities, and detonation pressures of the pyrazole salts.



**Figure 5.** Comparison of the impact sensitivity of the pyrazole salts.

#### 4. Conclusion

A series of salts composed of simple amine cations ( $\text{NH}_4^+$ ,  $\text{N}_2\text{H}_5^+$  and  $\text{NH}_2\text{OH}^+$ ) and pyrazole-based anions were designed. The densities, heats of formation (HOFs) and energetic performance of the salts were studied. The impact sensitivity of the salts were predicted using the elemental composition and relevant correction factor. The results show that for the promotion of densities of the pyrazole-based salts,  $\text{NH}_4^+$  has the best among the three amine cations. When one substituent is  $-\text{NO}_2$  group, another substitution of the  $-\text{NO}_2$ ,  $-\text{N}_3$  or  $-\text{NF}_2$  group is helpful for increasing the densities of the salts. The substitution of  $-\text{NF}_2$  group shows the best promotion for the densities of the salts, especially when the  $-\text{NF}_2$  group attaches to the pyrazole ring with  $-\text{N}_3$  group. The salts of  $\text{N}_2\text{H}_5^+$  exhibit the best HOFs due to its highest nitrogen content. When one substituent is  $-\text{NO}_2$  group, incorporating another  $-\text{NO}_2$ ,  $-\text{N}_3$ ,  $-\text{NF}_2$  and  $-\text{CN}$  group into salts is

favorable for improving the HOFs of the salts, with -N<sub>3</sub> group being the most favorable.

The detonation performance of these salts indicates that NH<sub>3</sub>OH<sup>+</sup> has better detonation performance than the other two amine cations. When one substituent is -NO<sub>2</sub> group, introducing another -NO<sub>2</sub>, -N<sub>3</sub> or -NF<sub>2</sub> group is an effective way for promoting the detonation properties of the salts. When the two substituents are -N<sub>3</sub> and -NF<sub>2</sub> group, there is a synergistic effect that is good for enhancing the detonation properties of the salts. By comparing with the commonly used explosives RDX and HMX, it can be found that the salts of **B3**, **B4**, **B5** or **B12** anion can be considered as potential high-energy ionic salts.

## References

1. Türker, L., Azo-bridged triazoles: Green energetic materials. *Defence Technology* **2016**, *12* (1), 1-15.
2. Sebastiao, E.; Cook, C.; Hu, A.; Murugesu, M., Recent developments in the field of energetic ionic liquids. *Journal of Materials Chemistry A* **2014**, *2* (22), 8153-8173.
3. Yin, P.; Shreeve, J. M., Chapter Four - Nitrogen-Rich Azoles as High Density Energy Materials: Reviewing the Energetic Footprints of Heterocycles. In *Advances in Heterocyclic Chemistry*, Academic Press: 2017; Vol. 121, pp 89-131.
4. Chavez, D. E., Energetic Heterocyclic N-Oxides. In *Heterocyclic N-Oxides*, Larionov, O. V., Ed. Springer International Publishing: Cham, 2017; pp 1-27.
5. Gao, H.; Shreeve, J. M., Azole-based energetic salts. *Chemical Reviews* **2011**, *111* (11), 7377-436.
6. Zhao, Z.; Du, Z.; Han, Z.; Zhang, Y.; He, C., Nitrogen-Rich Energetic Salts: Both Cations and Anions Contain Tetrazole Rings. *Journal of Energetic Materials* **2015**, *34* (2), 183-196.
7. Klapötke, T. M.; Witkowski, T. G., Covalent and Ionic Insensitive High-Explosives. *Propellants, Explosives, Pyrotechnics* **2016**, *41* (3), 470-483.
8. Schmidt, M. W.; Gordon, M. S.; Boatz, J. A., Triazolium-Based Energetic Ionic Liquids. *The Journal of Physical Chemistry A* **2005**, *109* (32), 7285-7295.
9. Gao, H.; Ye, C.; Piekarski, C. M.; Shreeve, J. M., Computational Characterization of Energetic Salts. *The Journal of Physical Chemistry C* **2007**, *111* (28), 10718-10731.
10. Ghule, V. D., Computational Screening of Nitrogen-Rich Energetic Salts Based on Substituted Triazine. *The Journal of Physical Chemistry C* **2013**, *117* (33), 16840-16849.
11. Zhao, X. X.; Li, S. H.; Wang, Y.; Li, Y. C.; Zhao, F. Q.; Pang, S. P., Design and synthesis of energetic materials towards high density and positive oxygen balance by N-dinitromethyl functionalization of nitroazoles. *Journal of Materials Chemistry A* **2016**, *4* (15), 5495-5504.
12. Yu, Q.; Yin, P.; Zhang, J.; He, C.; Imler, G. H.; Parrish, D. A.; Shreeve, J. M., Pushing the Limits of Oxygen Balance in 1,3,4-Oxadiazoles. *Journal of the American Chemical Society* **2017**, *139* (26), 8816-8819.
13. Xu, Y.; Shen, C.; Lin, Q.; Wang, P.; Jiang, C.; Lu, M., 1-Nitro-2-trinitromethyl substituted imidazoles: a new family of high performance energetic materials. *Journal of Materials Chemistry A* **2016**, *4* (45), 17791-17800.
14. Tang, Y.; He, C.; Gao, H.; Shreeve, J. M., Energized nitro-substituted azoles through ether bridges. *Journal of Materials Chemistry A* **2015**, *3* (30), 15576-15582.
15. Ravi, P.; Gore, G. M.; Tewari, S. P.; Sikder, A. K., Quantum Chemical Studies on the Structure and Performance Properties of 1,3,4,5-Tetranitropyrazole: A Stable New High Energy Density Molecule. *Propellants, Explosives, Pyrotechnics* **2012**, *37* (1), 52-58.
16. Huang, H.; Shi, Y.; Liu, Y.; Yang, J., 1,2,4,5-Dioxadiazine-functionalized [N-NO<sub>2</sub>]<sup>-</sup> furazan energetic salts. *Dalton Transactions* **2016**, *45* (39), 15382-15389.
17. Klapötke, T. M.; Mayr, N.; Stierstorfer, J.; Weyrauther, M., Maximum compaction of ionic organic explosives: bis(hydroxylammonium) 5,5'-dinitromethyl-3,3'-bis(1,2,4-oxadiazolate) and its derivatives. *Chemistry - A European Journal* **2014**, *20* (5), 1410-7.

18. Dharavath, S.; Zhang, J.; Imler, G. H.; Parrish, D. A.; Shreeve, J. M., 5-(Dinitromethyl)-3-(trinitromethyl)-1,2,4-triazole and its derivatives: a new application of oxidative nitration towards gem-trinitro-based energetic materials. *Journal of Materials Chemistry A* **2017**, 5 (10), 4785-4790.
19. Yin, P.; Parrish, D. A.; Shreeve, J. M., Energetic multifunctionalized nitraminopyrazoles and their ionic derivatives: ternary hydrogen-bond induced high energy density materials. *Journal of the American Chemical Society* **2015**, 137 (14), 4778-86.
20. Daling, I. L.; Vatsadze, I. A.; Shkineva, T. K.; Kormanov, A. V.; Struchkova, M. I.; Suponitsky, K. Y.; Bragin, A. A.; Monogarov, K. A.; Sinditskii, V. P.; Sheremetev, A. B., Novel Highly Energetic Pyrazoles: N-Trinitromethyl-Substituted Nitropyrazoles. *Chemistry – An Asian Journal* **2015**, 10 (9), 1987-96.
21. Yin, P.; Zhang, J.; Mitchell, L. A.; Parrish, D. A.; Shreeve, J. M., 3,6-Dinitropyrazolo[4,3-c]pyrazole-Based Multipurpose Energetic Materials through Versatile N-Functionalization Strategies. *Angewandte Chemie International Edition* **2016**, 55 (41), 12895-12897.
22. Ghule, V. D.; Deswal, S.; Devi, A.; Kumar, T. R., Computer-Aided Design of Energetic Tris(tetrazolyl)amine Derivatives and Salts. *Industrial & Engineering Chemistry Research* **2016**, 55 (4), 875-881.
23. Glasser, L.; Jenkins, H. D. B., Volume-Based Thermodynamics: A Prescription for Its Application and Usage in Approximation and Prediction of Thermodynamic Data. *Journal of Chemical & Engineering Data* **2011**, 56 (4), 874-880.
24. Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G. A.; Nakatsuji, H.; Caricato, M. Li. X.; Hratchian, H. P.; Izmaylov, A. F.; Bloino, J.; Zheng, G.; Sonnenberg, J. L.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Montgomery, J. J. A.; Peralta, J. E.; Ogliaro, F.; Bearpark, M.; Heyd, J. J.; Brothers, E.; Kudin, K. N.; Staroverov, V. N.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Rega, N.; Millam, J. M.; Klene, M.; Knox, J. E.; Cross, J. B.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Martin, R. L.; Morokuma, K.; Zakrzewski, V. G.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Dapprich, S.; Daniels, A. D.; Farkas, O.; Foresman, J. B.; Ortiz, J. V.; Cioslowski, J.; Fox, D. J. *Gaussian 09, Revision A.02*; Gaussian: Wallingford, CT, 2009.
25. Xiang, F.; Wu, Q.; Zhu, W.; Xiao, H., Comparative Theoretical Studies on Energetic Ionic Salts Composed of Heterocycle-Functionalized Nitraminofurazanate-Based Anions and Triaminoguanidinium Cation. *Journal of Chemical & Engineering Data* **2013**, 59 (2), 295-306.
26. Xiang, F.; Zhu, W.; Xiao, H., Theoretical studies of energetic nitrogen-rich ionic salts composed of substituted 5-nitroiminotetrazolate anions and various cations. *Journal of Molecular Modeling* **2013**, 19 (8), 3103-18.
27. Rice, B. M.; Hare, J. J.; Byrd, E. F. C., Accurate Predictions of Crystal Densities Using Quantum Mechanical Molecular Volumes. *The Journal of Physical Chemistry A* **2007**, 111 (42), 10874-10879.
28. Pan, J. F.; Lee, Y. W., Crystal density prediction for cyclic and cage compounds. *Physical Chemistry Chemical Physics* **2004**, 6 (3), 471.
29. Qiu, L.; Xiao, H.; Gong, X.; Ju, X.; Zhu, W., Crystal density predictions for nitramines based on quantum chemistry. *Journal of Hazardous Materials* **2007**, 141 (1), 280-288.
30. Politzer, P.; Martinez, J.; Murray, J. S.; Concha, M. C., An electrostatic correction for improved crystal density predictions of energetic ionic compounds. *Molecular Physics* **2010**, 108 (10), 1391-1396.
31. Lu, T.; Chen, F., Multiwfn: A multifunctional wavefunction analyzer. *Journal of Computational Chemistry* **2012**, 33 (5), 580-592.
32. Jenkins, H. D. B.; Tudela, D.; Glasser, L., Lattice Potential Energy Estimation for Complex Ionic Salts from Density Measurements. *Inorganic Chemistry* **2002**, 41 (9), 2364-2367.
33. Kamlet, M. J.; Jacobs, S. J., Chemistry of Detonations. I. A Simple Method for Calculating Detonation Properties of C–H–N–O Explosives. *The Journal of Chemical Physics* **1968**, 48 (1), 23-35.

- 340 34. Keshavarz, M. H., A New General Correlation for Predicting Impact Sensitivity of Energetic Compounds. *Propellants,*  
341 *Explosives, Pyrotechnics* **2013**, 38 (6), 754-760.
- 342 35. Badgajar, D. M.; Talawar, M. B.; Asthana, S. N.; Mahulikar, P. P., Advances in science and technology of modern  
343 energetic materials: An overview. *Journal of Hazardous Materials* **2008**, 151 (2), 289-305.
- 344 36. Li, X.; Tang, Z.; Zhang, X.; Yang, X., The heats of formation in a series of nitroester energetic compounds: A theoretical  
345 study. *Journal of Hazardous Materials* **2009**, 165 (1), 372-378.
- 346 37. Lide D R., Handbook of chemistry and physics, 84th ed. Boca Raton: CRC Press, 2003-2004(Section 5).
- 347 38. Curtiss, L. A.; Raghavachari, K.; Trucks, G. W.; Pople, J. A., Gaussian-2 theory for molecular energies of first- and  
348 second-row compounds. *The Journal of Chemical Physics* **1991**, 94 (11), 7221-7230.
- 349 39. Talawar, M. B.; Sivabalan, R.; Mukundan, T.; Muthurajan, H.; Sikder, A. K.; Gandhe, B. R.; Rao, A. S., Environmentally  
350 compatible next generation green energetic materials (GEMs). *Journal of Hazardous Materials* **2009**, 161 (2), 589-607.
- 351 40. Li, J., Relationships for the Impact Sensitivities of Energetic C-Nitro Compounds Based on Bond Dissociation Energy.  
352 *The Journal of Physical Chemistry B* **2010**, 114 (6), 2198-2202.