

Original Research Article**Free Radical Scavenging Activity of *Jatropha curcas* Leaves,
Phytochemical And Antibacterial Analysis of its Butanol Fraction.****ABSTRACT**

The use of different plant parts for the prevention and cure of ailments, infections and diseases as an alternative to the use of orthodox medicine is gaining more attention. Moreso, resistance to orthodox drugs has been confirmed in various literatures. The leaves of *Jatropha curcas*, an ever-green perennial plant was used in this study. They were washed, air-dried and pulverized for active extraction of the composition of the plant in butanol. Fresh leaves of the plant were tested for its DPPH scavenging activity (indicator of antioxidant), while the butanol leaf extract was accessed for the presence or absence of phytochemicals and was also tested against some gram positive and gram negative bacteria. The result of the free radical scavenging activity of the leaves was higher to that of ascorbic acid (a standard antioxidant). The qualitative analysis showed the presence of alkaloids, tannins, saponin, flavonoids, steroid and cardiac glycoside. The quantitative analysis on the other hand showed the presence of alkaloids in an appreciable amount, followed by saponin while others had relatively low values. The average values of the antibacterial (inhibitory) activity as compared with standard antibiotics used had no significant difference except for *pseudomonas sp.* which was very low to the antibiotics used. It

could be concluded that *Jatropha curcas* is of great medicinal potential which could be explored further.

Keywords; *Jatropha curcas*, Antibacterial, Scavenging activity, Phytochemicals.

1. INTRODUCTION

Ethnomedicine refers to the study of traditional medicinal practice which is concerned with the cultural interpretation of health diseases and illness and also addresses the health care seeking process and healing practices[1].

Ethnomedicinal plants have been used since ancient time for human healthcare and still remain the most widely used medication system in developing and least developed nations [2].

The reliance of people on ethnomedicine has been for reasons of cost-effectiveness, acceptability, biomedical benefits and accessibility. However, there has been a continuous growth of demand for herbal medicine globally [3].

Synthetic drugs have become expensive and not easily accessible by the less privileged. Moreso, resistance to most synthetic drugs is a serious health concern in the world today. In addition, people have returned to traditional medicine and natural plants are now used as drugs for various ailments based on their folkloric uses. Plant derived-bioactive compounds have received considerable attention due to their therapeutic potential as antimicrobial, anti-inflammatory properties and antioxidant activities [4].

Jatropha curcas leaves has been used as cure for various ailments like; skin infection, diarrhea, cancer e.t.c.

Therefore, this research is targeted at evaluating the free radical scavenging activities of *Jatropha curcas* leaves, phytochemicals present in its butanol fraction as well as the antibacterial activities of its butanol fraction in order to ascertain some of its claims in traditional healing.

1.1 Free radicals and DPPH scavenging activities

Free radicals are fundamental to any biochemical process and represent an essential part of aerobic life and metabolism [5]. Reactive oxygen (ROS) and reactive nitrogen species (RNS) are products of normal cellular metabolism. The most common ROS include superoxide anion Peroxy nitrite anion [6]. These reactive species play an important role in pathogenesis, cardiovascular diseases, and rheumatoid ulcerative colitis. It is possible to reduce the risk of chronic diseases and prevent diseases progression by either enhancing the body's natural antioxidant defenses or by supplementing with dietary antioxidants [7]. Antioxidant offer resistance against oxidative stress by scavenging the free radicals inhibiting the lipid peroxidation and by many other mechanisms and thus prevent the diseases progression. DPPH is a stable nitrogen centered free radical commonly used for testing radical scavenging activity of the compound or plant extracts. When stable DPPH radical accept an electron from the antioxidant compound the violet color of the DPPH reduce to yellow color or red. Diphenylpicrylhydrazyl radical which was measured colorimetrically. Substance which are able to perform this reaction can be considered as antioxidant and therefore a radical scavenger [8].

1.2 Antibacterial

Antibacterial also known as antibiotics is a type of antimicrobial drug used in the treatment and prevention of bacterial infection. They may kill or inhibit the growth of bacterial. They are not effective against viruses such as common cold or influenza. Antibiotics may be given as a preventive measure (prophylactic) and this is usually limited to at risk populations such as those with a weakened immune system (particularly in HIV cases to prevent pneumonia) [9].

2.0 MATERIALS AND METHODS

2.1 Experimental material

Some of the experimental apparatus used for this work included;

Beakers, test tubes, petri dishes, conical flasks, boiling tubes, Bunsen burner, measuring cylinder

Experimental material included; *J. curcas* leaves,

Reagents included;

HCl, H₂SO₄, Ferric chloride solution, Potassium iodide, Mayer's reagent, Butanol, Ethanol, Ethyl acetate, Dilute ammonia, Potassium ferrocyanide, Acetic acid, Olive oil, MacConkey agar, Nutrient agar.

Instruments included

Weighing balance, UV - spectrophotometer

2.2 Experimental procedure

Sample collection: The plant sample was collected around the quarters of the Federal college of Animal Health and production technology. The plant samples was air-dried to preserve some phytochemical constituent which are thermolabile and could be denatured by heat and also to prevent physiological change of the leaves.

The air-dried samples were pulverized, using an electric blender in other to reduce them to fine particles for effective extraction of the bioactive compounds.

Exactly, 100g of the leaves were weighed into a beaker and butanol was poured into it until the samples were fully immersed, the extraction was carried out by decanting the butanol (supernatant) every 24 hours. This was replaced with fresh volumes of butanol and was repeated for 10days.

The samples were concentrated using rotary evaporator until all the butanol had evaporated and the extract was obtained.

2.3. DPPH free radical scavenging Activity

100g of fresh sample was weighed into a beaker, 100ml of ethanol was added and shaken vigorously for 2min, it was stirred with a magnetic stirrer for 15min, it was allowed to stand for 2hours for proper extraction, then centrifuged at 2500rpm for 10min. The supernatant was poured into another beaker and concentrated by

evaporating in a water bath at 80⁰C. The concentrated extract was kept for further assay.

1mM DPPH was prepared in ethanol (394.32mg DPPH dissolved in 1litre of ethanol. It was dissolved with 10mg of the concentrated sample extract in 10ml of ethanol (1mg/ml) 1.5ml of the extract was pipette into a test tube. 1.5ml DPPH solution prepared was added into the test tube. The spectrophotometer was zeroed with ethanol as blank. Thereafter the absorbance\optical density of the control (DPPH solution) and that of the sample was read at 517nm.

$$\text{DPPH Scavanged \%} = \frac{\text{absorbance of control} - \text{absorbance of test sample} \times 100}{\text{absorbance of control} + \text{sample weight}}$$

This procedure was also carried out for Vitamin C (ascorbic acid) a standard antioxidant.

2.4 Qualitative phytochemical Analysis

2.4.1 Test for Alkaloids:

0.5g of the plant sample was dissolved in 5ml dilute HCl in a steam bath and filtered. 1ml of the above filterate was treated with few drops of Mayer's reagent. A creamy white precipitate was observed which indicated the presence of alkaloids.

2.4.2 Test for Tannin

Exactly 0.5g of each plant extract was stirred with about 20mi of distilled water and then filtered. Few drops of 0.1% ferric chloride solution were added to the filtrate. A dirty green precipitate was observed, indicative of the presence of tannin.

121 **2.4.3 Test for saponin**

122 2 gram of each portion was boiled with 20ml of distilled water filtered. To the
123 filtrate about 3ml of distilled water was further added and shake vigorously for
124 about 5min frothing which persisted on warming was observed. This was a positive
125 test for saponin.

126 **2.4.4 Test for flavonoids**

127 A portion of the powdered plant sample was heated with 10ml ethyl acetate over a
128 steam bath for 3min. it was filtered and 4ml of the filtrate was shaken with 1ml of
129 dilute ammonia .A yellow coloration indicating the presence of flavonoids was
130 observed

131 **2.4.5 Test for steroids**

132 About 0.2g of the plant extract, 2ml of acetic acid was added and the solution
133 was cooled well, Conc. H_2SO_4 was added to the solution. A violet to blue color
134 indicates the presence of a steroidal ring.

135

136 **2.5 Quantitative phytochemical analysis**

137 **2.5.1 Test for Alkaloids**

138 2g of the extracted sample was weighed into a 100ml beaker and 20ml of 80%
139 absolute alcohol added to give a smooth paste. The mixture was transferred to a
140 250ml flask and more alcohol added to make up to 100ml and 1g magnesium

oxide added. The mixture was digested in boiling water for 1.5 hours under a reflux air condenser with occasional shaking. The mixture was filtered while hot through a small bucher funnel. The residue was returned to the flask and redistilled for 30minutes with 50ml water to replace the alcohol will be evaporated adding hot water to replace the alcohol lost. When all the alcohol had been removed, 3 drops of 10% HCl was added. The whole solution was later transferred into a 250ml volumetric flask 5ml of zinc acetate solution and 5ml of Potassium Ferro cyanide solution was added thoroughly mixed to give homogenous solution.

The flask was allowed to stand for a few minutes filtered through a dry filter paper and 10ml of the filtrate was transferred into a separatory funnel and alkaloids present were extracted vigorously by shaking with five succession portions of chloroform. The residue obtained was dissolved in 10ml hot distilled water and transferred into a kjeldalh tube with the addition of 0.20g sucrose and 10ml Conc. H_2SO_4 and 0.02g selenium for digestion to colorless solution to determine %N by kjeldahl distillation method. % N got was converted to % total alkaloid by multiplying with a factor 3.26 i.e

$$\% \text{ total alkaloid} = \%N + 3.26$$

$$\% \text{ alkaloids} = \% N + 3.26.$$

2.5.2 Test for flavonoids

Exactly 0.50g of the extracted sample was weighed into a 100ml beaker and 80ml of 95% ethanol added and stirred with a glass rod to prevent lumping. The mixture was filtered through a whatman NO 1 filter paper into a 100ml volumetric flask and made up to mark with ethanol. 1ml of the extract was pipette into 500ml volumetric flask , four drops of conc. HCl added via a dropping pipette after which 0.5g of magnesium turning was added to develop a magenta red coloration . Standard solution were read on digitallabomed 200 spectrophotometer at a wavelength of 520nm . the percentage flavonoids was calculated using the formula

$$\% \text{ flavonoid} = \frac{\text{absorbance of sample} \times \text{average gradient} \times \text{dilution factor}}{\text{weight of sample} \times 10000}$$

2.5.3 Test for tannin

Exactly 0.20g extracted sample was measured into a 500ml of beaker , 20ml of 50% methanol was added and covered with paraffin and placed in water bath at 77-80°C for 1 hour . It was shaken thoroughly to ensure uniform mixing. The extract was quantitavely filtered using a double layered whatman NO. 41 filter paper into a 100ml volumetric flask, 20ml water added, 2.5ml folin-denis reagent and 10ml of 17% Na_2CO_3 were added and mixed properly . the mixture was made up to mark with water mixed well and and allow to stand for 20mins, the bluish-green color which developed at the end of range 0-10ppm were treated similarly as 1ml sample above .

The absorbance of the tannic acid standard solutions as well as samples were read after color development on a spectrophotometer at a wavelength of 70mm. %tannin was calculated using the formula:

$$\% \text{ tannin} = \frac{\text{absorbance of sample} + \text{average gradient factor} + \text{dilution factor}}{\text{Weight sample} \times 10000}$$

2.5.4 Test for saponin

1g of sample was weighed into a 250ml beaker and 10ml of butyl alcohol was added. The mixture was shaken on a UDY shaker for 5hrs to ensure uniform mixing. Thereafter the mixture was filtered through a whatman NO.1 filter paper into a 100ml beaker and 20ml of 40% saturated solution of magnesium carbonate was added. The mixture obtained with saturated MgCO_3 was again filtered through a whatman NO. 1 filter paper to obtain a clear colorless solution was pipette into a 500ml volumetric flask and 2ml of 5% FeCl_3 solution was added and made up to mark with distilled water. It was allowed to stand for 30 min for blood red color to development in a Jenway V6300 spectrophotometer at a wavelength of 380nm.

$$\% \text{ saponin} = \frac{\text{absorbance of sample} + \text{average gradient} + \text{dilution factor}}{\text{Weight sample} \times 10000}$$

2.5.5. Test for steroid

0.5g of sample extract was weighed into a 100ml beaker 20ml of chloroform-methanol (2:1) mixture was added to dissolve the extract upon shaking for 30mins

on a shaker . The whole mixture was filtered through a whatman NO. 1 filter paper into another dry clean 100ml conical flask /beaker.

The resultant residue was reportedly treated with chloroform- methanol mixture until it is free of steroids. 1ml of the filtrate was pipette into a 300ml test tube and 5ml of alcoholic KOH was added and shaken thoroughly to obtain a homogenous mixture. The mixture was later placed in a water bath set at 37°C- 40°C for 90mins. It was cooled to room temperature and 10ml of petroleum ether was added followed by the addition of 5ml of chard reagent to residue in dry bottle and absorbance taken at a wavelength of 620nm on spectronic 20D spectrophotometer. Standard steroids of concentration of 0.4mg/l were prepared from 100mg/ml stock steroid was calculated using the formular:

$$\frac{\text{absorbance of sample} + \text{average gradient} + \text{dilution factor}}{\text{Weight sample} \times 10000}$$

2.5.6 Test for glycosides

10ml of extract was pipetted into a 250ml conical flask. 50ml chloroform was added and shaken on a vortex mixer for 1hour, the mixture was filtered into 100ml conical flask and pyridine, 2ml of 2% sodium nitroprusside were added and shaken thoroughly for 10 minutes. 3ml of 20% NaOH was later added to give a brownish yellow color. The absorbance of the sample and the standards were read on a spectronic 20D digital spectrophotometer at a wavelength of 510nm. Percentage glycoside was calculated using the formula:

$$\frac{\text{absorbance of sample} + \text{average gradient} + \text{dilution factor}}{\text{Weight sample} \times 10000}$$

2.6 Antibacterial susceptibility test

2.6.1 Preparation of culture media

MacConkey agar of 24.00g and 14.00g of nutrient agar were weighed into two different with aluminum foil and shaken thoroughly and then left to disperse. They were then sterilized at 121°C for 15mins in autoclave. They were then poured into petri dishes and left to solidify. After which it was incubated for 24hrs at 37°C.

2.6.2 Isolation of organism

The inoculating loops were sterilized and were used to pick the samples. They were then used to streak the surface of each labeled petri-dishes containing the prepared media. The streaked media were then kept for the bacterial susceptibility test.

2.6.3 Bacterial susceptibility test

24hrs old standardized culture of bacterial was subcultured into distilled water and 1ml of the broth was used to flood the surface of the media prepared and allowed to dry. A sterile cork borer of 1cm was used to make holes and 1ml of the extract was dropped into each hole of the labeled dishes. Antibiotics were also placed on the surface of the media containing the cultured bacterial which were then incubated at 37°C for 24hrs, the minimum zone of inhibition of the extract and antibiotics were compared and records were taken.

3 RESULTS AND DISCUSSIONS

TABLE 1: DPPH free radical scavenging activity of fresh *Jatropha curcas* leaves

Parameter	Fresh leaves(mg/kg)	Ascorbic acid (mg/kg)
Antioxidant	74.73	31.01

Table 1 reveals the DPPH scavenging activity (indicator of the antioxidant) in fresh *Jatropha curcas* leaves in which the value was relatively high compare to the antioxidant ability of ascorbic acid a standard antioxidant which proves that the leaves have the ability of scavenging free radicals and the oxidation process in the body. These free radicals are responsible for some diseases and therefore *Jatropha curcas* has the ability to cure disease since it has the ability to inhibit the oxidation processes in human body that are responsible for these diseases.

TABLE 2: Qualitative and Quantitative analysis of butanol fraction of *Jatropha curcas* leaves.

Parameters	Qualitative	Quantitative (%)	Inference
Alkaloids	+	0.5670	present
Phlobatannins	-	0.0005	absent
Tannins	+	0.0018	present
Saponin	+	0.3160	present
Flavonoids	+	0.0024	present
Steroid	+	0.0037	present
Cardiac glycosides	+	0.1880	present
Phenol	+	0.0760	present

The qualitative and quantitative phytochemical analysis of the butanol extract of the leaves of *J. curcas* (Table 2), showed the presence of most phytochemicals like alkaloid, tannin, saponin, flavonoid, glycoside and phenol. This corroborates the findings of Thomas *et al* [10].

The medicinal value of this plant lies in its phytochemical constituents since each phytochemical is said to have a definite physiological and pharmacological action on the human body. Different classes of phytochemicals have been found to possess wide range of activities which helps in prevention and protection against diseases. The presence of alkaloid in the butanol extract could make the plant active against malaria, asthma and support its use as an analgesic.

Phytochemicals such as flavonoids, phenol are effective as antioxidant while saponin and glycosides are effective as anti-inflammatory and antibiotics respectively [11].

TABLE 3: Average value of the antibacterial activity of butanol extract of *Jatropha curcas* leaves.

Isolate	Zone of inhibition (mm)	
	Antibiotics	Butanol extract
Gram +ve		
<i>Staphylococcus sp.</i>	32(gen), 29(ofl)	30
<i>Bacillus cereus</i>	22(gen), 20(ofl)	20
<i>Clostridium sp.</i>	15 (gen), 20(ofl)	15
Gram -ve		
<i>Escherichia coli</i>	28(gen), 25(ofl)	41
<i>Proteus sp.</i>	20(ch), 20(ch)	17
<i>Pseudomonas sp</i>	40(cpx), 35(pef, ofl)	22

KEY

Cpx-ciprofloxacin

Pef-pefloxacin

Ofl-ofloxacin

301 Ch-chloraphenicol
302 Gen –gentamycin
303

304 Antibacterial activity of the extract (Table 3) revealed that the minimum zones of inhibition
305 (mm) values were not significantly different when compared with the antibiotics used
306 which shows that the leaves could be used as antibacterial agents with the exception
307 of *pseudomonas sp.* in which the its inhibitory zone value was relatively low to that
308 of the antibiotics used .

309 Igbinosa *et al.*,2009 opined that presence of biologically active principle (as confirmed in
310 table 2) may aid the antimicrobial activities of *J. curcas* as these secondary metabolite
311 exerts antimicrobial activity through different mechanisms.

312 This indicates that the plant extract is effective against most of the bacteria used in this
313 study. Therefore, the plant extracts has medicinal properties which supports its use in the
314 treatment of various forms of infections.

315 **4. CONCLUSION**

316 It can therefore be concluded from these research that *Jatropha curcas* leaves has
317 bioactive compounds (phytochemicals) as well as a high free radical scavenging activity
318 (indicator of antioxidant activity). Therefore, *Jatropha curcas* has great medicinal potential
319 and this could justify the use of its leaves in folklore therapy.

320

321

322

REFERENCES

1. Krippner S. Models of ethnomedicinal healing. Paper presented at the ethnomedicine conferences, Munich ,Germany. 2003, April 26-27 and October 11-12.
2. Abebe D: The role of ethnomedicinal plants in health care coverage of ethopia, the possible benefits of integration. *Journal of ethnobiologyand ethnomedicine*. 2001; 9:32.
3. Srivastava R. Studying the information needs of medicinal plant stakeholders in Europe. U.K. *Journal of applied and environmental microbiology*. 2000, (2):42-45
4. Rathee P, Hema C, Sushila R and Dharmender R. Mechanism of action of flavonoids as anti-inflammatory. *Journal of nutrient*. 2009, 2:(98).
5. Tiwari A. Imbalance in antioxidant defense and human diseases, *Journal of multiple approach of natural antioxidants therapy*. 2001, 81:1179-1187.
6. Sudha G, Sangeethapriya M, Indhushee R and Vadiuukkaras S. In vitro free radical scavenging activity of raw pepino fruit. *International Journal of pharamacy* 2011; (7) :137-140.
7. Stanner SA, Hugher J, Kelly CN and Butlriss JA. Review of the epidemiological evidence for the antioxidant hypothesis. *Journal public health nutrition*. 2000, 7: 401-422.
8. Dehpour AA, Ebrahimzadel MA, Nabari SF, Fazelin M and Eslami B. In vitro antioxidant and free radical scavenging activity of *Diospyros lotus* and *Pyrus boissieriana* growing in Iran. *Pharmacognosy magazine*, 2009, 4:123-127.

- 343 9. Aminov RI. A brief history of the antibiotic era. Lessons and challenges for the future.
344 *Journal of frontiers in microbiology* . 2010, 1:134.
- 345 10. Thomas T, Sah NK and Sharma PB. Therapeutic biology of *Jatropha curcas*: A mini
346 review. *Curr Pharm Biotechnol*; 2008, 9(4):24-315.
- 347 11. Edeoga H, Okwu DE and Mbaebre BO. Phytochemical constituents of some Nigerian
348 plants. *African journal of biotechnology* 2005, 44(7):685-688.
- 349 12. Igbinosa OO, Igbinosa EO and Aiyegoro OA. Antimicrobial activity and phytochemical
350 screening of stem bark extracts from *J. curcas* (linn). *African journal of pharmacy and*
351 *pharmacology* 2009, 3:58-62 rm.Bio. Sci. 2015: 6(2): (B)108-120.