

Isolation and Characterization of an isolated flavonoid from *Averrhoa bilimbi*

Abstract

Averrhoa bilimbi is a tropical plant which is commonly known as Bilimbi. The plant has an huge economic importance since its leaves, flower, fruit, bark, seeds, roots or the whole plant are used to treat variety of diseases in the alternative system of medicine. In the present work attempt was made to isolate a flavonoid compound from *Averrhoa bilimbi*. From the methanolic extract of the fruits of *Averrhoa Bilimbi*, a pentahydroxyl flavanonol has been isolated as a major compound for the first time in this plant. The isolate was purified, analyzed and characterized by using UV, FTIR, Mass, NMR, HPTLC and HPLC. The R_f value for HPTLC was found to be 0.24, λ max of UV spectra was obtained at 277 nm and retention time in HPLC was 2.55. The structure of this isolated compound has been characterized as dihydromyricetin i.e. '(2*R*,3*R*)-3,5,7-trihydroxy-2-(3,4,5-trihydroxyphenyl)-2,3-dihydrochromen-4-one' with molecular formula C₁₅H₁₂O₈ and molecular mass 320.0529. The structure is established on the basis of 1D and 2D Nuclear Magnetic Resonance (NMR) and also High Resolution Mass Spectral i.e HRMS data.

Key words: *Averrhoa bilimbi*, HPTLC, HPLC, UV, and FTIR

1. Introduction

Averrhoa bilimbi belonging to the family Oxalidaceae. It is a small tree which grows up to the height of 15 meters. Fruits are equally cylindrical with five wide rounded longitudinal lobes, and produced in groups. At the time of maturity there will be maximum increase in the weight of the fruits and its dimensions and also there will be change in color from green to light yellow. The fruits of *A. bilimbi* possess antibacterial, antiscorbutic, astringent and postpartum protective properties. The decoction of the leaves is being used as medicine for treating fever, inflammation of the rectum, diabetes, mumps and pimples. The paste of leaves is being used for the treatment of itches, boils, rheumatism, cough and syphilis. The juice of preserved fruits is being used for the treatment of scurvy, stomach ache, bilious colic, whooping cough, and hypertension. Moreover, the syrup of flowers is being given to treat children's cough. The plant is known for its antidiabetic, antihyperlipidemic and antibacterial activity [1,2]. The fruit extracts contain Saponins, Flavonoids, and Triterpenoid. The phyto constituents of *A. bilimbi* includes amino acids,

citric acid, cyanidin-3-O- β -D-glucoside, phenolics, potassium ion, vitamin A and sugars [3]. The present work has been carried out in the view to isolate flavonoids from the methanolic extract of the fruits of *Averrhoa Bilimbi*.

38

39 **2. Materials and Methods**

40

41 **2.1. Procurement of Plant Material**

42 The whole plant was collected from Palakkad, Kerala in South India. The specimen was
43 identified by Department of Pharmacognosy, Sanjo College of Pharmaceutical Studies,
44 Palakkad, Kerala. A voucher specimen was prepared in our laboratory and maintained
45 with voucher no. PPG/DPC/01, for further reference. Immediately after collection, the
46 fruit was washed thoroughly with water and then sliced, shade-dried at room temperature.
47 The sliced fruit was then pulverized to form a coarse powder and used for extraction.

48

49 **2.2. Preparation of fruit extract**

50 The sample of about 250 gm was taken in a thimble and kept it in a soxhlet apparatus. It
51 was consecutively extracted with 250 ml of each petroleum ether, hexane and methanol till
52 the extraction was complete. The methanolic extract was used for the screening and
53 isolation of compound [4].

54

55 **2.3. Preliminary phytochemical screening**

56 Qualitative phytochemical tests (colour reaction) were performed in methanolic extract to
57 find out the presence of important classes of phytochemicals such as alkaloids, flavonoids,
58 tannins, amino acids, phenolic compounds and triterpenes which were then further
59 confirmed by thin layer chromatography.

60

61 **2.4. Detection of phytoconstituents by TLC**

62 The spotted TLC plates were developed with the help of different mobile phases to find
63 out various groups phyto-constituents. The composition of the chemicals in the mobile
64 phases is shown in Table 1. Linear ascending development was carried out in 10 x 10 cm
65 twin trough glass chamber which was previously saturated with the mobile phase. The
66 chromatogram chamber was saturated with mobile phase for about 20 minutes at $25 \pm 2^\circ\text{C}$
67 with a relative humidity of $60 \pm 5\%$. Ten milliliters of the mobile phase was used for the
68 development of chromatogram. The mobile phase was allowed to migrate a distance of

about 85 mm from the place of sample application. After the development of chromatogram the TLC plate was dried and the chromatogram was viewed at 254 nm and 366 nm to visualize and to find out various phyto-constituents [5].

Table 1. Protocol for detection of various compounds by TLC

S. No.	Compounds	Mobile Phase	Visualizing Agent
1.	Alkaloids	Toluene : Methanol : Diethyl amine (8:1:1)	Dragendorff reagent
2.	Flavanoids	Toluene : Ethyl acetate : Formic acid (7:3:0.1)	NP/PEG Reagent
3.	Tannins	Ethyl acetate : Acetic acid : Ether : Hexane (4:2:2:2)	Fast Blue Salt B
4.	Triterpenes	Toluene : Chloroform : Ethanol (4:4:1)	Anisaldehyde sulphuric acid
5.	Amino acids	1-Butanol : Acetic acid : Water: Formic acid (28:9:8:2)	Ninhydrin
6.	Essential oil	Toluene : Ethyl acetate (8.5:1.5)	Anisaldehyde sulphuric acid

2.5. Screening of phytochemical groups using High Performance Thin Layer Chromatography (HPTLC)

HPTLC is ideal for the analysis of herbal drugs because of its flexibility, reliability and simplicity. A Camag HPTLC instrument comprises of automatic spotter (Linomat V) furnished with a 100 μ L syringe associated to a nitrogen cylinder, Scanner-III, twin-trough developing chambers, and UV viewing cabinet with dual wavelength UV lamps for short (254 nm) and long (366 nm) wavelengths (Camag, Switzerland) were used. HPTLC plates used were of aluminium backed silica gel 60 F₂₅₄ with 0.2mm thickness. HPTLC plates were washed by predevelopment solvent like methanol and activated at 110°C for 5min for complete solvent removal. Specific mobile phases were used for the development of each phyto-constituents [6-8].

2.5.1 Sample application

Sample was spotted on pre-coated TLC plate in the form of narrow bands (8 mm) with 10 mm from the lowest and at least 15 mm from left and right edges of the plate using Linomat V spotter. Samples were applied under incessant dry stream of nitrogen gas at constant application amount 10 μ L. HPTLC was performed using mobile phase Toluene:

Ethyl acetate: Formic acid (7:3:0.1) on precoated silica gel 60 F254 plates as stationary phase.

2.6 HPLC

Thermo HPLC system consisted of Quaternary gradient pumps (LC – 10ATvp); Photodiode Array (PDA) detector (SPD – M10Avp) with built-in system controller was used. The analysis was performed on a 250 x 4.6 mm, 5 µm particle size CNW, Athena C18-WP column. The data acquisition was done on ChromQuest 5 software. The isolated compound from HPTLC was analyzed by using Methanol: Water: Acetonitrile (40:40:20) as mobile phase and UV detector set at 254 nm. The injection volume was 20 µl, flow rate was 1 ml/min and run time was 10 minutes. The retention time of the fruit extract was compared with that of the isolated compound acting as the reference standard.

2.7.Isolation of flavonoid compound from methanolic extract

A specific phytochemical compound with R_f 0.24 was identified in the screening of flavonoid compounds and that compound was selected for further study. Extract was subjected to repetitive HPTLC using aluminium backed silica gel 60 F₂₅₄ as stationary phase (20 × 10 cm plates) and Toluene: Ethyl acetate: Formic acid (7:3:0.1 v/v/v) as mobile phase. A band under 254 nm at R_f value 0.24 was identified and were scraped. The compound was separated from silica gel by treating with methanol, filtered through Whatman filter paper, and filtrates were combined, concentrated, and dried. Isolated compound was subjected to HPTLC, HPLC, UV spectroscopy, IR spectroscopy, and LC MS.

2.8.Characterization of isolated compound[9]

2.8.1 UV Spectroscopy

The absorbance of the isolated compound was read using one cm cell in a UV – Vis - NIR spectrophotometer (Varian, Cray 5000, and Netherlands). The instrument have a spectral range of 175 nm to 3300 nm, wavelength accuracy of ± 0.1 nm (UV –Vis), ± 0.4 nm (NIR), Wavelength reproducibility of 0.025nm and a limiting resolution of 0.05nm(UV-Vis), 0.2nm(NIR).The maximum range of absorbance of isolated compound in the methanolic solution was noted by comparing it against HPLC grade methanol as a blank.

2.8.2. Fourier Transform Infra Red spectrometer (FTIR)

FTIR analysis was performed using Thermo Nicolet, Avatar 370 spectrophotometer system, which was used to detect the characteristic peaks and their functional groups. The Spectral range was between 4000-400 cm^{-1} and resolution was 4 cm^{-1} with KBr beam splitter, DTGS Detector and HATR Assembly for convenience of measurement. The finger print region extended between 400 – 1600 cm^{-1} . The spectrum of the isolated compound was elucidated against a blank of HPLC grade methanol.

2.8.3. LC-MS

The LC-MS system (Varian, USA-410 Prostar Binary LC with 500 MS IT) consisted of two pumps (LC-10ATvp), PDA detectors (SPD-M10Avp) and auto sampler (SIL-HTA) with built-in system controller. The analytical column was a C18, 250x4.6 mm ID, 5 μ particle size (Lichrospher 100 RP-18e, Merck, Germany) protected by a compatible guard column. For the characterization of isolated compound the HPLC method was same as that used in HPLC with CNW, Athena C18-WP column.

2.8.4 NMR/MS Studies

The 1D and 2D NMR spectral data were acquired using standard pulse sequences on Bruker Avance DRX 500 MHz or Varian INOVA 600 MHz instrument instruments. The NMR spectra were performed in $\text{C}_2\text{D}_6\text{SO}$ ($\text{d}_6\text{-DMSO}$). The chemical shifts were given in δ (ppm), and coupling constants were informed in Hz. MS and MS/MS data were produced with a Thermo LTQ-FTMS mass spectrometer (100,000 resolutions) fortified with a Nano spray ionization source. The samples were diluted with methanol and introduced via infusion using the onboard syringe pump.

3. Results and Discussion

3.1.Preliminary Phytochemical Screening

Methanolic extract was subjected to preliminary qualitative tests for the discovery of major phyto-constituents groups using standard protocols. The analysis have shown the presence of alkaloids, Flavanoids, Saponins, Triterpenoids, Tannins and Phenolic compounds. The results are presented in Table 2.

160

Table 2. Preliminary Phytochemical Studies of *Averrhoa bilimbi*

S. No.	Constituents	Test	Aqueous Extract	Methanolic Extract
1.	Alkaloids	Dragendroff's test	+	+
		Mayer's test	+	+
		Wagner's test	-	+
2.	Flavanoids	With 1% KOH	-	+
		With H ₂ SO ₄	+	+
		Legal's test	-	-
3.	Amino acids	Ninhydrin test	+	+
4.	Triterpenoids	Salkowski's test	+	+
		Libermann's Burchard test	-	+
5.	Tannins and Phenolic compounds	FeCl ₃ test	-	+

161

162 **3.2.Screening of phytochemical groups using HPTLC**

163 The results obtained from HPTLC analysis of the methanolic extract of *Averrhoa bilimbi*
 164 with respect to Alkaloids, Flavanoids, Tannins, Triterpenes, Amino acids and Essential oil
 165 are given below (Table 3).

166 Table 3. Rf Values of various phytoconstituents present in *Averrhoa bilimbi*

Sr. No.	Compounds	Rf Values
1.	Alkaloids	0.14, 0.45, 0.62, 0.91
2.	Flavanoids	0.24, 0.29, 0.62
3.	Tannins	0.74, 0.79
4.	Triterpenes	0.31, 0.37, 0.44, 0.47, 0.69, 0.89
5.	Amino acids	0.55, 0.64, 0.79, 0.86
6.	Essential oil	0.10, 0.12, 0.39, 0.53, 0.62, 0.68

167

168

169

170


171

172

173

174 **3.3.HPTLC of flavonoid compound**


175
176 Figure 1. HPTLC of the isolated flavanoid compound I



177
178
179
180 **3.4.HPLC of isolated compound**


181 The highest sharp peak with 2.55 retention time is of isolated compound I.

182 Figure 2 HPLC chromatogram of isolated compound I




3.5. UV spectrum of isolated compound

Figure 3. UV Spectrum of isolated compound I




3.6. IR spectrum of isolated compound

Figure 4. IR spectrum of isolated compound I



3.7.LCMS of isolated compound

Figure 5. LCMS of isolated compound



3.8 NMR Spectra:

Identification and spectroscopic data of dihydromyricetin (5, 7, 3', 4', 5'-pentahydroxyl flavanonol, 1) Off-White powder; ¹H-NMR (600 MHz, d₆-DMSO, δ ppm) and ¹³C-NMR (150 MHz, d₅pyridine/d₄-methanol/d₆-DMSO, δ ppm) spectroscopic data see Table 3a; HRMS (M+Na) + m/z 343.0426 (calcd. for C₁₅H₁₂O₈Na: 343.0424).




Fig. 6. Structure of dihydromyricetin

Table 3a. ¹H and ¹³C NMR Spectral data (chemical shifts and coupling constants) for dihydromyricetin

Position	NMR Data in d ₆ -DMSO	
	δH	δC
2	4.91 (1H, d, 12.6)	83.3
3	4.42 (1H, dd, 12.8, 6.4)	71.7
4		197.7
5		163.4
6	5.86 (1H, d, 2.4)	95.9
7		166.8
8	5.91 (1H, d, 2.1)	95.0
9		162.6
10		100.5
1'		127.2
2',6'	6.40 (2H, s)	106.9
3',5'		145.7
4'		133.5
3-OH	5.76 (1H, 6.2)	

Compound 1 was isolated in the form of an off-white powder. The molecular formula of the isolated compound has been deduced as C₁₅H₁₂O₈ from the adduct ion corresponding to [M+Na]⁺ ion observed at m/z 343.0426. This composition was further supported by the ¹³C NMR spectral data. The UV spectrum of compound 1 showed λ max at 277nm suggested a flavonoid structure. The ¹H NMR spectra data of compound 1 has been acquired in d₆-DMSO. The ¹H NMR spectra data of compound 1 showed doublet and doublet of doublets at δ 4.42 and 4.91 in d₆-DMSO corresponding to a proton each suggested the 2, 3-dihydroflavonol or 3-hydroxyflavanone skeleton in the structure of compound 1. The presence of 2,3 dihydroflavonol was further supported by the ¹³C NMR spectral data which showed the presence of oxymethine groups resonating between δ 71.7 and 85.8. In addition, the ¹H NMR spectra data of 1 also showed the presence of two meta-coupled aromatic protons as doublets between δ 5.86 and 6.50, and an additional two metacoupled aromatic protons δ 6.40 and 7.24 as singlets corresponds to a pentahydroxyl flavanonol scaffold. on the basis of COSY, HMQC and HMBC correlations the ¹H and ¹³C NMR values for all the protons and carbons for the compound 1 were assigned and are tabulated in Table 1. The HMBC correlations established the position of all the five hydroxyl groups at 5, 7, 3', 4', 5' positions as shown in Figure 6. The structure of compound 1 was determined unambiguously as dihydromyricetin (5, 7, 3', 4', 5'-pentahydroxyl flavanonol) on the basis of 1D and 2D NMR spectroscopic data [10].

Conclusion

Based on the UV, FTIR, LCMS and NMR analysis the isolated compound was found to be dihydromyricetin i.e (2*R*,3*R*)-3,5,7-trihydroxy-2-(3,4,5-trihydroxyphenyl)-2,3-dihydrochromen-4-one with molecular formula C₁₅H₁₂O₈ and molecular mass 320.0529. Further this compound can be explored to find out the mechanism behind its pharmacological activities like anitdiabetic, and antihyperlipidemic activites.

References

1. Anitha Roy, Geetha RV, Lakshmi T, Averrhoa bilimbi Linn–Nature's Drug Store- A Pharmacological Review. Int. J of Drug Develop & Res. 2011; 3(3):101-06.
2. Ambili S, Subramoniam A, Nagarajan NS, Studies on the Antihyperlipidemic properties of Averrhoa bilimbi fruit in rats. Department of Phytochemistry and

- Phytopharmacology, Tropical Botanic Garden and Research Institute, Palode, Thiruvananthapuram, Kerala State, India. *Planta Medica* 2009, 75(1):55-58.
3. Ashok Kumar K, Gousia SK, Anupama M, Naveena Lavanya Latha J. A review of phytochemical constituents and biological assays of *Averrhoa bilimbi*. 2013;3(4):136-39.
 4. Harbone J.B., *Methods of Plant Analysis Chapter II* In: *Phytochemical methods: A guide to modern techniques of plant analysis* Toppan Company Ltd, Japan, 1973, (1), pp 4 – 5.
 5. H. Wagner., S.Bladt, *Plant Drug Analysis, A Thin Layer Chromatography Atlas* , Second edition. Germany. 1996. 112-124.
 6. Eike Reich and Anne Schibli. *High – Performance Thin – Layer Chromatography for the Analysis of Medicinal Plants*. Theime Medical Publishers, Newyork. 2007. 110-150.
 7. Lukasz Ciesla and Monika Waksmundzka-Hajnos, Springer-Verlag, Berlin,Heidelberg., *Multidimensional and Multimodal Separations by HPTLC in Phytochemistry MM. Srivastava (ed.), High-Performance Thin-Layer Chromatography (HPTLC)*, 2011; 69-92. DOI 10.1007/978-3-642-14025-9_5.
 8. Kirti M. Kulkarni, Leena S. Patil, Mrs. Vineeta V. Khanvilkar, Dr. Vilasrao J. Kadam, “ *Fingerprinting Techniques in Herbal Standardization*” , *Indo American Journal of Pharmaceutical Research*, Vol 4, Issue 02,2014, ISSN NO: 2231-6876.
 9. Dr. Dheeraj Singh, Dr. Anjula Sachan, Dr. Hemant Singh, Dr. Ranjendra Nath, Dr. R. K. Dixit, “*Extraction, Isolation and Characterization of Phytochemicals*”, *WJPR* 2015; 4, (5), 2703-2717.
 10. Zhang YS, Zhang QY, Wang B, Li LY, Zhao YY. Chemical constituents from *Ampelopsis grossedentata*. *Journal of Chinese Pharmaceutical Sciences* 2006; 15:211-214.